



US 20080318225A1

(19) **United States**

(12) **Patent Application Publication**
Brentnall et al.

(10) **Pub. No.: US 2008/0318225 A1**

(43) **Pub. Date: Dec. 25, 2008**

(54) **COMPOSITIONS AND METHODS FOR
DETECTING CANCER**

(76) Inventors: **Teresa A. Brentnall**, Seattle, WA
(US); **Ru Chen**, Seattle, WA (US);
Katherine L. Pogue-Geile,
Pittsburgh, PA (US); **David C.**
Whitcomb, Allison Park, PA (US);
Mary Patricia Bronner, Moreland
Hills, OH (US); **Carol A. Otey**,
Chapel Hill, NC (US)

Correspondence Address:
BOZICEVIC, FIELD & FRANCIS LLP
1900 UNIVERSITY AVENUE, SUITE 200
EAST PALO ALTO, CA 94303 (US)

(21) Appl. No.: **11/840,112**

(22) Filed: **Aug. 16, 2007**

Related U.S. Application Data

(60) Provisional application No. 60/838,746, filed on Aug.
18, 2006.

Publication Classification

(51) **Int. Cl.**
C12Q 1/68 (2006.01)
G01N 33/53 (2006.01)
G01N 33/535 (2006.01)
(52) **U.S. Cl.** **435/6; 435/7.1; 435/7.92**

(57) **ABSTRACT**

The present invention provides methods and compositions involving detecting the presence of and/or assessing the risk of cancer in a subject. These methods include methods of detecting and diagnosing cancer in an individual; methods of identifying individuals at risk of developing a cancer; and methods of staging a cancer. The methods generally involve detecting a palladin gene nucleotide sequence alteration that has been found to be associated with cancer and/or detecting a level of a palladin mRNA and/or protein in a biological sample. The present invention further provides nucleic acid probes, nucleic acid primers, and antibodies, as well as kits comprising one or more of the same, for use in a subject method.

FIG. 1A

SEQ ID NO:1

```

1 cctgagtcac ccggcgggcg aggtataaag cccgatacct gccccgcgcc cgggtccgcg
61 agccccgtgc agctcccgtc cgctccggac gcggaatcgg gcagcagcgg gaggcggccc
121 ggagagccga gggaccctct gaagctccag caactccaga accaaatccg actggagcag
181 gaggccggcg ctccggcagcc tccgccagcc ccgcgcagcg cgcgcacctc gcccccttc
241 ccgccgcgcg ccgccttccc cgagctcgcg gcctgcacgc cgcgccgctc cccggagccc
301 atgagcgcgc tggcctcccg ctccgcccc gccatgcagt cctccggctc cttcaactac
361 gcgcgcccc aqcagttcat cgccgcgcag aacctcgggc ccgcgtcggg ccacggcacg
421 ccggcctcca gccccagctc gtcagcctc ccgtcgccta tgtcccgcag gccgaggcag
481 ttccggcgcg cccccgtgcc gcccttcgcg cagcccttcg gcgctgagcc cgaggccccg
541 tggggctcct cctgcgcgct gcccccgcc ccgccacccc cggctttcag ccccacggct
601 gccttcccgg tgcccgcagt gttcccactg ccgccgccac caccgccgct cccgagcccc
661 ggacaggcgt cccactgctc gtcgcctgcc acccgcttcg gccacggcca gacgcccgcg
721 gccttcctca gcgctctgct gccctcgcag ccgccgccgg cggccgtcaa cgccctgggg
781 ctgcccaagg gtgtcacccc cgcaggattt ccaaagaagg ccagtagaac tgctagaata
841 gcctccgatg aggaaattca aggcacaaa gatgctgtta ttcaagacct ggaacgaaaa
901 cttcgcttca aggaggacct cctgaacaat ggccagccga ggtaacata cgaagaaga
961 atggctcgtc gactgctagg tgctgacagt gcaactgtct ttaatttca ggagccagaa
1021 gaggaaacag ctaatcagga atacaaagtc tccagctgtg aacagagact catcagtgaa
1081 atagagtaca ggctagaaag gtctcctgtg gatgaatcag gtgatgaagt tcagtatgga
1141 gatgtgcctg tggaaaatgg aatggcacca ttttttgaga tgaagctgaa acattacaag
1201 atctttgagg gaatgccagt aactttcaca tgtagagtgg ctggaaatcc aaagccaaag
1261 atctattggt ttaaagatgg gaagcagatc tctccaaaga gtgatcacta caccattcaa
1321 agagatctcg atgggacctg ctccctccat accacagcct ccacctaga tgatgatggg
1381 aattatacaa ttatggctgc aaacctcag ggccgcacatc gttgtactgg acggctaagt
1441 gtacaggctg tcaaccaaag agtgcaagt ccccggtctc cctcaggcca cctcagctgc
1501 agaaggcctc gttctagatc aagggacagt ggagacgaaa atgaaccaat tcaggagcga
1561 ttcttcagac ctcaacttctt gcaggctcct ggagatctga ctgttcaaga aggaaaactc
1621 tgcagaatgg actgcaaagt cagtgggtta ccaaccccag atctaagctg gcaactagat
1681 ggaaagcccg tacgccctga cagtgtcac aagatgctgg tgctgagaa cgggggtgcac
1741 tctctgatca tagagccagt cacgtcacgt gatgccggca tctacacatg tatagctacc
1801 aaccgagcag gacagaactc attcagcctg gagcttgtgg ttgctgctaa agaagcacac
1861 aaacccccctg tgtttattga gaagctccaa aacacaggag ttgctgatgg gtaccagtg
1921 cggctggaat gtcgtgtatt gggagtgcga ccacctcaga tattttggaa gaaagaaaat
1981 gaatcactca ctacacagcac tgaccgagtg agcatgcacc aggacaacca cggctacatc
2041 tgccctgctc ttcagggagc cacaaaagaa gatgctgggt ggtatactgt gtcagccaag
2101 aatgaagcag ggattgtgtc ctgtactgcc aggctggacg tttacacca gtggcatcag
2161 cagtcacaga gcaccaagcc aaaaaagta cggccctcag ccagtcgcta tgcagcactt
2221 tcggaccagg gactagacat caaagcagcg tccaacctg aggccaacc atctcacctg
2281 aactgaata ctgccttggg agaaagtgag gacctgtaat ccagcattct tgtaaagct
2341 gaaacactga aacagccatt gccttgacca acatattcct ttgtcacatt atgtaaaag
2401 cagaaacata cctttgacta taagaaatta aaaaaaaca caaaataat atttttctta
2461 cttgatatac caaacttagt ttaagttagt aatgctaata caaatatac cattgcacag
2521 aaaatacaca tttactgtcc aatttaaac tttggaattg ctgtgattaa agtgatcaaa
2581 atgccaaaat actaaaggaa atcaattgtt cacaggtaac tacaatttgt attatctaca
2641 agtgccttta aacacaagat ataggtgctg tgtagcctga tagtgtgaaa tgtttaatga
2701 gggagtgtga ccacaaacag tactacaatg attctgaagc acagtgtatt cagacagata
2761 cagtgaacca agtgcaatat gtaaggatga aagaagaaga gatgacaaag aaatccaagt
2821 aaatgccttg tctttgcaaa tgtttttata ttaaatacata aggaaggaaac tacttgctt
2881 aaatgttaat atcaaaagag ttttctaaca aggttaatac cttagtctt aacattttt
2941 ttctttatgt gtagtgtttt catgctacct tggtaggaaa cttatttaca aaccatatta

```

FIG. 1B

3001 aaaggctaatt ttaaataataa ataataataaa gtgctctgaa taaagcagaa atatattaca
3061 gttcattcca cagaaagcat ccaaaccacc caaatgacca aggcatatat agtatttgga
3121 ggaatcaggg gtttggaagg agtagggagg agaatagaagg aaaatgcaac cagcatgatt
3181 atagtgtggt caattagata aaagtagaag gcacaggaga ggtagcaaag gccaggcttt
3241 tctttggttt tcttcaaaca taggtgaaaa aaacactgcc attcacaagt caaggaacc
3301 agggccagct ggaagtgtgg agcacacatg ctgtggagca cacatgctgt ggagattgca
3361 gtgtgtctga ggtttgtgta gtagtggaag attttaggta tgtagagcaa gttgaaaatg
3421 gattgagact gcatgggagg ataaatgaga aattgcctgt agcatctagt ctacttgaag
3481 gaagtggaga cataaggaga gacaaaaaca ggtttgtgcc ataaagtatt tttcaaaga
3541 caccaagatg tggtaaatga aaattattag ttcacttccc tgctgccatg aaactttgcc
3601 ttaagaaggt gctggattcc aaggtttgta aaggcatctc ggtaaagact gctttttgaa
3661 tgcataatgat tttgcatcag ctagactgag ttgattctga ccagacttga tggttttaag
3721 tcggaaccga taaattttaa aaaggagaaa aaataatttg acctagtagt ataaaacatg
3781 aggttttaat ggtactttgc tatgaaaaga aaacactgta ttccttatgc aaaacacatg
3841 tatctttcat tatttataag tggcctctct tagctcagtt actcaattca tacgtagtat
3901 tttttaaaat aattttatat ctgtgtacca ccccatatat ttcataattac tgtttcacat
3961 gtacagcttt ctacttcttt gtaagaacac caaccaacca aggtttaagt gattaatagg
4021 cttgagcacc ggggtggcaga tgttctatgc agtgtgggtc aagtttcttt gaccgcactt
4081 atatgcattg ctaatatgga atttaagata ccatacacag tctctcatgg acctatctct
4141 attgtagaat tatgacttat gtcttacttg gcaaattttt ctgaatgtga cctttttttg
4201 ctgatttgct gggtttgga ttaactagca ttattttgcc acctttatat tgtatttata
4261 aaaaaaaagt actatcaatc aatcatacta ctttggttgg ttgtgctggt gtaatgtgga
4321 tttaacatca ataaatattt gacaaat

FIG. 2




FIG. 3A




FIG. 3C




FIG. 3B

Gene Name	Sporadic	Family X
1. KIAA0992: palladin IMAGE:2542582	21.3	14.564
2. KIAA0992 protein IMAGE:469720	13.021	2.357

FIG. 4

Chromosome 4q31-




FIG. 5

Human	S	Q	T	P	A	A	F
Chimp	S	Q	T	P	A	A	F
Mouse	S	Q	T	P	A	A	F
Rat	S	Q	T	P	A	A	F
Chicken	G	Q	S	P	A	A	F
Zebra Fish	C	Q	T	P	P	A	P
Family X	S	Q	T	S	A	A	F

FIG. 6




FIG. 7

Palladin Gene Expression in Epithelial Ductal cells




FIG. 8A




FIG. 8B




FIG. 10

621 ggacaggcgt cccactgctc gtcgctgcc acccgcttcg gccacggcca gacglccgcg
gccttcctca gcgctctgct gccctcgag ccgccgccgg cggccgtcaa cgccctgggg 780
(SEQ ID NO:63)

Forward primer: 5'-ggacaggcgtccactgctc-3' (SEQ ID NO:64)

Reverse primer: 5'-cccaggcggttgacggccg-3' (SEQ ID NO:65)

FIG. 11

BAA76836 palladin protein

PESPGGRGIKPDTC PAPGPRSP LQLPLAPDAESGSSGRRPGEPR
DPLKLOQLQNQIRLEQEAGARQPPAPRSAPPSPFFPPPAFPELAACTPPASPEPMS
ALASRSAPAMQSSGSFNYPKQFIAAQN LGPASGHGTPASSPSSSSLPSPMSPTPRQ
FGRAPVPPFAQPF GAEP EAPWGSSSPSPPPPPPPVFSPTAAFPVPDVFPLPPPPPLP
SPGQASHC SSPATRFHG **QT PAA**FLSALLPSQPPAAVNALGLPKGVTPAGFPKKASR
TARIASDEEI QGTKDAVI QDLERKLRFKEDLLNNGQPRLT YEERMARRLLGADSATVF
NIQEPEEETANQEYKVSSCEQRLI SEIEYRLERSPVDESGDEVQYGDVPVENGMAPPF
EMKCLKHYKIFEGMPVTFTCRVAGNPKPKIYWFKDGKQISP KSDHYTIQRDL DGTC SLH
TTASTLDDDGN YTIMAANPQGRISCTGRLMVQAVNQRGRSPRSPSGHPHVRRPRRSR
DSGDENEPIQERFFRPHFLQAPGDLTVQEGKLCRMDCKVSGLPTPD **LSWQLD**GKPVRP
DSAHKMLVRENGVHSLIIEPVT SRDAGIYTCIATNRAGQNSFSLELVVAAKEAHKPPV
FIEKLQNTGVADGYPVRLECRVLGVPPPQIFWKKENESLTHSTDRVSMHQDNHGYICL
LIQGATKEDAGWYTVSAKNEAGIVSCTARLDVYTQWHQSQSTKPKKVRPSASRYAAL
SDQGLDIKAAFQPEANPSHLTLNTALVESEDL (SEQ ID NO:99)

FIG. 12

NP_057165.3 palladin amino acid sequence

```

1 msgtsshesf ydsldsmdqee skntdffpgl saflsqeein ksldlarrai adsetedfds
61 ekeisqifst spaslcehps hketklgeha srrpqdnrst pvqplaekqt ksisppvskr
121 kpamsp11tr psyirslrka ekrqaktpst nvkpktphqr kggpqsq1cd kaanlieelt
181 sifkaakprn rspngessp dslylspknq psallsasas qspmedqgem erevkspgar
241 hcyqdnq1da vphnrkshpq phsalhfpaa prfiqklrsq evaegsrvyl ecrvtgnptp
301 rvrwfcegke lhntpdqiqh ceggdlhtli iaefeddtg rytclatnps gsdttsaevf
361 iegasstdsd seslafksra gampqaqkkt tsvsltigss spktgvttav iqplsvpvqq
421 vhsptsylcr pdgtttayfp pvftkelqnt avaegqv1v1 ecrvrgappl qvqwfirgse
481 iqdspdfri1 qkkprstaep eeictlviae tfpedagift csarndygsa tstaqlv1ts
541 antencsy1e mgesnndhfq hfpppppile tsslelaskk pseiqqv1np elgl1sraalq
601 mqfn1aaeret ngvhpsrgvn glingkansn kslptpav1l sptkeppp1l akpklgfpkk
661 asrtari1asd eeiqg1tkdav iqdlerklrf kedllnnggp rlyteermar rllgadsatv
721 fniqepee1et anqeykvssc eqrliseiey r1erspvdes gdevqyg1dvp vengmapffe
781 mklkhyk1ife gmpvtftcrv agnpkpk1iyw fkdgkqispk sdhytiqr1dl dgtcslhtta
841 stldddg1nyt imaanpqgri sctgrlmvqa vnqrgrs1prs psghphv1rrp rsrsrdsgde
901 nepiqer1ffr phflqapgd1 tvqegklcrm dckvsgl1ptp dlswq1ldgkp vrpdsahkml
961 v1rengvhsli iepvtsrdag iytciatnra gqnsfsle1lv vaakeahkpp vfieklqntg
1021 vadgypvr1le crvlgvpppq ifwkkenes1l thstdrvsmh qdnhgyic1l1 iggatkedag
1081 wytvsak1nea giv1sctarld vyisrh (SEQ ID NO:87)

```

FIG. 13A

NM_016081
 Palladin mRNA

```

1  gtgaccacgg accaggcagt ctctaatagaa taggcaaggc cacaacctcc atttctcccag
61  aaaagaagaa atgctcatct gaaattcatc acctctctgg agtcttcaaa ctgaccaagc
121 attgaaaaga acacagtttc agaaaacagt ttccagtgcc tctggccttc ctactgaaag
181 cagacacaga gtgcatgaag accgttcaaa tatgtcaggg acctcctccc atgagtcctt
241 ctatgactcc ctctcagaca tgcaggaaga aagcaagaat actgacttct tcccgggcct
301 ttctgctttc ctcagccagg aagagataaa caagagcttt gacctggccc ggagagccat
361 agccgactcc gaaacagaag attttgactc ggaaaaggag atctcgcaga ttttcagtac
421 ttctcctgca agcctctgtg aacatccttc ccataaggag accaaattgg gtgaacacgc
481 ctcgaggaga cctcaggata acaggtcaac acctgtccag cctctggcag agaacaacaa
541 taagagtatc tcttcacctg tttcaaagag gaaacctgcc atgtcacccc tgctcaccag
601 gcccagctac atccggagcc tccgaaaggg tgaaaagcgt ggtgcaaaa ctcccagcac
661 aaacgtaaag cccaaaacgc cacatcaaag aaagggtggc ccccagagcc agctgtgtga
721 caaggcagct aatttaattg aggagctaac atccatattt aaagccgcaa agcaagaata
781 cagaagccca aatggggagt cctcgtcacc agacagtggg tacctgtctc ttaaaaatca
841 gccgtcagcc ctgctgagtg cctcagccag ccagagccct atggaagacc aaggggagat
901 ggaaagagag gtcaagtccc ctggggccag gcattgctac caggacaacc aggacttggc
961 agtgccacac aaccgcaagt ctcacccaca gcccacagc gccctccact tcccagctgc
1021 acctcgattc atccaaaagc tgaggagcca agaagtagca gaaggagcc gagtttatct
1081 ggagtgtaga gtcactggaa accccactcc tcgagtcaga tggttctgtg aagggaaaga
1141 actgcacaac actcctgata ttcaaatacca ctgtgagggg ggggacctcc ataccctgat
1201 catagcagag gcctttgagg acgacacagg tcgctacacc tgtttggcta cgaatccag
1261 cggctcagac acaacatctg ctgaggtggt cattgaaggt gccagtcaa cagattctga
1321 cagtgaaggt ttagctttca aatcaagagc tggagctatg ccacaagctc aaaagaaaac
1381 aacttctggt tccttgacaa taggatcatc atctcaaag acaggggtga ccacagctgt
1441 gattcaacca ctgtctgtcc ctgtgcaaca ggtcacagc ccaactcat atctctgccg
1501 acctgatgga accactactg cctactttcc tcctgttttt acaaaggaac tgcaaaacac
1561 agccgtggcg gaaggccagg tggtggttct ggagtgcggg gtccgtgggg caccctctct
1621 gcaggtccag tggtttcggc aagggagtga aatccaagac tctccagatt tccgaattct
1681 acagaaaaaa cctagatcta cagctgaacc tgaggagatt tgcaccctag ttatcgctga
1741 gactttccct gaagatgcag ggatctttac atgttcagca agaaatgatt atggatcagc
1801 aaccagcact gccagctgg ttgtcacctc agccaacact gaaaactgta gttacgagtc
1861 aatgggagaa tccaacaatg accacttcca acactttcca cctcccctc caatcttggg
1921 gacaagttcc ttggagttgg cttcaaagaa accatctgag atccagcagg tgaacaacc
1981 tgagttagcc ctgagcaggg cagcccttca aatgcaattc aatgtgctg agagggaaac
2041 gaacggagtc catcccagcc gtggagtaaa tggactgatt aacggcaaag ctaacagtaa
2101 taaatctctt ccaacaccag ctgtcctgct ttcaccact aaggagccac cacctctgct
2161 tgccaaacca aaactaggat ttcaaagaa ggccagtaga actgctagaa tagcctccga
2221 tgaggaaatt caaggcacia aggatgctgt tattcaagac ctggaacgaa aacttcgctt
2281 caaggaggac ctctgaaca atggccagcc gaggttaaca tacgaagaaa gaatggctcg
2341 tcgactgcta ggtgctgaca gtgcaactgt ctttaatatt caggagccag aagaggaaac
2401 agctaatcag gaatacaaa tctccagctg tgaacagaga ctcatcagtg aaatagagta
2461 caggctagaa aggtctcctg tggatgaatc aggtgatgaa gttcagtatg gagatgtgcc
2521 tgtgaaaaat ggaatggcac cattctttga gatgaagctg aaacattaca agatctttga
2581 gggaaatgcca gtaactttca catgtagagt ggtggaaat ccaaagccaa agatctattg
2641 gtttaaaagt gggaaagcaga tctctccaaa gactgatcac tacaccattc aaagagatct
2701 cgatgggacc tgctccctcc ataccacagc ctccacccta gatgatgatg ggaattatac
2761 aattatggct gcaaaccctc agggccgcat cagttgtact ggacggctaa tggtagagcc
2821 tgtcaaccaa agaggtcgaa gtccccggtc tcctcaggc catcctcatg tcagaaggcc
2881 tcgttctaga tcaagggaca gtggagacga aatgaacca attcaggagc gattcttcag
2941 acctcacttc ttgcaggctc ctggagatct gactgttcaa gaaggaaac tctgcagaat
3001 ggactgcaaa gtcagtgggt taccaccccc agatctaagc tggcaactag atggaagcc
3061 cgtacgccct gacagtgtct acaagatgct ggtgcgtgag aacggggtgc actctctgat
3121 catagagcca gtcacgtcac gtgatgccgg catctacaca tgtatagcta ccaaccgagc
3181 aggacagAAC tcattcagcc tggagcttgt ggtgctgct aaagaagcac acaaaccccc
3241 tgtgtttatt gagaagctcc aaaacacagg agttgctgat gggtagccag tgcggctgga
3301 atgtcgtgta ttgggagtgC caccacctca gataattttg aagaaagaaa atgaatcact
    
```


FIG. 13B

```

3361 cactcacagc actgaccgag tgagcatgca ccaggacaac cacggctaca tctgcctgct
3421 cattcagggg gccacaaaag aagatgctgg gtggtatact gtgtoagcca agaatgaagc
3481 agggattgtg tcctgtactg ccaggctgga cgtttacatt tctcgacatt aatagtgaac
3541 cacaccagga gaacaaatac ccaaccaggt ggcatcagca gtcacagagc accaagccaa
3601 aaaaagtacg gccctcagcc agtcgctatg cagcactttc ggaccagggg ctagacatca
3661 aagcagcgtt ccaacctgag gccaacccat ctcacctgac actgaatact gccttggtag
3721 aaagtgagga cctgtaatcc agcattcctg ttaaagctga aacctgaaa cagccattgc
3781 cttgaccaac atattccttt gtcacattat gtaaaaggca gaaacatacc tttgactata
3841 agaaattaaa aaaaaaacac caaaataata ttttcttac ttgatatacc aaacttagtt
3901 taagtagata atgctaatac aaatatacac attgcacaga aaatacacat ttactgtcca
3961 atttaaaact ttggaattgc tgtgattaaa gtgatcaaaa tgccaaaata ctaaaggaaa
4021 tcaattgttc acaggtaact acaatttgta ttatctacaa gtgcctttaa acacaagata
4081 taggtgctgt gtagcctgat agtgtgaaat gtttaatgag ggagttgtac cacaaacagt
4141 actacaatga ttctgaagca cagtgtattc agacagatac agtgaaccaa gtgcaatatg
4201 taaggatgaa agaagaagag atgacaaaga aatccaagta aatgccttgt ctttgcaaat
4261 gtttttatat taaatcataa ggaaggaact acttgcctta aatgttaata tcaaaagagt
4321 tttctaacaa ggtaataacc ttagtcttta acattttttt tctttatgtg tagtgttttc
4381 atgctacctt ggtaggaaac ttattttacaa accatattaa aaggctaatt taaatataaa
4441 taatataaag tgctctgaat aaagcagaaa tatattacag ttcattoac agaaagcatc
4501 caaaccaccc aaatgaccaa ggcatatata gtatttgag gaatcagggg tttggaagga
4561 gtagggagga gaatgaagga aaatgcaacc agcatgatta tagtgtgttc atttagataa
4621 aagtaaaagg cacaggagag gtagcaaagg ccaggctttt ctttggtttt cttcaaacat
4681 aggtgaaaaa aacactgcca ttcacaagtc aaggaacca gggccagctg gaagtgtgga
4741 gcacacatgc tgtggagcac acatgctgtg gagattgcag tgtgtctgag gtttgtgtag
4801 tagtggaaaga ttttaggtat gtagagcaag ttgaaaatgg attgagactg catgggtggca
4861 taaatgagaa attgcctgta gcatctagtc tacttgaagg aagtggagac ataaggagag
4921 acaaaaacag gtttgtgcca taaagtatth tttcaaagac accaagatgt ggtaaatgaa
4981 aattattagt tcacttccct gctgccatga aactttgcct taagaagggtg ctggattcca
5041 aggtttgtaa aggcatctcg gtaaagactg ctttttgaat gcatatgatt ttgcatcagc
5101 tagactgagt tgattctgac cagacttgat ggttttaagt cggaacccat aaatttttaa
5161 aaggagaaaa aataatttga cctagtagta taaaacatga ggctttaatg gtactttgct
5221 atgaaaagaa aacactgtat tccttatgca aaacacatgt atctttcatt atttataagt
5281 gccctctctt agctcagtta ctcaattcat acgtagtatt ttttaaata attttatatc
5341 tgtgtaccac cccatatatt tcatattact gtttcacatg tacagcttct tacttctttg
5401 taagaacacc aaccaaccaa ggtttaagtg attaatagcc ttgagcaccg ggtggcagat
5461 gttctatgca gtgtggttca agtttctttg accgcactta tatgcattgc taatatggaa
5521 tttaagatac catacacagt ctctcatgga cctatctcta ttgtagaatt atgacttatg
5581 tcttacttgc caaatttttc tgaatgtgac ctttttttgc tgatttgctg ggtttgggat
5641 taactagcat tattttgcca cctttatatt gtatttataa aaaaaaaagt actatcaatc
5701 aatcactacta ctttggattg ttgtgctggt gtaatgtgga tttaacatca ataaatattt
5761 gacaaaataat agttgcagtt ttgtgaagca aaataaatat tcagtttta (SEQ ID NO:20)

```

FIG. 14

GenBank AC080188; Homo sapiens BAC clone RP11-635L1

183421 agccacctct tgtactactg aaggaggaat ttatgcagac ttcttagcta ccagtgattt
183481 cactctgttt taatacaaaa atttacatgt atttctttta tgatttaggt cagtggtta
183541 ccaacccag atctaagctg gcaactagat ggaagcccg tacgccctga cagtgcacac
183601 aagatgctgg tgcgtgagaa cggggtgcac tctctgatca tagagccagt cacgtcacgt
183661 gatgcggca tctacacatg tatagtacc aaccgagcag gacagaacto attcagcctg
183721 gagcttgtgg ttgctggtag gctcatctgt gaatccttgc tctctgacag aatgaacatc
183781 agacttacia atgtaacta attctacatt actaaccaat acggaaaata (SEQ ID NO:101)

FIG. 15

GenBank AC084353; Homo sapiens BAC clone RP11-592K15

26881 tgcgtcagat gagagcagca caatcacctc ttctttaaca acttcacaca acacagggat
26941 tctcagaaga ctctgacagt gtgaaatcac ttggtgaact agtggcatct tcttatgttt
27001 ttcctctctt tcccctcct tagccaacac tgaaaactgt agttacgagt caatgggaga
27061 atccaacaat gaccacttcc aacactttcc acctccccct ccaatcttgg agacaagtcc
27121 cttggagtty gcttcaaaga aaccatctga gatccagcag gtgaacaacc ctgagttagg
27181 cctgagcagg gcagcccttc aaatgcaatt caatgctgct gagagggaaa cgaacggagt
27241 ccattcccagc cgtggagtaa atggactgat taacggcaaa gctaacagta ataaatctct
27301 tccaacacca gctgtcctgc tttcacccac taaggagcca ccacctctgc ttgccaaacc (SEQ ID NO:102)

FIG. 16

GenBank AC080188; Homo sapiens BAC clone RP11-635L1

```

139501  tgagagtgaa atgggggagc atacttcaact gacttagcgc ttggiccaga atgaactgct
139561  gggcccacac ttgggacctgc taagtggtgc ccgttccctg ggtgttctct gcagctggcc
139621  taagtcccaa gcgatgtcca caagctgagc tcaactcctgg aatacacgtt cctggccggt
139681  ccatctctga aggtgtcaact tctctttttc cccccaggga cctctctgaag ctccagcaac
139741  tccagaacca aatccgactg gaggaggagg cgggctctcg gcagcctcag ccagccccgc
139801  gcagcgcgcc gccctcgccc cccttcccgc cggcgcgcgc cttecccag ctgcgggect
139861  gcacgcgcgc cgggtccccc gagcccata ggcgctggc ctcccgcctc gcccccgcca
139921  tgcagtcctc cggctccttc aactacgagc gcccaagca gttcatagcc ggcagaaacc
139981  tcgggcccgc gtcgggcccac ggcacgcggg cctccagccc cagctcgtcc agcctcccgt
140041  cgcccattgc cccgacgcgc aggcagttcg gcggcgcgc cgtgcgcgcc ttggcgcagc
140101  ccttcggcgc tgagcccag gcccctggtg gctcctcctc gccgtgcgcc ccgccccccg
140161  cacccccggc cttcagcccc agggctgcct tcccggtgcc cgaagtgttc ccaactggcc
140221  cgccaccacc gccgctcccc agcccgggac aggcgtccca ctgctcgtcg cctgcccacc
140281  gcttcggcca cagccagagc gccgcggcct tctcagcgc tctgctgccc tcgcagcccg
140341  cgcggcgggc cgtcaaacgc ctggggctgc ccaagggtgt cacccccggc tgagtaaccg
140401  ccgcgggtcct ccacttccct gccctcctgc ctccgggtgc cctgggactc ccacatctcc
140461  atacacgcgc tcccctcagc ctgcaaccca gaggcacca gtaacatttc acacatttct
140521  ctccgtgcga tgtaaaaatt cttaacggca atttgactca gtgattcttg cgtagccact

```

(SEQ ID NO:100)

FIG. 17




FIG. 18




FIG. 19

AAATGCTTTACTCTTCTCACCCCTAAGGGCTCCTGAATTTAGCTTTTTATTACACAGAACC
CAGTAAACTAATTAATGGAAATCTAAAAATATCCTTCCTCTTCTTAGAAAAGAAATATA
CTGCATGATGATTTTGTGCTTTTTCGATTTAATGTACACACAATCGTTTTTACACTGTATGTG
CTTCCCTCCCCACAGCCCCAGTCCCAGCCCCAGTCTCGCAGTTCCATTCCCTTCAGGGCA
GCTTCCATTCTACCGTGAGTTTTTATTGCTTTCCCCTAAAACGCTGCCCTGGAGTCGCGT
TTATTTGGACACTCTTTAGTCTGTAAAGGAAAAAACACAAACACCCGGAGGAGAATCATCC
TAACAAGCCGACAACATATCTATGGGCCTAAAAAACTTAAATTCCTTTCCCGTTATTAGCA
CTTCTCGATCGTTCCAGGAAAGCCCCGATTTGTAGCTTGGCTGCCCCAGGAAGGGGCTGC
GACCAGAAAGATGCAAAGGGGGGACAAAGGTTACGAGTGGGTGCCGGGAAGCCTCCCTAA
CCTGGGGGCGCGCTCCCAGAGCTGCGAGCCACGCCCTCCTCTCTCCCGCCCGCGCGCCGG
CGGACTCTTATTTTTGAAGGGCGCGGGTGAAGGCTCGGAGCCTCCTGAGTCACCCGG
CGGGCGAGGTATAAAGCCCGATACCTGCCCGCGCCCGGTCCGCGGAGCCCCGCTGCAG
CTCCCGCTCGCTCCGGACCGGAATCGGGCAGCAGCGGGAGGCGGCCCGGAGAGCCGAGG
TAGGCGCGGGGAATCGGCCCTGAGGCTGGTGAAGAAATGTGTGAATTAAGGGAGTGCA
GCGTGCAGGGTGGGCGCGAGTCCGGAGTGCAGGGCTCGGGACAGGGTGGGGGCGGGGAG
GACAAGAGACTCGTTCAGCGCACTTCCACCCGGCCGGTTCCTTCCCGAGCCCTCTCCCTC
CCCCGGCCCGCGGGAGCACAGATCCCTGAAAAGCGGAGCTCCAGATAGGAATGTCTGCATC
ATCTTGGCAGGGCCAGACGGGAGAAGATGCTCTGTGTCACGACGATCCCTCCCGCTC
TCCTAACTCTTCTCTGAACACCTCGCACCCCCACGGCCCCCGCCCCCTCTCTTAATCTTG
GCCACGCTCCCTGAGGGGTGAGACCTGGGGTGGGAGAGCAGAGAAAGCGGGGCTGGCAGA
GACAGGGGTGAGGGCCGGGAGAGCGATGACATCATTCCTTGGCAGCTGGCGGGGAGAGG
GTGGGGACGGAGGGGTCCCCCGACCTGAGCCGAGCCCTCCCTTCCCAAGCCCATCTCC
TTTCTTTGGGATGGTTCACCCACCTGGGGCCCATTCAAAGGAAGTTTGGCGTAACTCG
GGGCGTCTTTCCCCGGCCGGGTGCTGTGCCCGGCCCGCAGTCCGTGTTTCTTCTGGGCC
TGGGGGCGCGTGGAGCCGGCGGCTCAGTCCCTCAGTCCCAGGCCTTGGCAGCCGCTTT
GTTACGCCGGGCGCGGGCCGAACCCGAACCTCAGCTGCAGCACTGGCAGCGCGCACCCGG
CCACGCGTGGGTTCCCCGGCCACGGCCGGGCCAACAAAGGGAGGGTCCCGTGAGCAGCCAGG
CTGGCCCTCCCCGGGCTGGGTGAGCTCCCAAAGCTGGAGCGCAGGGCTCTAGGCCGGCC
CGCCCCGGGGCTGGGGCTTCCCGCCCGTTTGTCTCATTTTAACTTTTGGGGCAGCTC
TGTTCCGAATGGTGAGTGGTCCAGACTCTAGAAAGGGTTTGGTCCCTTGGCCGGGATT
GAGGCTTGGACCTTTGCTCTCTTGTGGGACGGAAGGGGGCGGCTGGGCAGGAGAGG
GGCGTCTCGSTCGGTCTCCCTGGGCTCGGGCATCTTCCACCCCTGCGGAGCCGGGTCT
TGCGCTGCGTGCCCCGAACAGGCCCGGGCACACCGAGTCCCGTTACTGGTCTTCAGGCGA
AATTCAGGACGGGTGTGCATTCATCAACCTGGGAGTCTCTAGGAGCCAGAGATGACTGG
CTCTACATTTAATGCGTAATCACAGAGAACTAGCACGAGAAAAGGACAGAGGGACCGA
CCCAAGCTTACAGAGACATGGAATAATATGATCCTGGTTCATAAGCCTTGAAAAGCTCCCC
TCCGCCAGGTTAAGGAATGGTGTACCAGCCTTGGCTGTGAGGATAATTTCTTGATTTGT
GTTATGCAGGCTCCTGTACAGGACTGTCTCCTTAACTCCCACTACCTAAGGAAAAGATAGC
CAGAGAATTTGCCAGGGATTTGGGTTTGGGTTGCTTCTAGAAAGACAGCATCTTCTTGC
TATTTTTTTTTTTCACATCCCCCAAAACAATTTTTTTTCCCCCTTCAGACTTGGCAGACTT
TGATTTAGATAAGAGGTTTGGGTTAATTTGATTTTCCCCATGTCTGTTTAGATCAATGGA
TATAGGAGATAGTTTGAAGCTAATGTACCCCTATGCCTTTGTGTGAGTGTAAAGATTTATA
ACTTGAGGGAGGCAGTTAATAAATAGCACATCCTTGAACCATAAACATACGAGATAGACA
AGAGCTAGATAATAAACAGTTTTTCCCCCAATCCCTAAGCTTTACTTATTTTTTTTACTG
ATTCTTGTACTTTGTCAATAGAAAAGGAGTGGGGTTGGTTTCCCAGTGACCGCATTCAT
(SEQ ID NO:110)

COMPOSITIONS AND METHODS FOR DETECTING CANCER

CROSS-REFERENCE

[0001] This application claims the benefit of U.S. Provisional Patent Application No. 60/838,746, filed Aug. 18, 2006, which application is incorporated herein by reference in its entirety.

BACKGROUND

[0002] Pancreatic cancer is the fourth leading cause of cancer death in the United States. It is difficult to detect, early to metastasize and resistant to treatment. Nearly every person diagnosed with pancreatic cancer will die from it, usually within 12 months of diagnosis. Familial clustering of pancreatic cancers is commonly recognized, occurring in at least 10% of all pancreatic cancer. The risk of pancreatic cancer increases further with each family member who is affected.

[0003] Current methods for diagnosing pancreatic cancer include computed tomography scanning, magnetic resonance imaging, positron emission tomography scanning, endoscopic ultrasonography, laparoscopy, endoscopic retrograde cholangiopancreatography, percutaneous transhepatic cholangiography, and biopsy.

[0004] Early detection and diagnosis are important for the successful treatment of cancer in general. Despite advances in detection of various cancers, there is an ongoing need for diagnostic methods and tools for detection of cancers. In particular, there is a need in the art for methods of detecting pancreatic cancer.

LITERATURE

[0005] U.S. Pat. No. 6,642,009; Parast and Otey (2000) *J. Cell Biol.* 150:643-655; Mykkänen et al. (2001) *Molec. Biol. Cell* 12:3060-3073; Boukhelifa et al. (2001) *Molec. Biol. Cell* 12:2721-2729; Boring et al. (1994) *CA Cancer J Clin* 44:7-26; Hruban et al. (1999) *Annals of Oncology* 10 Suppl 4:69-73; Silverman et al. (1999) *British Journal of Cancer* 80:1830-7; Eberle et al. (2002) *Am J Hum Genet.* 70:1044-8; Ghadirian et al. (1991) *Int J Pancreatol* 10:183-96; Lynch et al. (1996) *Semin Oncol.* 23:251-75; Pohl et al. (2005) *Cancer Res.* 65:1994; Rachlin and Otey (2006) *J. Cell Sci.* 119:995-1004; WO 04/016809; WO 03/025138; WO 02/059377; U.S. Patent Publication No. 2004/0029114; WO 06/017635; WO 05/094306; WO 06/002433; U.S. Patent Publication No. 2006/0024692; WO 04/031413; WO 04/031410; U.S. Patent Publication No. 2004/0076955; WO 03/039443.

SUMMARY OF THE INVENTION

[0006] The present invention provides methods and compositions involving detecting the presence of and/or assessing the risk of cancer in a subject. These methods include methods of detecting and diagnosing cancer in an individual; methods of identifying individuals at risk of developing a cancer; and methods of staging a cancer. The methods generally involve detecting a palladin gene nucleotide sequence alteration that has been found to be associated with cancer and/or detecting a level of a palladin mRNA and/or protein in a biological sample. The present invention further provides

nucleic acid probes, nucleic acid primers, and antibodies, as well as kits comprising one or more of the same, for use in a subject method.

BRIEF DESCRIPTION OF THE DRAWINGS

[0007] FIGS. 1A and 1B provide a nucleotide sequence (AB023209; SEQ ID NO:1) of an mRNA encoding palladin. Mutations associated with pancreatic cancer are underlined and in bold text.

[0008] FIG. 2 depicts a pedigree of Family X.

[0009] FIGS. 3A-C depict identification of 4q32-34 genes differentially expressed in sporadic pancreatic cancer and in Family X pre-cancer.

[0010] FIG. 4 depicts location and identification of the Family X mutation.

[0011] FIG. 5 depicts an alignment of actinin binding site amino acid sequences of palladin from various species (human, chimp, mouse, rat: SEQ ID NO:103; chicken: SEQ ID NO:104; zebra fish: SEQ ID NO:105; and Family X: SEQ ID NO:106). The Pro→Ser substitution is boxed.

[0012] FIG. 6 depicts a blot of proteins extracted from various human cells and detected with polyclonal antibody against palladin raised in rabbit.

[0013] FIG. 7 is a graph depicting palladin expression in human ductal pancreatic epithelial (HDPE) primary cultures with neoplastic progression from normal ("HDPE Norm") to pancreatic intraepithelial neoplasia stage I ("HDPE PanIN I") to HDPE PanIN stage III ("HDPE PanIN III") to cancer ("HDPE PC").

[0014] FIG. 8A schematically depicts a model of interaction of palladin with actinin and ezrin.

[0015] FIG. 8B schematically depicts various palladin isoforms.

[0016] FIG. 9 depicts abnormal protein expression of palladin and α -actinin proteins in sporadic pancreatic cancer cell lines.

[0017] FIG. 10 depicts an exemplary palladin target nucleic acid (SEQ ID NO:63). The sequence corresponds to nucleotides 621-780 of a SEQ ID NO:1, and includes a C→T substitution (in bold and underlined) at a position corresponding to nucleotide 715 in SEQ ID NO:1. FIG. 10 also depicts an exemplary primer pair (SEQ ID NOs:64 and 65) for amplifying the exemplary target nucleic acid.

[0018] FIG. 11 depicts a palladin amino acid sequence (BAA76836.1; SEQ ID NO:99). Mutations associated with pancreatic cancer are underlined and in bold text.

[0019] FIG. 12 depicts a palladin amino acid sequence (NP_057165.3; SEQ ID NO:87).

[0020] FIGS. 13A and 13B depict a palladin mRNA nucleotide sequence (NM_016081; SEQ ID NO:20).

[0021] FIG. 14 depicts a genomic DNA sequence from GenBank Accession No. AC080188 (*Homo sapiens* BAC clone RP11-635L1; SEQ ID NO:101), including exon 9 sequences (depicted in bold text). Nucleotide 183561 (underlined and in bold text) corresponds to nucleotide 1671 of SEQ ID NO:1.

[0022] FIG. 15 depicts a genomic DNA sequence from GenBank Accession No. AC084353 (*Homo sapiens* BAC clone RP11-592K15; SEQ ID NO:102), including nucleotide 27251 (underlined and in bold text), which corresponds to nucleotide 2060 of the sequence depicted in FIG. 12.

[0023] FIG. 16 depicts a genomic DNA sequence from GenBank Accession No. AC080188 (*Homo sapiens* BAC clone RP11-635L1, including exon 2 sequences (depicted in

bold text). Nucleotide 140301 (underlined and in bold text) corresponds to nucleotide 715 of SEQ ID NO:1.

[0024] FIG. 17 depicts relative log expression, normalized to GAPDH and to a standard sample using the ddCt method, of palladin mRNA in white blood cells from normal individuals (“Nor” and “RS”), and individuals with pancreatic cancer (“PG” or “PA”).

[0025] FIG. 18 depicts the effect of palladin mutation on cell mobility.

[0026] FIG. 19 depicts the nucleotide sequence of a palladin 90 kDa isoform promoter, including a 12-base pair insertion. The promoter sequence is boxed; the 12-bp insertion is in bold.

DEFINITIONS

[0027] The terms “polynucleotide,” “oligonucleotide,” “nucleic acid” and “nucleic acid molecule” are used interchangeably herein to include a polymeric form of nucleotides, either ribonucleotides or deoxyribonucleotides. The terms refer only to the primary structure of the molecule. Thus, the terms include triple-, double- and single-stranded DNA, as well as triple-, double- and single-stranded RNA. RNA includes, e.g., messenger RNA. The terms “polynucleotide,” “oligonucleotide,” “nucleic acid” and “nucleic acid molecule” also include modifications, such as by methylation and/or by capping, and unmodified forms of the polynucleotide. More particularly, the terms “polynucleotide,” “oligonucleotide,” “nucleic acid” and “nucleic acid molecule” include polydeoxyribonucleotides (containing 2-deoxy-D-ribose), polyribonucleotides (containing D-ribose), any other type of polynucleotide which is an N- or C-glycoside of a purine or pyrimidine base, and other polymers containing nonnucleotidic backbones, for example, polyamide (e.g., peptide nucleic acids (PNAs)) and polymorpholino (commercially available from the Anti-Virals, Inc., Corvallis, Oreg., as Neugene) polymers, and other synthetic sequence-specific nucleic acid polymers providing that the polymers contain nucleobases in a configuration which allows for base pairing and base stacking, such as is found in DNA and RNA.

[0028] Unless specifically indicated otherwise, there is no intended distinction in length between the terms “polynucleotide,” “oligonucleotide,” “nucleic acid” and “nucleic acid molecule” and these terms will be used interchangeably. These terms refer only to the primary structure of the molecule. Thus, these terms include, for example, 3'-deoxy-2',5'-DNA, oligodeoxyribonucleotide N3' P5' phosphoramidates, 2'-O-alkyl-substituted RNA, double- and single-stranded DNA, as well as double- and single-stranded RNA, DNA:RNA hybrids, and hybrids between PNAs and DNA or RNA, and also include known types of modifications, for example, labels which are known in the art, methylation, “caps,” substitution of one or more of the naturally occurring nucleotides with an analog, internucleotide modifications such as, for example, those with uncharged linkages (e.g., methyl phosphonates, phosphotriesters, phosphoramidates, carbamates, etc.), with negatively charged linkages (e.g., phosphorothioates, phosphorodithioates, etc.), and with positively charged linkages (e.g., aminoalkylphosphoramidates, aminoalkylphosphotriesters), those containing pendant moieties, such as, for example, proteins (including nucleases, toxins, antibodies, signal peptides, poly-L-lysine, etc.), those with intercalators (e.g., acridine, psoralen, etc.), those containing chelators (e.g., metals, radioactive metals, boron, oxidative metals, etc.), those containing alkylators, those with modified

linkages (e.g., alpha anomeric nucleic acids, etc.), as well as unmodified forms of the polynucleotide or oligonucleotide. In particular, DNA is deoxyribonucleic acid.

[0029] Throughout the specification, abbreviations are used to refer to nucleotides (also referred to as bases), including abbreviations that refer to multiple nucleotides. As used herein, G=guanine, A=adenine, T=thymine, C=cytosine, and U=uracil. In addition, R=a purine nucleotide (A or G); Y=a pyrimidine nucleotide (C or T (U)); S=C or G; W=A or T (U); M=A or C; K=G or T (U); V=A, C or G; and N=any nucleotide (A, T (U), C, or G). Nucleotides can be referred to throughout using lower or upper case letters. It is also understood that nucleotides sequences provided for DNA in the specification also represent nucleotide sequences for RNA, where T is substituted by U.

[0030] The terms “deoxyribonucleic acid” and “DNA” as used herein mean a polymer composed of deoxyribonucleotides.

[0031] The terms “ribonucleic acid” and “RNA” as used herein refer to a polymer composed of ribonucleotides. Where sequences of a nucleic acid are provided using nucleotides of a DNA sequence, it is understood that such sequences encompass complementary DNA sequences and further also encompass RNA sequences based on the given DNA sequence or its complement, where uracil (U) replaces thymine (T) in the DNA sequence or its complement.

[0032] Two nucleotide sequences are “complementary” to one another when those molecules share base pair organization homology. “Complementary” nucleotide sequences will combine with specificity to form a stable duplex under appropriate hybridization conditions. For instance, two sequences are complementary when a section of a first sequence can bind to a section of a second sequence in an anti-parallel sense wherein the 3'-end of each sequence binds to the 5'-end of the other sequence and each A, T(U), G, and C of one sequence is then aligned with a T(U), A, C, and G, respectively, of the other sequence. RNA sequences can also include complementary G=U or U=G base pairs. Thus, two sequences need not have perfect homology to be “complementary” under the invention. Usually two sequences are sufficiently complementary when at least about 85%, at least about 90%, at least about 95%, or at least about 98% of the nucleotides share base pair organization over a defined length of the molecule.

[0033] As used herein the term “isolated,” when used in the context of an isolated compound, refers to a compound of interest that is in an environment different from that in which the compound naturally occurs. “Isolated” is meant to include compounds that are within samples that are substantially enriched for the compound of interest and/or in which the compound of interest is partially or substantially purified. The term “isolated” encompasses instances in which the recited material is unaccompanied by at least some of the material with which it is normally associated in its natural state, where the isolated compound constitutes at least about 50%, at least about 75%, or at least about 90%, or more, by weight of the total protein in a given sample. For example, the term “isolated” with respect to a polynucleotide generally refers to a nucleic acid molecule devoid, in whole or part, of sequences normally associated with it in nature; or a sequence, as it exists in nature, but having heterologous sequences in association therewith; or a molecule disassociated from the chromosome.

[0034] “Purified” as used herein means that the recited material comprises at least about 75%, at least about 80%, or

at least about 90% by weight of the total material. As used herein, the term “substantially pure” refers to a compound that is removed from its natural environment and is at least 60% free, at least 75% free, at least 90% free, at least 95% free, at least 98% free, or at least 99% or more free, from other components with which it is naturally associated.

[0035] A polynucleotide “derived from” or “specific for” a designated sequence, such as a target sequence of a target nucleic acid, refers to a polynucleotide sequence which comprises a contiguous sequence of approximately at least about 6 nucleotides, at least about 8 nucleotides, at least about 10-12 nucleotides, or at least about 15-20 nucleotides corresponding to, i.e., identical or complementary to, a region of the designated nucleotide sequence. The derived polynucleotide will not necessarily be derived physically from the nucleotide sequence of interest, but may be generated in any manner, including, but not limited to, chemical synthesis, replication, reverse transcription or transcription, which is based on the information provided by the sequence of bases in the region (s) from which the polynucleotide is derived or specific for. Polynucleotides that are derived from” or “specific for” a designated sequence include polynucleotides that are in a sense or an antisense orientation relative to the original polynucleotide.

[0036] A polynucleotide or polypeptide has a certain percent “sequence identity” to another polynucleotide or polypeptide, meaning that, when aligned, that percentage of bases or amino acids are the same, and in the same relative position, when comparing the two sequences. Sequence similarity can be determined in a number of different manners. To determine sequence identity, sequences can be aligned using the methods and computer programs, including BLAST, available over the world wide web at ncbi.nlm.nih.gov/BLAST. See, e.g., Altschul et al. (1990), *J. Mol. Biol.* 215: 403-10. Another alignment algorithm is FASTA, available in the Genetics Computing Group (GCG) package, from Madison, Wis., USA, a wholly owned subsidiary of Oxford Molecular Group, Inc. Other techniques for alignment are described in *Methods in Enzymology*, vol. 266: Computer Methods for Macromolecular Sequence Analysis (1996), ed. Doolittle, Academic Press, Inc., a division of Harcourt Brace & Co., San Diego, Calif., USA. Of particular interest are alignment programs that permit gaps in the sequence. The Smith-Waterman is one type of algorithm that permits gaps in sequence alignments. See *Meth. Mol. Biol.* 70: 173-187 (1997). Also, the GAP program using the Needleman and Wunsch alignment method can be utilized to align sequences. See *J. Mol. Biol.* 48: 443-453 (1970).

[0037] A nucleic acid is “hybridizable” to another nucleic acid, such as a cDNA, genomic DNA, or RNA, when a single stranded form of the nucleic acid can anneal to the other nucleic acid under the appropriate conditions of temperature and solution ionic strength. Hybridization and washing conditions are well known and exemplified in Sambrook, J., Fritsch, E. F. and Maniatis, T. *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor (1989), particularly Chapter 11 and Table 11.1 therein; and Sambrook, J. and Russell, W., *Molecular Cloning: A Laboratory Manual*, Third Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor (2001). The conditions of temperature and ionic strength determine the “stringency” of the hybridization. Stringency conditions can be adjusted to screen for moderately similar fragments, such as homologous sequences from distantly

related organisms, to highly similar fragments, such as genes that duplicate functional enzymes from closely related organisms.

[0038] Hybridization conditions and post-hybridization washes are useful to obtain the desired determine stringency conditions of the hybridization. One set of illustrative post-hybridization washes is a series of washes starting with 6×SSC (where SSC is 0.15 M NaCl and 15 mM citrate buffer), 0.5% SDS at room temperature for 15 minutes, then repeated with 2×SSC, 0.5% SDS at 45° C. for 30 minutes, and then repeated twice with 0.2×SSC, 0.5% SDS at 50° C. for 30 minutes. Other stringent conditions are obtained by using higher temperatures in which the washes are identical to those above except for the temperature of the final two 30 minute washes in 0.2×SSC, 0.5% SDS, which is increased to 60° C. Another set of highly stringent conditions uses two final washes in 0.1×SSC, 0.1% SDS at 65° C. Another example of stringent hybridization conditions is hybridization at 50° C. or higher and 0.1×SSC (15 mM sodium chloride/1.5 mM sodium citrate). Another example of stringent hybridization conditions is overnight incubation at 42° C. in a solution: 50% formamide, 5×SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5×Denhardt’s solution, 10% dextran sulfate, and 20 µg/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1×SSC at about 65° C. Stringent hybridization conditions and post-hybridization wash conditions are hybridization conditions and post-hybridization wash conditions that are at least as stringent as the above representative conditions.

[0039] In certain embodiments, the stringency of the wash conditions that set forth the conditions which determine whether a nucleic acid is specifically hybridized to a probe. Wash conditions used to identify nucleic acids may include, e.g.: a salt concentration of about 0.02 molar at pH 7 and a temperature of at least about 50° C. or about 55° C. to about 60° C.; or, a salt concentration of about 0.15 M NaCl at 72° C. for about 15 minutes; or, a salt concentration of about 0.2×SSC at a temperature of at least about 50° C. or about 55° C. to about 60° C. for about 15 to about 20 minutes; or, the hybridization complex is washed twice with a solution with a salt concentration of about 2×SSC containing 0.1% SDS at room temperature for 15 minutes and then washed twice by 0.1×SSC containing 0.1% SDS at 68° C. for 15 minutes; or, equivalent conditions. Stringent conditions for washing can also be, e.g., 0.2×SSC/0.1% SDS at 42° C. In instances wherein the nucleic acid molecules are deoxyoligonucleotides (“oligos”), stringent conditions can include washing in 6×SSC/0.05% sodium pyrophosphate at 37° C. (for 14-base oligos), 48° C. (for 17-base oligos), 55° C. (for 20-base oligos), and 60° C. (for 23-base oligos). See Sambrook, Ausubel, or Tijssen (cited below) for detailed descriptions of equivalent hybridization and wash conditions and for reagents and buffers, e.g., SSC buffers and equivalent reagents and conditions.

[0040] Hybridization requires that the two nucleic acids contain complementary sequences, although depending on the stringency of the hybridization, mismatches between bases are possible. The appropriate stringency for hybridizing nucleic acids depends on the length of the nucleic acids and the degree of complementation, variables well known in the art. The greater the degree of similarity or homology between two nucleotide sequences, the greater the value of the melting temperature (T_m) for hybrids of nucleic acids having those sequences. The relative stability (corresponding to higher

T_m) of nucleic acid hybridizations decreases in the following order: RNA:RNA, DNA:RNA, DNA:DNA. For hybrids of greater than 100 nucleotides in length, equations for calculating T_m have been derived (see Sambrook et al., supra, 9.50-9.51). For hybridizations with shorter nucleic acids, i.e., oligonucleotides, the position of mismatches becomes more important, and the length of the oligonucleotide determines its specificity (see Sambrook et al., supra, 11.7-11.8). Typically, the length for a hybridizable nucleic acid is at least about 10 nucleotides. Illustrative minimum lengths for a hybridizable nucleic acid are: at least about 15 nucleotides; at least about 20 nucleotides; and at least about 30 nucleotides. Furthermore, the skilled artisan will recognize that the temperature and wash solution salt concentration may be adjusted as necessary according to factors such as length of the probe.

[0041] The terms “peptide,” “polypeptide,” and “protein” are used interchangeably herein, and refer to a polymeric form of amino acids of any length, which can include coded and non-coded amino acids, chemically or biochemically modified or derivatized amino acids, and polypeptides having modified peptide backbones.

[0042] A “biological sample” encompasses a variety of sample types obtained from an individual and can be used in a diagnostic or monitoring assay. The definition encompasses blood and other liquid samples of biological origin, solid tissue samples such as a biopsy specimen or tissue cultures or cells derived therefrom and the progeny thereof. The definition also includes samples that have been manipulated in any way after their procurement, such as by treatment with reagents, solubilization, or enrichment for certain components, such as polynucleotides or polypeptides. The term “biological sample” encompasses a clinical sample, and also includes cells in culture, cell supernatants, cell lysates, serum, plasma, biological fluid, and tissue samples. In many embodiments, a biological sample will include cells (e.g., pancreatic cells; non-pancreatic cells).

[0043] In some embodiments, a biological sample will include tissue. In some embodiments, a biological sample will include RNA (e.g., mRNA) obtained from cells or tissue; a cDNA copy of an RNA obtained from cells or tissues; an amplified copy of an mRNA or a cDNA obtained from cells or tissues; and the like. In some embodiments, a biological sample will include genomic DNA obtained from cells or tissue. The tissue may appear histologically normal, or may appear histologically abnormal (e.g., neoplastic).

[0044] In some embodiments, a biological sample will include tissue (e.g., pancreatic tissue; non-pancreatic tissue). In some embodiments, a biological sample will include RNA (e.g., mRNA) obtained from cells or tissue (e.g., pancreatic cells; non-pancreatic cells; pancreatic tissue; non-pancreatic tissue); a cDNA copy of an RNA obtained from cells or tissues; an amplified copy of an mRNA or a cDNA obtained from cells or tissues; and the like. In some embodiments, a biological sample will include genomic DNA obtained from cells or tissue (e.g., pancreatic cells; non-pancreatic cells; pancreatic tissue; non-pancreatic tissue).

[0045] An “indicator cell” as used herein is a cell which is subjected to analysis for a parameter that is indicative of the presence of cancer in a subject from whom the indicator cell was obtained. Where the parameter is the presence or absence of a palladin mutation, the indicator cell is a cell from blood or a tissue suspected of containing a pre-cancerous or cancerous cell having a palladin mutation (e.g., pancreatic cell,

breast cell, head and neck squamous cell, and the like). Where the parameter is the presence or absence of an aberrant expression level of palladin, the indicator cell can be a cell from blood or a tissue suspected of having an aberrant expression level of palladin or a cell other than a cancerous or pre-cancerous cell, e.g., a white blood cell. “Indicator cell” as used herein is most often used in the context of detection of palladin expression levels in a white blood cell, but could represent any cell in the body.

[0046] A “DNA-dependent DNA polymerase” is an enzyme that synthesizes a complementary DNA copy from a DNA template. Examples include DNA polymerase I from *E. coli* and bacteriophage T7 DNA polymerase. All known DNA-dependent DNA polymerases require a complementary primer to initiate synthesis. Under suitable conditions, a DNA-dependent DNA polymerase may synthesize a complementary DNA copy from an RNA template.

[0047] A “DNA-dependent RNA polymerase” or a “transcriptase” is an enzyme that synthesizes multiple RNA copies from a double-stranded or partially-double stranded DNA molecule having a (usually double-stranded) promoter sequence. The RNA molecules (“transcripts”) are synthesized in the 5' to 3' direction beginning at a specific position just downstream of the promoter. Examples of transcriptases are the DNA-dependent RNA polymerase from *E. coli* and bacteriophages T7, T3, and SP6.

[0048] An “RNA-dependent DNA polymerase” or “reverse transcriptase” is an enzyme that synthesizes a complementary DNA copy from an RNA template. All known reverse transcriptases also have the ability to make a complementary DNA copy from a DNA template; thus, they are both RNA- and DNA-dependent DNA polymerases. A primer is required to initiate synthesis with both RNA and DNA templates.

[0049] “RNase H” is an enzyme that degrades the RNA portion of an RNA:DNA duplex. These enzymes may be endonucleases or exonucleases. Most reverse transcriptase enzymes normally contain an RNase H activity in addition to their polymerase activity. However, other sources of the RNase H are available without an associated polymerase activity. RNA degradation mediated by an RNase H may result in separation of RNA from an RNA:DNA complex, or the RNase H may cut the RNA at various locations such that portions of the RNA melt off or permit enzymes to unwind portions of the RNA.

[0050] As used herein, the term “target nucleic acid region” or “target nucleic acid” or “target molecules” refers to a nucleic acid molecule with a “target sequence” to be detected (e.g., in a method involving nucleic acid hybridization and/or amplification). The target nucleic acid may be either single-stranded or double-stranded and may or may not include other sequences besides the target sequence (e.g., the target nucleic acid may or may not include nucleic acid sequences upstream or 5' flanking sequence, may or may not include downstream or 3' flanking sequence, and in some embodiments may not include either upstream (5') or downstream (3') nucleic acid sequence relative to the target sequence. Where detection is by amplification, these other sequences in addition to the target sequence may or may not be amplified with the target sequence.

[0051] The term “target sequence” or “target nucleic acid sequence” refers to the particular nucleotide sequence of the target nucleic acid to be detected (e.g., through amplification). The target sequence may include a probe-hybridizing region contained within the target molecule with which a

probe will form a stable hybrid under desired conditions. The “target sequence” may also include the complexing sequences to which the oligonucleotide primers complex and be extended using the target sequence as a template. Where the target nucleic acid is originally single-stranded, the term “target sequence” also refers to the sequence complementary to the “target sequence” as present in the target nucleic acid. If the “target nucleic acid” is originally double-stranded, the term “target sequence” refers to both the plus (+) and minus (-) strands. Moreover, where sequences of a “target sequence” are provided herein, it is understood that the sequence may be either DNA or RNA. Thus where a DNA sequence is provided, the RNA sequence is also contemplated and is readily provided by substituting “T” of the DNA sequence with “U” to provide the RNA sequence.

[0052] The term “primer” or “oligonucleotide primer” as used herein, refers to an oligonucleotide which acts to initiate synthesis of a complementary nucleic acid strand when placed under conditions in which synthesis of a primer extension product is induced, e.g., in the presence of nucleotides and a polymerization-inducing agent such as a DNA or RNA polymerase and at suitable temperature, pH, metal ion concentration, and salt concentration. Primers are generally of a length compatible with its use in synthesis of primer extension products, and are in many embodiments in the range of between 8 nucleotides and 100 nucleotides (nt) in length, such as 10 nt to 75 nt, 15 nt to 60 nt, 15 nt to 40 nt, 18 nt to 30 nt, 20 nt to 40 nt, 21 nt to 50 nt, 22 nt to 45 nt, 25 nt to 40 nt, and so on, e.g., in the range of between 18 nt and 40 nt, between 20 nt and 35 nt, between 21 and 30 nt in length, inclusive, and any length between the stated ranges. Primers can be in the range of between 10-50 nucleotides long, such as 15-45, 18-40, 20-30, 21-25 nt and so on, and any length between the stated ranges. In some embodiments, the primers are not more than about 10, 12, 15, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, 55, 60, 65, or 70 nucleotides in length.

[0053] Primers are in many embodiments single-stranded for maximum efficiency in amplification, but may alternatively be double-stranded. If double-stranded, the primer is in many embodiments first treated to separate its strands before being used to prepare extension products. This denaturation step is typically effected by heat, but may alternatively be carried out using alkali, followed by neutralization. Thus, a “primer” is complementary to a template, and complexes by hydrogen bonding or hybridization with the template to give a primer/template complex for initiation of synthesis by a polymerase, which is extended by the addition of covalently bonded bases linked at its 3' end complementary to the template in the process of DNA synthesis.

[0054] A “primer pair” as used herein refers to first and second primers having nucleic acid sequence suitable for nucleic acid-based amplification of a target nucleic acid. Such primer pairs generally include a first primer having a sequence that is the same or similar to that of a first portion of a target nucleic acid, and a second primer having a sequence that is complementary to a second portion of a target nucleic acid to provide for amplification of the target nucleic acid or a fragment thereof. Reference to “first” and “second” primers herein is arbitrary, unless specifically indicated otherwise. For example, the first primer can be designed as a “forward primer” (which initiates nucleic acid synthesis from a 5' end of the target nucleic acid) or as a “reverse primer” (which initiates nucleic acid synthesis from a 5' end of the extension

product produced from synthesis initiated from the forward primer). Likewise, the second primer can be designed as a forward primer or a reverse primer.

[0055] As used herein, the term “probe” or “oligonucleotide probe”, used interchangeably herein, refers to a structure comprised of a polynucleotide, as defined above, which contains a nucleic acid sequence complementary to a nucleic acid sequence present in the target nucleic acid analyte (e.g., a nucleic acid amplification product). The polynucleotide regions of probes may be composed of DNA, and/or RNA, and/or synthetic nucleotide analogs. Probes are generally of a length compatible with its use in specific detection of all or a portion of a target sequence of a target nucleic acid, and are in many embodiments in the range of between 8 nt and 100 nt in length, such as 8 to 75 nt, 10 to 74 nt, 12 to 72 nt, 15 to 60 nt, 15 to 40 nt, 18 to 30 nt, 20 to 40 nt, 21 to 50 nt, 22 to 45 nt, 25 to 40 nt in length, and so on, e.g., in the range of between 18-40 nt, 20-35 nt, or 21-30 nt in length, and any length between the stated ranges. In some embodiments, a probe is in the range of between 10-50 nucleotides long, such as 15-45, 18-40, 20-30, 21-28, 22-25 and so on, and any length between the stated ranges. In some embodiments, the primers are not more than about 10, 12, 15, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, 55, 60, 65, or 70 nucleotides in length.

[0056] Probes contemplated herein include probes that include a detectable label. For example, when an “oligonucleotide probe” is to be used in a 5' nuclease assay, such as the TAQMAN™ assay, the probe includes at least one fluorescer and at least one quencher which is digested by the 5' endonuclease activity of a polymerase used in the reaction in order to detect any amplified target oligonucleotide sequences. In this context, the oligonucleotide probe will have a sufficient number of phosphodiester linkages adjacent to its 5' end so that the 5' to 3' nuclease activity employed can efficiently degrade the bound probe to separate the fluorescers and quenchers. When an oligonucleotide probe is used in the TMA technique, it will be suitably labeled, as described below.

[0057] Probes and primers contemplated herein include those useful in various amplification and/or detection systems, including those in which primers and probes are provided as bi-functional molecules. Exemplary amplification and/or detection systems include Sunrise™ primer-based systems, Molecular Beacons, the Taqman™ system, an Amplifluor™ hairpin primer-based system, a Scorpions technology (e.g., bi-functional molecules containing a PCR primer element covalently linked to a probe element), and a Light Upon Extension or LUX™-based system. Further exemplary detection systems include those based on a melt-curve analysis, and using intercalating dyes such as the fluorescent dye SYBR Green.

[0058] As used herein, the terms “label” and “detectable label” refer to a molecule capable of detection, including, but not limited to, radioactive isotopes, fluorescers, chemiluminescers, chromophores, enzymes, enzyme substrates, enzyme cofactors, enzyme inhibitors, chromophores, dyes, metal ions, metal sols, ligands (e.g., biotin, avidin, streptavidin or haptens), intercalating dyes and the like. The term “fluorescer” refers to a substance or a portion thereof which is capable of exhibiting fluorescence in the detectable range.

[0059] The terms “hybridize” and “hybridization” refer to the formation of complexes between nucleotide sequences which are sufficiently complementary to form complexes via Watson-Crick base pairing. Where a primer “hybridizes” with target (template), such complexes (or hybrids) are suf-

ficiently stable to serve the priming function required by, e.g., the DNA polymerase to initiate DNA synthesis.

[0060] The term “stringent conditions” refers to conditions under which a primer will hybridize preferentially to, or specifically bind to, its complementary binding partner, and to a lesser extent to, or not at all to, other sequences. Put another way, the term “stringent hybridization conditions” as used herein refers to conditions that are compatible to produce duplexes on an array surface between complementary binding members, e.g., between probes and complementary targets in a sample, e.g., duplexes of nucleic acid probes, such as DNA probes, and their corresponding nucleic acid targets that are present in the sample, e.g., their corresponding mRNA analytes present in the sample.

[0061] The term “assessing” includes any form of measurement, and includes determining if an element is present or not. The terms “determining,” “measuring,” “evaluating,” “assessing,” and “assaying” are used interchangeably and includes quantitative and qualitative determinations. Assessing may be relative or absolute. “Assessing the presence of” includes determining the amount of something present, and/or determining whether it is present or absent. As used herein, the terms “determining,” “measuring,” and “assessing,” and “assaying” are used interchangeably and include both quantitative and qualitative determinations.

[0062] The term “binds specifically,” in the context of antibody binding, refers to high avidity and/or high affinity binding of an antibody to a specific polypeptide i.e., epitope of a polypeptide, e.g., a palladin polypeptide. For example, antibody binding to an epitope on a specific subject deacylase or fragment thereof is stronger than binding of the same antibody to any other epitope, particularly those which may be present in molecules in association with, or in the same sample, as the specific polypeptide of interest, e.g., binds more strongly to a specific palladin polypeptide than to any other palladin epitopes so that by adjusting binding conditions the antibody binds almost exclusively to the specific palladin epitope and not to any other palladin epitope, or to any other polypeptide which does not comprise the epitope. Antibodies that bind specifically to a polypeptide may be capable of binding other polypeptides at a weak, yet detectable, level (e.g., 10% or less of the binding shown to the polypeptide of interest). Such weak binding, or background binding, is readily discernible from the specific antibody binding to a palladin polypeptide, e.g. by use of appropriate controls. In general, specific antibodies bind to a given polypeptide with a binding affinity of 10^{-7} M or more, e.g., 10^{-8} M or more (e.g., 10^{-9} M, 10^{-10} M, 10^{-11} M, etc.). In general, an antibody with a binding affinity of 10^{-6} M or less is not useful in that it will not bind an antigen at a detectable level using conventional methodology currently used.

[0063] The terms “cancer,” “neoplasm,” and “tumor” are used interchangeably herein to refer to cells that exhibit relatively autonomous growth, so that they exhibit an aberrant growth phenotype characterized by a significant loss of control of cell proliferation. Cancerous cells can be benign or malignant.

[0064] The terms “individual,” “host,” “subject,” and “patient,” used interchangeably herein, refer to a mammal, including human and non-human primates; rodents (e.g., mice, rats, etc.); livestock (e.g., bovine, ovine, caprine, porcine, etc., mammals); mammalian pets (e.g., canines, felines, etc.); horses; lagomorphs; etc. In some embodiments, an individual is a human.

[0065] Before the present invention is further described, it is to be understood that this invention is not limited to particular embodiments described, as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present invention will be limited only by the appended claims.

[0066] Where a range of values is provided, it is understood that each intervening value, to the tenth of the unit of the lower limit unless the context clearly dictates otherwise, between the upper and lower limit of that range and any other stated or intervening value in that stated range, is encompassed within the invention. The upper and lower limits of these smaller ranges may independently be included in the smaller ranges, and are also encompassed within the invention, subject to any specifically excluded limit in the stated range. Where the stated range includes one or both of the limits, ranges excluding either or both of those included limits are also included in the invention.

[0067] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can also be used in the practice or testing of the present invention, the preferred methods and materials are now described. All publications mentioned herein are incorporated herein by reference to disclose and describe the methods and/or materials in connection with which the publications are cited.

[0068] It must be noted that as used herein and in the appended claims, the singular forms “a,” “and,” and “the” include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to “a nucleic acid probe” includes a plurality of such probes and reference to “the nucleic acid primer pair” includes reference to one or more nucleic acid primer pairs and equivalents thereof known to those skilled in the art, and so forth. It is further noted that the claims may be drafted to exclude any optional element. As such, this statement is intended to serve as antecedent basis for use of such exclusive terminology as “solely,” “only” and the like in connection with the recitation of claim elements, or use of a “negative” limitation.

[0069] The publications discussed herein are provided solely for their disclosure prior to the filing date of the present application. Nothing herein is to be construed as an admission that the present invention is not entitled to antedate such publication by virtue of prior invention. Further, the dates of publication provided may be different from the actual publication dates which may need to be independently confirmed.

DETAILED DESCRIPTION

[0070] The present invention provides methods for detecting and diagnosing cancer in an individual; methods of identifying individuals at risk of developing a cancer; and methods of staging a cancer and/or metastases. The methods generally involve detecting a palladin gene nucleotide sequence alteration (e.g., a substitution, an insertion, a deletion, a change in methylation status, etc.) that has been found to be associated with cancer and/or detecting a level of a palladin mRNA or palladin polypeptide in a cell or tissue. The present invention further provides nucleic acid probes and nucleic acid primers, as well as kits comprising same, for use

in a subject method. It is to be understood that this invention is not limited to the particular methods and materials described herein.

[0071] The present invention provides, for example, methods for detecting and diagnosing pancreatic cancer in an individual; methods of identifying individuals at risk of developing pancreatic cancer; and methods of staging pancreatic cancer. The methods generally involve detecting a palladin gene nucleotide sequence alteration that has been found to be associated with pancreatic cancer and/or detecting a level of a palladin mRNA or palladin polypeptide in a cell or tissue. The present invention further provides nucleic acid probes and nucleic acid primers, as well as kits comprising same, for use in a subject method.

[0072] The following observations were made: 1) a C→T nucleotide substitution at position 715 of the human palladin gene is associated with increased risk of developing pancreatic cancer; 2) a G→T nucleotide substitution at position 1671 of the human palladin gene is associated with increased risk of developing pancreatic cancer; 3) a C→T nucleotide substitution at position 2060 of the human palladin gene is associated with cancerous phenotype in pancreatic cells; 4) palladin mRNA, e.g., palladin mRNA that encodes the 90 kD isoform of palladin, is overexpressed in pre-cancerous and pancreatic cells; 5) the level of palladin mRNA, e.g., palladin mRNA that encodes the 90 kD isoform of palladin, in a pancreatic cell increases with neoplastic progression; 6) lower than normal levels of palladin mRNA in white blood cells is associated with pancreatic cancer; 7) palladin mRNA is abnormally expressed in a variety of cancers; and 8) a 12-base pair insertion in the promoter of the 90 kDa isoform of palladin is associated with an increased risk of developing pancreatic cancer.

[0073] Cancers that can be detected and/or diagnosed using a subject method include, but are not limited to, pancreatic cancer, breast cancer, and head and neck squamous cell carcinoma.

[0074] Methods for detecting a cancerous or precancerous cell, e.g., a cancerous or precancerous pancreatic cell, as well as methods for identifying individuals at risk of developing a cancer such as pancreatic cancer, may involve detecting a nucleotide substitution in a palladin target nucleic acid (such as palladin genomic DNA; palladin mRNA; a cDNA copy of a palladin mRNA; an amplified copy of a palladin mRNA or cDNA) present in a biological sample. Examples of palladin gene nucleotide substitutions associated with a cancerous or precancerous cell (e.g., a precancerous pancreatic cell, a cancerous pancreatic cell) and/or an increased risk of developing cancer (e.g., pancreatic cancer) include the following: 1) a C→T nucleotide substitution at position 715 of AB023209 (see, e.g., FIG. 1A); 2) a G→T nucleotide substitution at position 1671 of AB023209 (see, e.g., FIG. 1A); and 3) a C→T nucleotide substitution at position 2060 of NM_016081 (see, e.g., FIGS. 13A and 13B). Mutations in palladin, however, are not limited to the ones listed herein.

[0075] Methods for detecting a cancerous or precancerous cell, e.g., a cancerous or precancerous pancreatic cell, as well as methods for identifying individuals at risk of developing a cancer such as pancreatic cancer, can also involve detecting an insertion in a palladin target nucleic acid (such as palladin genomic DNA; palladin mRNA; a cDNA copy of a palladin mRNA; an amplified copy of a palladin mRNA or cDNA) present in a biological sample. Examples of palladin gene nucleotide sequence insertions associated with a cancerous or

precancerous cell (e.g., a precancerous pancreatic cell, a cancerous pancreatic cell) and/or an increased risk of developing cancer (e.g., pancreatic cancer) include an insertion in the promoter region of the gene encoding the 90 kDa isoform of palladin.

[0076] Methods for detecting a cancerous or precancerous cell, e.g., a cancerous or precancerous pancreatic cell, as well as methods for identifying individuals at risk of developing a cancer such as pancreatic cancer, can also involve detecting a change in the methylation status of palladin target nucleic acid (such as palladin genomic DNA; palladin mRNA; a cDNA copy of a palladin mRNA; an amplified copy of a palladin mRNA or cDNA) present in a biological sample. For example, hypomethylation of a palladin gene promoter associated with a cancerous or precancerous cell (e.g., a precancerous pancreatic cell, a cancerous pancreatic cell) and/or an increased risk of developing cancer (e.g., pancreatic cancer).

[0077] Detection of a nucleotide sequence alteration in a palladin target nucleic acid can be carried out using, e.g., nucleic acid hybridization with a nucleic acid probe specific for the nucleotide substitution. Detection of a nucleotide sequence alteration in a palladin target nucleic acid can also be carried out by first amplifying a target palladin nucleic acid that comprises the nucleotide substitution, using a nucleic acid primer pair; then detecting the nucleotide substitution by hybridization with a nucleic acid probe.

[0078] For example, detection of the 715 C→T nucleotide substitution can be carried out using, e.g., nucleic acid hybridization with a nucleic acid probe specific for the 715 C→T nucleotide substitution. Detection of the 715 C→T nucleotide substitution can also be carried out by first amplifying a target palladin nucleic acid that comprises the 715 C→T nucleotide substitution, using a nucleic acid primer pair; then detecting the 715 C→T nucleotide substitution by hybridization with a nucleic acid probe.

[0079] Methods for detecting a cancerous or precancerous cell, as well as methods of staging a cancer, may involve detecting a level of palladin mRNA present in a biological sample (for example, a cell, or a biological sample comprising a cell). Detecting a level of palladin mRNA present in a cell may involve nucleic acid hybridization with a nucleic acid probe that detects a target palladin nucleic acid that is overexpressed in a cancerous cell (e.g., a cancerous pancreatic cell). Detecting a level of palladin mRNA present in a cell may involve nucleic acid hybridization with a nucleic acid probe that detects a target palladin nucleic acid that is expressed at lower than normal levels in an indicator cell, e.g., a cell other than a cancer cell. Detecting a level of palladin mRNA can also be carried out using any of a variety of nucleic acid amplification methods, e.g., a quantitative polymerase chain reaction or other nucleic acid amplification method.

[0080] Methods for detecting a cancerous or precancerous cell, as well as methods of staging a cancer, may involve detecting a level of palladin protein/polypeptide present in a biological sample. Detecting a level of palladin protein/polypeptide present in a cell or tissue or fluid may involve protein binding with an antibody probe that detects a target palladin protein that is overexpressed in a pre-cancerous or cancerous biologic sample (for example, cancerous pancreatic tissue). Examples of methods for protein detection include, but are not limited, to Western Blotting, enzyme-linked immunosorbent assays, immunostaining, and in-situ hybridization. Other methods of detecting polypeptides

include direct detection through the use of proteomics, such as mass spectroscopy measurements, and the like.

[0081] As discussed above, subject detection, diagnostic, and staging methods generally involve nucleic acid hybridization, and/or nucleic acid amplification, and/or protein detection. The present invention provides nucleic acid probes and nucleic acid primers that are suitable for use in such methods. In some embodiments, nucleic acid probes and nucleic acid primer pairs are useful for detecting a palladin nucleic acid comprising a nucleotide sequence alteration that is associated with pancreatic cancer. In other embodiments, nucleic acid probes and nucleic acid primer pairs are useful for detecting a palladin nucleic acid that is over-expressed in a pancreatic cell. Probes and primers useful in detecting a palladin nucleic acid comprising a nucleotide substitution can also be used in detecting palladin expression levels. Kits that include a nucleic acid probe, a nucleic acid primer pair, an antibody reagent, or two or more of the foregoing, are also provided.

Target Nucleic Acids

[0082] Target palladin nucleic acids include palladin nucleic acids that include one or more palladin gene nucleotide sequence alterations (e.g., substitutions, deletions, insertions, etc.) that are associated with an increased risk of developing cancer (e.g., pancreatic cancer, breast cancer, head and neck squamous cell carcinoma, etc.) and/or are associated with a precancerous or cancerous state of a cell such as a pancreatic cell and/or are indicative of the presence of a cancerous cell in the individual. For example, a palladin gene nucleotide sequence alteration will in some cases be associated with abnormal levels of palladin mRNA and/or polypeptide in a pancreatic cell.

[0083] In some embodiments, abnormal levels of a palladin mRNA that, when present in a cell, are associated with a precancerous or cancerous state of the cell, are levels that are significantly higher or lower than normal levels of the palladin mRNA found in a non-cancerous cell of the same cell type. In some embodiments, abnormal levels of a palladin mRNA that, when present in a test cell, are indicative of the presence of a cancerous cell in the individual from whom the test cell was obtained, are levels that are significantly higher or lower than normal levels of the palladin mRNA typically found in the test cell in an individual who does not have cancer.

[0084] An abnormally high level of a palladin mRNA that, when present in a cell, is associated with a precancerous or cancerous state of the cell, is a level that is at least about 25%, at least about 50%, at least about 100% (or 2-fold), at least about 3-fold, at least about 5-fold, at least about 10-fold, at least about 25-fold, or at least about 50-fold, or more, higher than the level of the palladin mRNA in a non-cancerous cell of the same cell type.

[0085] For example, an abnormally high level of a palladin mRNA that, when present in a pancreatic cell, is associated with a precancerous or cancerous state of the cell, is a level that is at least about 25%, at least about 50%, at least about 100% (or 2-fold), at least about 3-fold, at least about 5-fold, at least about 10-fold, at least about 25-fold, or at least about 50-fold, or more, higher than the level of the palladin mRNA in a non-cancerous pancreatic cell.

[0086] An abnormally low level of a palladin mRNA that, when present in a cell, is associated with a precancerous or cancerous state of the cell, is a level that is about 75% or less,

about 60% or less, about 50% or less, about 25% or less, or about 10% or less, than the level of the palladin mRNA in a non-cancerous cell of the same cell type

[0087] For example, an abnormally low level of a palladin mRNA that, when present in a pancreatic cell, is associated with a precancerous or cancerous state of the cell, is a level that is about 75% or less, about 60% or less, about 50% or less, about 25% or less, or about 10% or less, than the level of the palladin mRNA in a non-cancerous pancreatic cell.

[0088] An abnormally low level of a palladin mRNA that, when present in a test cell (also referred to as an "indicator cell"), is indicative of the presence of a cancerous cell in the individual from whom the test cell was obtained, is a level that is about 75% or less, about 60% or less, about 50% or less, about 25% or less, or about 10% or less, than the level of the palladin mRNA typically found in the test cell in an individual who does not have cancer. As one non-limiting example, it was found that the level of palladin mRNA in white blood cells (WBC) from individuals with pancreatic cancer was abnormally low when compared to the level of palladin mRNA present in WBC from normal individuals (e.g., individuals without pancreatic cancer).

[0089] Non-limiting examples of palladin gene nucleotide substitutions associated with a precancerous or cancerous cell (e.g., a precancerous pancreatic cell, a cancerous pancreatic cell) and/or an increased risk of developing a cancer such as pancreatic cancer include the following: 1) a C→T nucleotide substitution at position 715 of AB023209 (see, e.g., FIG. 1A); 2) a G→T nucleotide substitution at position 1671 of AB023209 (see, e.g., FIG. 1A); and 3) a C→T nucleotide substitution at position 2060 of NM_016081 (see, e.g., FIGS. 13A and 13B).

[0090] It should be noted that in the diagnostic methods based on detection of palladin mutations or palladin expression levels, while detection of a palladin mutation and/or aberrant (abnormal) palladin expression levels indicates the presence of a cancerous or pre-cancerous cell in the subject, the detection of the absence of a palladin mutation or the absence of aberrant (abnormal) expression levels does not necessarily indicate the absence of a cancerous or pre-cancerous state.

715 C→T Nucleotide Substitution

[0091] Target palladin nucleic acids include palladin nucleic acids that include the 715 C→T substitution; and palladin nucleic acids that are overexpressed in cancerous and precancerous cells (e.g., cancerous and precancerous pancreatic cells). A target palladin nucleic acid that is overexpressed in a cancerous or a precancerous cell (e.g., a cancerous or a precancerous pancreatic cell) may or may not include the 715 C→T substitution. In some embodiments, a target palladin nucleic acid that is overexpressed in a cancerous or precancerous cell (e.g., a cancerous or a precancerous pancreatic cell) is a palladin mRNA that encodes the 90 kD isoform of palladin. In some embodiments, a target palladin nucleic acid that is overexpressed in a cancerous or a precancerous cell (e.g., a cancerous or a precancerous pancreatic cell) includes nucleotide sequences found in exon 2 of the palladin gene, e.g., all or part of the sequence depicted in nucleotides 132 through 803 of the sequence shown in FIG. 1A.

[0092] In some embodiments, a nucleic acid probe hybridizes to and provides for detection of a palladin nucleic acid comprising a nucleotide substitution that is associated with cancer (e.g., pancreatic cancer). In some embodiments, a

nucleotide substitution associated with cancer (e.g., pancreatic cancer) is a C-to-T (C→T) substitution at a location corresponding to nucleotide 715 of SEQ ID NO:1. As depicted in FIG. 1, the underlined and bolded "C" at position 715 is the wild-type sequence; substitution of the C at position 715 with a T is associated with cancer (e.g., pancreatic cancer). In some embodiments, the target nucleic acid is a palladin mRNA, e.g., a C→T substitution at a position corresponding to nucleotide 715 of SEQ ID NO:1 is detected in a palladin mRNA. In other embodiments, the target nucleic acid is a cDNA, e.g., a C→T substitution at a position corresponding to nucleotide 715 of SEQ ID NO:1 is detected in a cDNA copy of a palladin mRNA. In other embodiments, the target nucleic acid is genomic DNA, e.g., a C→T substitution at a position corresponding to nucleotide 715 of SEQ ID NO:1 is detected in palladin genomic DNA, e.g., exon 2 of the palladin gene. For example, a genomic DNA comprising exon 2 sequences and including nucleotide 140301 of the sequence set forth in GenBank Accession No. AC080188 (corresponding to nucleotide 715 of SEQ ID NO:1) is depicted in FIG. 16.

[0093] In some embodiments, the target nucleic acid comprises a nucleotide sequence that has at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 98% nucleotide sequence identity with a stretch of at least about 25, at least about 50, at least about 100, at least about 150, at least about 200, at least about 250, at least about 300, at least about 350, at least about 400, at least about 450, at least about 500, at least about 550, at least about 600, or at least about 670 contiguous nucleotides of nucleotide 132 to nucleotide 803 of SEQ ID NO:1 (or a complement thereof), which is an exon that includes the C→T substitution at position 715.

[0094] In some embodiments, the target nucleic acid comprises a nucleotide sequence of from about nucleotide 132 to about nucleotide 803 of SEQ ID NO:1 (or a complement thereof). In some embodiments, the target nucleic acid comprises from about nucleotide 132 to about nucleotide 802, from about nucleotide 135 to about nucleotide 800, from about nucleotide 140 to about nucleotide 800, from about nucleotide 150 to about nucleotide 800, from about nucleotide 175 to about nucleotide 800, from about nucleotide 200 to about 800, from about nucleotide 250 to about nucleotide 800, from about nucleotide 300 to about nucleotide 800, from about nucleotide 350 to about nucleotide 800, from about nucleotide 400 to about nucleotide 800, from about 450 to about nucleotide 800, from about nucleotide 500 to about nucleotide 800, from about nucleotide 550 to about nucleotide 800, from about nucleotide 600 to about nucleotide 800, from about nucleotide 650 to about nucleotide 800, or from about nucleotide 700 to about nucleotide 800 of SEQ ID NO:1, or the complement of any of the foregoing. In some embodiments, the target nucleic acid will comprise a C at a position corresponding to nucleotide 715 of SEQ ID NO:1; in other embodiments, the target nucleic acid will comprise a T at a position corresponding to nucleotide 715 of SEQ ID NO:1.

[0095] In some embodiments, the target nucleic acid comprises a nucleotide sequence of from about nucleotide 132 to about nucleotide 802 of SEQ ID NO:1 (or a complement thereof). In some embodiments, the target nucleic acid comprises from about nucleotide 132 to about nucleotide 802, from about nucleotide 135 to about nucleotide 750, from about nucleotide 140 to about nucleotide 750, from about nucleotide 150 to about nucleotide 750, from about nucleotide

175 to about nucleotide 750, from about nucleotide 200 to about 750, from about nucleotide 250 to about nucleotide 750, from about nucleotide 300 to about nucleotide 750, from about nucleotide 350 to about nucleotide 750, from about nucleotide 400 to about nucleotide 750, from about 450 to about nucleotide 750, from about nucleotide 500 to about nucleotide 750, from about nucleotide 550 to about nucleotide 750, from about nucleotide 600 to about nucleotide 750, from about nucleotide 650 to about nucleotide 750, from about nucleotide 700 to about nucleotide 750 of SEQ ID NO:1, or the complement of any of the foregoing. In some embodiments, the target nucleic acid will comprise a C at a position corresponding to nucleotide 715 of SEQ ID NO:1; in other embodiments, the target nucleic acid will comprise a T at a position corresponding to nucleotide 715 of SEQ ID NO:1.

[0096] In some embodiments, the target nucleic acid comprises a nucleotide sequence of from about nucleotide 132 to about nucleotide 803 of SEQ ID NO:1 (or a complement thereof). In some embodiments, the target nucleic acid comprises from about nucleotide 132 to about nucleotide 803, from about nucleotide 135 to about nucleotide 725, from about nucleotide 140 to about nucleotide 725, from about nucleotide 150 to about nucleotide 725, from about nucleotide 175 to about nucleotide 725, from about nucleotide 200 to about 725, from about nucleotide 250 to about nucleotide 725, from about nucleotide 300 to about nucleotide 725, from about nucleotide 350 to about nucleotide 725, from about nucleotide 400 to about nucleotide 725, from about 450 to about nucleotide 725, from about nucleotide 500 to about nucleotide 725, from about nucleotide 550 to about nucleotide 725, from about nucleotide 600 to about nucleotide 725, from about nucleotide 650 to about nucleotide 725, from about nucleotide 675 to about nucleotide 725, or from about nucleotide 700 to about nucleotide 725 of SEQ ID NO:1, or the complement of any of the foregoing. In some embodiments, the target nucleic acid will comprise a C at a position corresponding to nucleotide 715 of SEQ ID NO:1; in other embodiments, the target nucleic acid will comprise a T at a position corresponding to nucleotide 715 of SEQ ID NO:1.

[0097] In some embodiments, a suitable target nucleic acid comprises at least the sequence 5'-GACGCCCGCG-3' (SEQ ID NO:2), or a complement thereof, or at least the sequence 5'-GACGICCGCG-3' (SEQ ID NO:3), or a complement thereof, where the bold and underlined nucleotides represent alternative residues at a position corresponding to 715 of SEQ ID NO:1. In some embodiments, a suitable target nucleic acid comprises at least the sequence 5'-GCCACGGCCAGACGCCCGCGGCCTTCCTCA-3' (SEQ ID NO:4), or the complement thereof, or at least the sequence 5'-GCCACG-GCCAGACGTCCGCGGCCTTCCTCA-3' (SEQ ID NO:5), where the bold and underlined nucleotides represent alternative residues at a position corresponding to 715 of SEQ ID NO:1.

[0098] In one non-limiting exemplary embodiment, the target nucleic acid comprises Target Region 1, where Target Region 1 is:

[0099] 5'-ggaccctctgaagctccagcaactcca-gaaccaaatccgactggagcaggaggccggcgcctcggcagcctccgccagc-cggcgagcgcgcggccctcgccccctccgcccgcggccgcttcc-cagactcgcggcctgcacgccg-cggcgtccccggagcccatgagcgcgctggcctcccgtcgcggcccgccatg-cagtcctccgctcctcaactacgcgcgcccaagcagttcatcgccgcgcagaaacctcggcccgctcggccagc-

gcacgccggcctccagcccagctcgtccagcctcc cgtcgcctatgtc-
 cccgacgccgaggcagttcggccgccccctgtccgccccctcgcgagcc
 ctfcgagcctgagcccagggc-
 cccgtggggctcctcctcgcgctcgcggcccccgccccggccaccc cgggtct-
 tcagccccacggctgccttcccgtgc-
 ccgacgtgttcccactgccgccccaccac
 cgccgctcccagcccgggacagggcgtc-
 ccactgctcgtcgcctgccacccgctcggccacgg ccagacg
Tccgcccctcctcagcgtctgtgcctcgcagccggcgc cgccggcgt-
 caac gccctggggctgcccaagggtgtcacccccgc-3' (SEQ ID NO:6;
 corresponding to nucleotides 132-803 of SEQ ID NO:1, but
 including the C→T substitution shown in bold and under-
 lined), or the complement thereof.

[0100] In one non-limiting exemplary embodiment the tar-
 get nucleic acid comprises Target Region 1a, where Target
 Region 1a is:

[0101] 5'-caccgccgctcccagcccgggacag-
 gctcccactgctcgtcgtccacccgctcggcca cgccagacg
Tccgcccctcctcagcgtctgtgcctcgcagccggcgc
 aacgcctgggg-3' (SEQ ID NO:7; corresponding to nucle-
 otides 641-780 of SEQ ID NO:1, but including the C→T
 substitution shown in bold and underlined), or the comple-
 ment thereof.

[0102] In one non-limiting exemplary embodiment, the tar-
 get nucleic acid comprises Target Region 1b, where Target
 Region 1b is:

[0103] 5'-gccttcccgg tcccagcgt gttcccactg ccgcgccac
 caccgcgct cccgagcccg ggacagcgt cccactgctc gtcgctgcc
 accccttcg gccacggcca-3' (SEQ ID NO:8; corresponding to
 nucleotides 601-710 of SEQ ID NO:1), or a complement
 thereof.

[0104] In one non-limiting exemplary embodiment, the tar-
 get nucleic acid comprises Target Region 1c, where Target
 Region 1c is:

[0105] 5'-tggggctcct cctcgcgctc gccccggcc ccgccaccc
 cggtctcag cccacagcgt gccttcccgg tcccagcgt gttcccactg
 ccgccccc acccgcgct cccgagccc-3' (SEQ ID NO:9; corre-
 sponding to nucleotides 541-660 of SEQ ID NO:1), or the
 complement thereof.

[0106] In one non-limiting exemplary embodiment, the tar-
 get nucleic acid comprises Target Region 1d, where Target
 Region 1d is:

[0107] 5'-ggacagcgt cccactgctc gtcgctgcc acccgttcg
 gccacggcca gacgtccgcg gcttctca gcgctctgt gcccgcag
 ccgccccc cgccgctca cgcctgggg-3' (SEQ ID NO:10; cor-
 responding to nucleotides 621-780 of SEQ ID NO:1, but
 including the C→T substitution), or the complement thereof.

[0108] In another non-limiting exemplary embodiment, the
 target nucleic acid comprises Target Region 2, where Target
 Region 2 is:

[0109] 5'-ggaccctctgaagctccagcaactcca-
 gaacaaatccgactggagcaggagccgcccgtcgg cagcctccgcccag-
 ccgcgcagcgcgccctcgcggccccctcccgcccgcggcccgccttc
 ccgagctcgcggcctgacggccc-
 ccgctccccggagcccactgagcgcgtggtccctccgctc cgccccgcatg-
 cagtctcctcgtccttcaactacgcgcccccaagcagttcatcgccgcg
 cagaacctcggcccgcgctcggccacg-
 gcaccccgcctcagccccagctcgtccagcctcc cgtcgcctatgtc-
 ccgagcccagggcagttcggccgccccctcgcgccccctcgcgagcc
 ctfcgagcctgagcccagggc-
 cccgtggggctcctcctcctcgcgctcgcggcccccgccccggccaccc cgggtct-
 tcagccccacggctgccttcccgtgc-
 ccgacgtgttcccactgccgccccaccac
 cgccgctcccagcccgggacagggcgtc-

ccactgctcgtcgcctgccaccgctcggccacgg ccagacg
Cccgcccctcctcagcgtcgtcgtccctcgcagccggccggcgcc
 gtaaac gccctggggctcccaggggtgtcacccccgc-3' (SEQ ID
 NO:11; corresponding to nucleotides 132-803 of SEQ ID
 NO:1), or the complement thereof.

1671 G→T Nucleotide Substitution

[0110] Target palladin nucleic acids include palladin
 nucleic acids that include the 1671 G→T substitution palla-
 din nucleic acids that are abnormally expressed in cancerous
 and precancerous cells (e.g., cancerous and precancerous
 pancreatic cells); and palladin nucleic acids that are abnor-
 mally expressed in non-cancerous cells that are indicative of
 the presence of a cancerous cell. A target palladin nucleic acid
 that is abnormally expressed in a cancerous or a precancerous
 cell (e.g., a cancerous or a precancerous pancreatic cell) or an
 indicator cell may or may not include the 1671 G→T substi-
 tution. In some embodiments, a target palladin nucleic acid
 that is overexpressed in a cancerous or precancerous cell (e.g.,
 a cancerous or a precancerous pancreatic cell) is a palladin
 mRNA that encodes the 90 kD isoform of palladin. In some
 embodiments, a target palladin nucleic acid that is overex-
 pressed in a cancerous or a precancerous cell (e.g., a cancer-
 ous or a precancerous pancreatic cell) includes nucleotide
 sequences found in exon 9 of the palladin gene, e.g., all or part
 of the sequence depicted in nucleotides 1639 through 1846 of
 the sequence shown in FIG. 1A.

[0111] In some embodiments, a nucleic acid probe hybrid-
 izes to and provides for detection of a palladin nucleic acid
 comprising a nucleotide substitution that is associated with
 cancer (e.g., pancreatic cancer). In some embodiments, a
 nucleotide substitution associated with cancer is a G-to-T
 (G→T) substitution at a location corresponding to nucleotide
 1671 of SEQ ID NO:1, and as depicted in FIG. 1A. As
 depicted in FIG. 1A, the underlined and bolded "G" at posi-
 tion 1671 is the wild-type sequence; substitution of the G at
 position 1671 with a T is associated with pancreatic cancer. In
 some embodiments, the target nucleic acid is a palladin
 mRNA, e.g., a G→T substitution at a position corresponding
 to nucleotide 1671 of SEQ ID NO:1 (GenBank AB023209) is
 detected in a palladin mRNA. In other embodiments, the
 target nucleic acid is a cDNA, e.g., a G→T substitution at a
 position corresponding to nucleotide 1671 of SEQ ID NO:1
 (GenBank AB023209) is detected in a cDNA copy of a pal-
 ladin mRNA. In other embodiments, the target nucleic acid is
 genomic DNA, e.g., a G→T substitution at a position corre-
 sponding to nucleotide 1671 of SEQ ID NO:1 is detected in
 palladin genomic DNA, e.g., exon 9 of the palladin gene. For
 example, a genomic DNA comprising exon 9 sequences and
 including nucleotide 183561 of the sequence set forth in
 GenBank Accession No. AC080188 (where nucleotide
 183561 corresponds to nucleotide 1671 of SEQ ID NO:1) is
 depicted in FIG. 1A.

[0112] In some embodiments, the target nucleic acid com-
 prises a nucleotide sequence that has at least about 70%, at
 least about 75%, at least about 80%, at least about 85%, at
 least about 90%, at least about 95%, or at least about 98%
 nucleotide sequence identity with a stretch of at least about
 25, at least about 50, at least about 100, at least about 150, or
 at least about 200 contiguous nucleotides of nucleotide 1639
 to nucleotide 1846 of SEQ ID NO:1 (or a complement
 thereof), which is an exon that includes the G→T substitution
 at position 1671.

[0113] In some embodiments, the target nucleic acid comprises a nucleotide sequence of from about nucleotide 1639 to about nucleotide 1846 of SEQ ID NO:1 (or a complement thereof). In some embodiments, the target nucleic acid comprises from about nucleotide 1639 to about nucleotide 1846, from about nucleotide 1639 to about nucleotide 1840, from about nucleotide 1639 to about nucleotide 1800, from about nucleotide 1639 to about nucleotide 1750, from about nucleotide 1639 to about nucleotide 1700, from about nucleotide 1645 to about nucleotide 1846, from about nucleotide 1650 to about nucleotide 1846, or from about nucleotide 1660 to about nucleotide 1846 of SEQ ID NO:1, or the complement of any of the foregoing. In some embodiments, the target nucleic acid will comprise a G at a position corresponding to nucleotide 1671 of SEQ ID NO:1; in other embodiments, the target nucleic acid will comprise a T at a position corresponding to nucleotide 1671 of SEQ ID NO:1.

[0114] In some embodiments, a suitable target nucleic acid comprises at least the sequence 5'-atctaagctggcaactagat-3' (SEQ ID NO:12), or a complement thereof, or at least the sequence 5'-atctaagctgtcaactagat-3' (SEQ ID NO:13), or a complement thereof, where the bold and underlined nucleotides represent alternative residues at a position corresponding to 1671 of SEQ ID NO:1. In some embodiments, a suitable target nucleic acid comprises at least the sequence 5'-ccaaccccagatctaagctggcaactagatggaaagccc-3' (SEQ ID NO:14), or the complement thereof, or at least the sequence 5'-ccaaccccagatctaagctgtcaactagatggaaagccc-3' (SEQ ID NO:15), where the bold and underlined nucleotides represent alternative residues at a position corresponding to 1671 of SEQ ID NO:1.

[0115] In one non-limiting exemplary embodiment, the target nucleic acid comprises Target Region 3, where Target Region 3 is:

[0116] 5'-gtcagtggttaccacccca-gatctaagctggcaactagatggaaagc-cgtacgcctgacagtgctcacaagatgctggtgcgtgag aacgggggtg-cactctctgatcatagagccagtcacgtcacgtgatgccgcatctacacatgtatagctaccaaccgagcaggacagaact cattcagcctggagctgtgtgtctg (SEQ ID NO:16; corresponding to nucleotides 1639-1846 of SEQ ID NO:1), or the complement thereof.

[0117] In one non-limiting exemplary embodiment, the target nucleic acid comprises Target Region 3a, where Target Region 3a is:

[0118] 5'-gtcagtggttaccacccagatctaagctgtcaactagatggaaagccgtacgcctgacagtgctcacaagatgctggtgcgtgagacgggggtgacactctctgatcatagagc-cagtcacgtcacgtgatgccgcatctacacatgtatagctaccaaccgagcaggacagaactc attcagcctggagctgtgtgtctg (SEQ ID NO:17; corresponding to nucleotides 1639-1846 of SEQ ID NO:1, but including the G→T substitution, shown in bold and underlined), or the complement thereof.

[0119] In one non-limiting exemplary embodiment, the target nucleic acid comprises Target Region 3b, where Target Region 3b is:

[0120] 5'-gtcagtggttaccacccca-gatctaagctggcaactagatggaaagc-cgtacgcctgacagtgctcacaagatgctggtgcgtgag aacgggggtg-cactctctgatcatagagccag (SEQ ID NO:18; corresponding to nucleotides 1639-1760 of SEQ ID NO:1), or the complement thereof.

[0121] In one non-limiting exemplary embodiment, the target nucleic acid comprises Target Region 3c, where Target Region 3c is:

[0122] 5'-gtcagtggttaccacccagatctaagctgtcaactagatggaaagccgtacgcctgacagtgctcacaagatgctggtgcgtgagacgggggtg-cactctctgatcatagagccag (SEQ ID NO:19; corresponding to nucleotides 1639-1760 of SEQ ID NO:1, but including the G→T substitution, shown in bold and underlined), or the complement thereof.

2060 C→T Nucleotide Substitution

[0123] Target palladin nucleic acids include palladin nucleic acids that include the 2060 C→T substitution; and palladin nucleic acids that are overexpressed in cancerous and precancerous cells (e.g., cancerous and precancerous pancreatic cells). A target palladin nucleic acid that is overexpressed in a cancerous or a precancerous cell (e.g., a cancerous or a precancerous pancreatic cell) may or may not include the 2060 C→T substitution. In some embodiments, a target palladin nucleic acid that is overexpressed in a cancerous or precancerous cell (e.g., a cancerous or a precancerous pancreatic cell) is a palladin mRNA that encodes the 140 kD isoform of palladin. In some embodiments, a target palladin nucleic acid that is overexpressed in a cancerous or a precancerous cell (e.g., a cancerous or a precancerous pancreatic cell) includes nucleotide sequences found in exon 10 of the palladin gene, e.g., all or part of the sequence depicted in nucleotides 1833-2175 of the sequence shown in FIGS. 13A and 13B (see, e.g., GenBank Accession No. NM_016081).

[0124] In some embodiments, a nucleic acid probe hybridizes to and provides for detection of a palladin nucleic acid comprising a nucleotide substitution that is associated with cancer, e.g., pancreatic cancer. In some embodiments, a nucleotide substitution associated with cancer is a C-to-T (G→T) substitution at a location corresponding to nucleotide 2060 of SEQ ID NO:20 (GenBank NM_016081, and as depicted in FIGS. 13A and 13B. As depicted in FIGS. 13A and 13B, the underlined and bolded "C" at position 2060 is the wild-type sequence; substitution of the C at position 2060 with a T is associated with cancer. In some embodiments, the target nucleic acid is a palladin mRNA, e.g., a C→T substitution at a position corresponding to nucleotide 2060 of SEQ ID NO:20 (GenBank NM_016081; FIGS. 13A and 13B) is detected in a palladin mRNA. In other embodiments, the target nucleic acid is a cDNA, e.g., a C→T substitution at a position corresponding to nucleotide 2060 of SEQ ID NO:1 (GenBank NM_016081) is detected in a cDNA copy of a palladin mRNA. In other embodiments, the target nucleic acid is genomic DNA, e.g., a C→T substitution at a position corresponding to nucleotide 2060 of SEQ ID NO:20 (GenBank NM_016081) is detected in palladin genomic DNA, e.g., exon 10 of the palladin gene. For example, a genomic DNA comprising nucleotide 27251 of the sequence set forth in GenBank Accession No. AC084353 (corresponding to nucleotide 2060 of the sequence depicted in FIGS. 13A and 13B) is depicted in FIG. 15.

[0125] In some embodiments, the target nucleic acid comprises a nucleotide sequence that has at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 98% nucleotide sequence identity with a stretch of at least about 25, at least about 50, at least about 100, at least about 150, at least about 200, at least about 250, at least about 300, or at least about 340 contiguous nucleotides of nucleotide 1833 to nucleotide 2175 of SEQ ID NO:20 (GenBank NM_016081; or a complement thereof), which is an exon that includes the C→T substitution at position 2060.

[0126] In some embodiments, the target nucleic acid comprises a nucleotide sequence of from about nucleotide 1833 to about nucleotide 2175 of SEQ ID NO:20 (GenBank NM_016081), or a complement thereof. In some embodiments, the target nucleic acid comprises from about nucleotide 1833 to about nucleotide 2090, from about nucleotide 1833 to about nucleotide 2100, from about nucleotide 1833 to about nucleotide 2150, from about nucleotide 1833 to about nucleotide 2160, from about nucleotide 1850 to about nucleotide 2175, from about nucleotide 1900 to about nucleotide 2175, from about nucleotide 1950 to about nucleotide 2175, or from about nucleotide 2000 to about nucleotide 2175 of SEQ ID NO:20 (GenBank NM_016081), or the complement of any of the foregoing. In some embodiments, the target nucleic acid will comprise a C at a position corresponding to nucleotide 2060 of SEQ ID NO:20 (GenBank NM_016081); in other embodiments, the target nucleic acid will comprise a T at a position corresponding to nucleotide 2060 of SEQ ID NO:20 (GenBank NM_016081).

[0127] In some embodiments, a suitable target nucleic acid comprises at least the sequence 5'-cccagcgtggagtaaat-3' (SEQ ID NO:21; GenBank NM_016081), or a complement thereof, or at least the sequence 5'-cccagcgtggagtaaat-3' (SEQ ID NO:22; corresponding to nucleotides 2054-2071 of SEQ ID NO:20, GenBank NM_016081), or a complement thereof, where the bold and underlined nucleotides represent alternative residues at a position corresponding to 2060 of SEQ ID NO:20 (GenBank NM_016081). In some embodiments, a suitable target nucleic acid comprises at least the sequence 5'-ggagtcacccagccgtggagtaaatggactgattaacgg-3' (SEQ ID NO:23; nucleotides 2045-2085 of SEQ ID NO:20, GenBank NM_016081, FIGS. 13A and 13B), or the complement thereof, or at least the sequence 5'-ggagtcacccagcgtggagtaaatggactgattaacgg-3' (SEQ ID NO:24; GenBank NM_016081), where the bold and underlined nucleotides represent alternative residues at a position corresponding to 2060 of SEQ ID NO:20 (GenBank NM_016081).

[0128] In one non-limiting exemplary embodiment, the target nucleic acid comprises Target Region 4, where Target Region 4 is:

[0129] 5'-ccaactgaaaactgtagtacgagtc-
caatgggagaatccaacaatgaccact-
tccaacactttccacctccccccaatcttggagacaa gttccttgagttggct-
tcaagaacaacctctgagatccagcaggtgaacaacctgagttagcctgag
cagggcagccttcaaatgca attcaatgctgctgagagggaac-
gaacggagtcacccagc
cgtggagtaaatggactgattaacggcaagctaacagtaataaatct ctcca-
caccagctgtcctgcttccaccac-
taaggagccaccactctgctgccaacccaaact (SEQ ID NO:25; cor-
responding to nucleotides 1833-2175 of SEQ ID NO:25
(GenBank NM_016081)), or the complement thereof.

[0130] In one non-limiting exemplary embodiment, the target nucleic acid comprises Target Region 4a, where Target Region 4a is:

[0131] 5'-ccaactgaaaactgtagtacgagtc-
caatgggagaatccaacaatgaccact-
tccaacactttccacctccccccaatcttggagacaa gttccttgagttggct-
tcaagaacaacctctgagatccagcaggtgaacaacctgagttagcctgag
cagggcagccttcaaatgca attcaatgctgctgagagggaac-
gaacggagtcacccagc
tgtggagtaaatggactgattaacggcaagctaacagtaataaatct ctcca-
caccagctgtcctgcttccaccac-
taaggagccaccactctgctgccaacccaaact (SEQ ID NO:26; cor-
responding to nucleotides 1833-2175 of SEQ ID NO:20
(GenBank NM_016081), but including the C→T substitui-

tion at a position corresponding to nucleotide 2060 of NM_016801), or the complement thereof.

[0132] In one non-limiting exemplary embodiment, the target nucleic acid comprises Target Region 4b, where Target Region 4b is:

[0133] 5'-ttcaatgctgctgagagggaacgaacg-
gagtcacatcccagccgtggagtaaatg-
gactgattaacggcaagctaacagtaataaatctctt (SEQ ID NO:27; cor-
responding to nucleotides 2018 to 2110 of SEQ ID NO:20
(GenBank NM_016081), or the complement thereof.

[0134] In one non-limiting exemplary embodiment, the target nucleic acid comprises Target Region 4c, where Target Region 4c is:

[0135] 5'-ttcaatgctgctgagagggaacgaacg-
gagtcacatcccagctgtggagtaaatg-
gactgattaacggcaagctaacagtaataaatctct t (SEQ ID NO:28;
corresponding to nucleotides 2018 to 2110 of SEQ ID NO:20
(GenBank NM_016081) but including the C→T substitution
at a position corresponding to nucleotide 2060 of
NM_016801), or the complement thereof.

Insertions in a Palladin Promoter

[0136] Target palladin nucleic acids include palladin nucleic acids that include a palladin gene promoter that includes a nucleotide sequence insertion, e.g., an insertion of one or more nucleotides not present in a palladin gene promoter present in a normal (non-cancerous) cell. Thus, the insertion in the palladin gene promoter is associated with an increased risk of developing cancer (e.g., pancreatic cancer), and/or the presence of a pre-cancerous or cancerous cell (e.g., a pre-cancerous or cancerous pancreatic cell).

[0137] In some embodiments, a nucleic acid probe hybridizes to a region of a palladin gene promoter that includes a nucleotide sequence insertion, e.g., an insertion of 1 nt to 5 nt, from 5 nt to 10 nt, from 10 nt to 15 nt, from 15 nt to 20 nt, from 20 nt to 25 nt, from 25 nt to 30 nt, from 30 nt to 35 nt, from 35 nt to 40 nt, from 40 nt to 50 nt, or more than 50 nt.

[0138] A target nucleic acid can include a palladin gene promoter of any isoform of palladin. For example, in some embodiments, the target nucleic acid is a promoter region of the palladin gene that encodes the 90 kDa isoform of palladin. FIG. 19 depicts a nucleotide sequence (SEQ ID NO:110) that includes a promoter for the gene encoding the 90 kDa isoform of palladin, where the promoter region is boxed (e.g., nucleotides 501-1987 of SEQ ID NO:110). The promoter region includes a 12-base pair insertion after nucleotide 704 of SEQ ID NO:110. The 12-bp insertion is associated with an increased risk of developing cancer and/or the presence of a pre-cancerous or cancerous cell. A promoter without the 12-base pair insertion would include nucleotides 501-704 and 717-1987 of SEQ ID NO:110. The 12-base pair insertion is nucleotides 705-716 of SEQ ID NO:110. An example of a promoter region of a palladin gene encoding the 90 kDa isoform of palladin, without the 12-base pair insertion, is SEQ ID NO:111, which includes nucleotides 501-704 and 717-1987 of SEQ ID NO:110.

[0139] In some embodiments, a target nucleic acid comprises a nucleotide sequence that has at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 98% nucleotide sequence identity with a stretch of at least about 25, at least about 50, at least about 100, at least about 150, at least about 200, at least about 250, at least about 500, at least about 750, at least about 1000, at least about 1250, or at least

about 1475 contiguous nucleotides of the nucleotide sequence set forth in SEQ ID NO:111. SEQ ID NO:111 provides a nucleotide sequence of a palladin promoter without nucleotide sequence insertions.

[0140] In some embodiments, a target nucleic acid comprises a nucleotide sequence that has at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or at least about 98% nucleotide sequence identity with a stretch of at least about 25, at least about 50, at least about 100, at least about 150, at least about 200, at least about 250, at least about 500, at least about 750, at least about 1000, at least about 1250, or at least 1487 contiguous nucleotides of the nucleotide sequence set forth in nucleotides 501-1987 of SEQ ID NO:110 (see FIG. 19). In some embodiments, a target nucleic acid includes a nucleotide sequence insertion, compared to the sequence set forth in SEQ ID NO:111. For example, in some embodiments, the target nucleic acid includes a nucleotide sequence insertion at a position corresponding to nucleotide 704 of the sequence set forth in SEQ ID NO:111.

[0141] In one non-limiting exemplary embodiment, a target nucleic acid comprises the nucleotide sequence of Target Region 5, where Target Region 5 is the nucleotide sequence set forth in SEQ ID NO:111.

[0142] In another non-limiting exemplary embodiment, a target nucleic acid comprises the nucleotide sequence of Target Region 5a, where Target Region 5a has the sequence:

[0143] 5'-CGGGCGAGGTATAAAGCCCGATACCTGCC-
CCCGCGCCCGGTCCGCGGAGCCCGCTG CAGCTC-
CCGCTCGCTCCGGACGCG-
GAATCGGCGCAGCAGCGGGAGGCGGCCCGGAG
AGCCGAGG-3' (SEQ ID NO:112), where the sequence corresponds to nucleotides 658-775 of the nucleotide sequence depicted in FIG. 19 and set forth in SEQ ID NO:110, and where the 12-base pair insertion is in bold; or the complement of the sequence.

Sources of Target Nucleic Acids

[0144] Where the detection methods involve detection of a palladin-encoding nucleic acid, the target nucleic acids are detected in samples obtained from a tissue comprising cells. In some embodiments, the cells are obtained from a tissue suspected of comprising cancer cells. For example, in some embodiments, the cells are obtained from pancreatic tissue. As another example, the cells are obtained from breast tissue and/or axillary lymph nodes. As yet another example, the cells are obtained from the upper aerodigestive tract (including cells, e.g., squamous cells, obtained from the lips, mouth, tongue, nose, throat, larynx, pharynx, and upper trachea; but not including cells obtained from the stomach or intestines), and tissues such as cervical lymph nodes. In other embodiments, the cells are indicator cells, e.g., the cells themselves are not suspected of comprising cancer cells, but serve as an indication of the presence of cancer cells in an individual. Such indicator cells include, e.g., white blood cells. For example, peripheral blood mononuclear cells are suitable indicator cells.

[0145] For example, target nucleic acids can be detected in samples obtained from pancreatic cells, particularly human pancreatic cells. Pancreatic cells suitable for analysis include, but are not limited to, ductal pancreatic epithelial cells; pancreatic acinar cells; pancreatic stroma and/or extra-cellular matrix; pancreatic stem cells; and pancreatic islet cells. In addition, other targets for palladin detection can include, but

are not limited to, biologic sources of tissue including peripheral white blood cells and pancreatic juice.

[0146] Head and neck cancers are malignant growths originating in the lip and oral cavity, nasal cavity, pharynx, larynx, thyroid, paranasal sinuses, salivary glands and cervical lymph nodes of the neck. Squamous cell carcinomas represent more than 90% of all head and neck cancers, and originate from the squamous cells that line the upper aerodigestive tract. Suitable sources of target nucleic acids in the context of head and neck squamous cell carcinoma include tissues obtained from any of: the lip, the oral cavity (mouth), nasal cavity, pharynx, larynx, thyroid, paranasal sinuses, salivary glands, and cervical lymph nodes of the neck. Suitable sources of target nucleic acids include a tissue sample from any of the aforementioned sources, where the tissue sample includes squamous cells. Suitable sources of target nucleic acids include cells obtained by oral rinse; oral swab; and tissue biopsy from any of the aforementioned sites.

[0147] In the context of breast cancer, suitable sources of target nucleic acids include breast cells and lymph node cells, e.g., cells obtained via fine needle aspiration biopsy; cells obtained via core needle biopsy; cells obtained from lymph nodes in the vicinity of the breast (e.g., axillary lymph nodes); and the like.

[0148] In the context of breast cancer, breast cancers that can be detected using a subject method include mammary carcinoma, adenocarcinoma, ductal carcinoma in situ, lobular (small cell) carcinoma, intraductal carcinoma, medullary breast cancer, mucinous breast cancer, tubular breast cancer, papillary breast cancer, Paget's disease, inflammatory breast cancer, and hormone dependent tumors of the breast.

Nucleic Acid Probes

[0149] As discussed above, the present invention provides detection, diagnostic, and staging methods, e.g., methods for detecting and diagnosing cancer (e.g., pancreatic cancer, breast cancer, head and neck squamous cell cancer, etc.) in an individual; methods of identifying individuals at risk of developing cancer (e.g., pancreatic cancer); and methods of staging cancer (e.g., pancreatic cancer, breast cancer, head and neck squamous cell cancer, etc.). The methods generally involve detecting a nucleotide sequence alteration (e.g., nucleotide substitution, a nucleotide sequence insertion, a nucleotide sequence deletion, a change in methylation status, etc.), a level of palladin mRNA, or both. In aspects where detection of the presence of palladin gene having a cancer-associated nucleotide substitution is desired, a subject method provides for detection one of more of the following exemplary nucleotide substitutions: 1) a C→T nucleotide substitution at a position corresponding to nucleotide 715 ("a 715 C→T substitution") of a human palladin nucleic acid, e.g., nucleotide 715 of a palladin nucleic acid as set forth in GenBank Accession No. AB023209 and depicted in FIG. 1A, where nucleotide 715 is underlined and in bold text; 2) a G→T nucleotide substitution at a position corresponding to nucleotide 1671 ("a 1671 G→T substitution") of a human palladin nucleic acid, e.g., nucleotide 1671 of a palladin nucleic acid as set forth in GenBank Accession No. AB023209 and depicted in FIG. 1A, where nucleotide 1671 is underlined and in bold text; 3) a C→T substitution at a position corresponding to nucleotide 2060 ("a 2060 C→T substitution") of a human palladin nucleic acid, e.g., a nucleotide 2060 of a palladin nucleic acid as set forth in GenBank Accession No. NM_016081 and depicted in FIGS. 13A and 13B,

where nucleotide 2060 is underlined and in bold text. In aspects where detection of the presence of palladin gene promoter having a cancer-associated nucleotide sequence insertion is desired, a subject method provides for detection of, e.g., a nucleotide insertion in a promoter of a palladin gene encoding palladin, e.g., encoding the 90 kDa isoform of palladin. In other aspects, a subject method provides for detection of a level of a palladin mRNA in a cell. Both of these methods (detecting a nucleotide sequence alteration; and detecting a palladin mRNA level) can be carried out using a method involving nucleic acid hybridization, amplification, or both.

[0150] Nucleic acid hybridization can be carried out using a nucleic acid probe that detects: a) a cancer-associated nucleotide sequence alteration in a palladin target nucleic acid; b) a level of a palladin mRNA that is abnormally expressed in a cancerous or precancerous cell; or c) both a cancer-associated nucleotide sequence alteration in a palladin target nucleic acid and a level of a palladin mRNA that is abnormally expressed in a cancerous or precancerous cell.

[0151] For example, nucleic acid hybridization can be carried out using a nucleic acid probe that detects: a) a pancreatic cancer-associated nucleotide sequence alteration in a palladin target nucleic acid; b) a level of a palladin mRNA that is abnormally expressed in a cancerous or precancerous pancreatic cell; or c) both a pancreatic cancer-associated nucleotide sequence alteration in a palladin target nucleic acid and a level of a palladin mRNA that is abnormally expressed in a cancerous or precancerous pancreatic cell.

[0152] For example, nucleic acid hybridization can be carried out using a nucleic acid probe that detects: a) a cancer-associated nucleotide substitution (e.g., a defined by a 715 C→T substitution, a 1671 G→T substitution, or a 2060 C→T substitution in a palladin target nucleic acid); b) a level of a palladin mRNA that is overexpressed in a cancerous or precancerous cell; or c) both a 715 C→T substitution a 1671 G→T substitution, or a 2060 C→T substitution in a palladin target nucleic acid and a level of a palladin mRNA that is overexpressed in a cancerous or precancerous cell. In addition, nucleic acid hybridization can be used to detect a nucleotide sequence insertion in a palladin promoter. The following is a description of exemplary probes that can be used in a subject method.

[0153] Where the method involves detection of a cancer-associated nucleotide sequence alteration (e.g., a nucleotide substitution, a nucleotide sequence insertion, a nucleotide sequence deletion, a change in methylation status, etc.) in palladin, suitable nucleic acid probes include nucleic acid probes that hybridize to and provide for detection of a palladin nucleic acid comprising a nucleotide substitution that is associated with cancer, e.g., a palladin 715 C→T substitution, a 1671 G→T substitution, or a 2060 C→T substitution; and nucleic acid probes that hybridize to and provide for detection of a palladin nucleic acid comprising a nucleotide sequence insertion that is associated with cancer. Where abnormal expression of palladin is to be detected, nucleic acid probes suitable for use include nucleic acid probes that hybridize to and provide for detection of a palladin nucleic acid that is overexpressed or underexpressed in an indicator cell, a cancerous cell, or a precancerous cell. The present invention provides such nucleic acid probes.

[0154] Suitable nucleic acid probes are in some embodiments in the range of between 10-50 nucleotides long, such as 10 to 50, 12 to 45, 15 to 40, 20 to 35, 25 to 30 nucleotides, and

the like. For example, probes will in some embodiments be in the range of between 18 to 40, 19 to 35, 20 to 30, 21 to 29, 22 to 28, 23 to 27, 24-25 nucleotides long, and any length between the stated ranges. Probes of about 20 to 22 nucleotides in length are of particular interest in some embodiments.

[0155] A suitable probe may be coupled to a label for detection. There are several methods and compositions known for derivatizing oligonucleotides with reactive functionalities which permit the addition of a label. For example, several approaches are available for biotinylating probes so that radioactive, fluorescent, chemiluminescent, enzymatic, or electron dense labels can be attached via avidin. See, e.g., Broken et al., *Nucl. Acids Res.* (1978) 5:363-384 which discloses the use of ferritin-avidin-biotin labels; and Chollet et al. *Nucl. Acids Res.* (1985) 13:1529-1541 which discloses biotinylation of the 5' termini of oligonucleotides via an aminoalkylphosphoramidate linker arm. Several methods are also available for synthesizing amino-derivatized oligonucleotides which are readily labeled by fluorescent or other types of compounds derivatized by amino-reactive groups, such as isothiocyanate, N-hydroxysuccinimide, or the like, see, e.g., Connolly (1987) *Nucl. Acids Res.* 15:3131-3139, Gibson et al. (1987) *Nucl. Acids Res.* 15:6455-6467 and U.S. Pat. No. 4,605,735 to Miyoshi et al. Methods are also available for synthesizing sulfhydryl-derivatized oligonucleotides which can be reacted with thiol-specific labels, see, e.g., U.S. Pat. No. 4,757,141 to Fung et al., Connolly et al. (1985) *Nucl. Acids Res.* 13:4485-4502 and Spoot et al. (1987) *Nucl. Acids Res.* 15:4837-4848. A comprehensive review of methodologies for labeling DNA fragments is provided in Matthews et al., *Anal. Biochem.* (1988) 169:1-25.

[0156] For example, probes may be fluorescently labeled by linking a fluorescent molecule to the non-ligating terminus of the probe. Guidance for selecting appropriate fluorescent labels can be found in Smith et al., *Meth. Enzymol.* (1987) 155:260-301; Karger et al., *Nucl. Acids Res.* (1991) 19:4955-4962; Haugland (1989) *Handbook of Fluorescent Probes and Research Chemicals* (Molecular Probes, Inc., Eugene, Ore.). Exemplary fluorescent labels include fluorescein and derivatives thereof, such as disclosed in U.S. Pat. No. 4,318,846 and Lee et al., *Cytometry* (1989) 10:151-164, and 6-FAM, JOE, TAMRA, ROX, HEX-1, HEX-2, ZOE, TET-1 or NAN-2, and the like.

[0157] Additionally, probes can be labeled with an acridinium ester (AE). Current technologies allow the AE label to be placed at any location within the probe. See, e.g., Nelson et al. (1995) "Detection of Acridinium Esters by Chemiluminescence" in *Nonisotopic Probing, Blotting and Sequencing*, Kricka L. J. (ed) Academic Press, San Diego, Calif.; Nelson et al. (1994) "Application of the Hybridization Protection Assay (HPA) to PCR" in *The Polymerase Chain Reaction*, Mullis et al. (eds.) Birkhauser, Boston, Mass.; Weeks et al., *Clin. Chem.* (1983) 29:1474-1479; Berry et al., *Clin. Chem.* (1988) 34:2087-2090. An AE molecule can be directly attached to the probe using non-nucleotide-based linker arm chemistry that allows placement of the label at any location within the probe. See, e.g., U.S. Pat. Nos. 5,585,481 and 5,185,439.

[0158] If a solid support is used in the assay (e.g., to capture amplicons of target nucleic acid using a probe), the oligonucleotide probe may be attached to the solid support in a variety of manners. For example, the probe may be attached to the solid support by attachment of the 3' or 5' terminal nucleotide of the probe to the solid support. In some embodiments,

the probe is attached to the solid support by a linker which serves to distance the probe from the solid support. The linker is in many embodiments at least 15-30 atoms in length, or at least 15-50 atoms in length. The required length of the linker will depend on the particular solid support used. For example, a six atom linker is generally sufficient when high cross-linked polystyrene is used as the solid support.

[0159] A wide variety of linkers are known in the art which may be used to attach the oligonucleotide probe to the solid support. The linker may be formed of any compound which does not significantly interfere with the hybridization of the target sequence to the probe attached to the solid support. The linker may be formed of a homopolymeric oligonucleotide which can be readily added on to the linker by automated synthesis. Alternatively, polymers such as functionalized polyethylene glycol can be used as the linker. In some embodiments, polymers such as functionalized polyethylene glycol are used because they do not significantly interfere with the hybridization of probe to the target oligonucleotide. In some embodiments, the linker is polyethylene glycol.

[0160] The linkages between the solid support, the linker and the probe are normally not cleaved during removal of base protecting groups under basic conditions at high temperature. Examples of suitable linkages include carbamate and amide linkages.

[0161] Examples of suitable types of solid supports for immobilization of the oligonucleotide probe include controlled pore glass, glass plates, polystyrene, avidin-coated polystyrene beads, cellulose, nylon, acrylamide gel and activated dextran.

[0162] Probes for use in detection of a pancreatic cancer-associated nucleotide substitution in a palladin gene, and probes useful in assessing palladin expression levels are described in more detail below.

[0163] Probes for Detecting a Cancer-Associated Nucleotide Substitutions in Palladin

[0164] In some embodiments, a suitable nucleic acid probe hybridizes to and provides for detection of a palladin nucleic acid (e.g., genomic DNA; mRNA; cDNA; amplified copies of any of the foregoing; etc.) comprising a nucleotide substitution associated with cancer (e.g., pancreatic cancer, breast cancer, head and neck squamous cell cancer, etc.).

Probes for Detecting a 715 C→T Substitution

[0165] Non-limiting examples of nucleic acid probes that hybridize to a nucleic acid comprising Target Region 1, and that detect the presence of the C→T substitution depicted in Target Region 1 include:

[0166] Probe 1: 5'-CGGCCAGACGTCCGCGGCCTT-3' (SEQ ID NO:29), or the complement thereof, e.g., 5'-AAGGCCGCGACGTCTGGCCG-3' (SEQ ID NO:30);

[0167] Probe 2: 5'-GCCACGGCCAGACGTCCGCGGCCTT-3' (SEQ ID NO:31), or the complement thereof, e.g., 5'-AAGGCCGCGACGTCTGGCCGTGGC-3' (SEQ ID NO:32);

[0168] Probe 3: 5'-CGGCCAGACGTCCGCGGCCTTCTC-3' (SEQ ID NO:33), or the complement thereof, e.g., 5'-GAGGAAGGCCGCGGACGTCTGGCCGTGGC-3' (SEQ ID NO:34); and

[0169] Probe 4: 5'-GCCACGGCCAGACGTCCGCGGCCTTCTC-3' (SEQ ID NO:35), or the complement thereof, e.g., 5'-GAGGAAGGCCGCGGACGTCTGGCCGTGGC-3' (SEQ ID NO:36).

[0170] Any of Probes 1-4, above, or any of the complements thereof, will in some embodiments include additional 5' and/or 3' sequences. The additional 5' and/or 3' sequences will in some embodiments provide for hybridization with the target region. In other embodiments, the additional 5' and/or 3' sequences will provide restriction endonuclease recognition sites, for cloning the probe into a vector.

Probes for Detecting a 1671 G→T Substitution

[0171] Non-limiting examples of nucleic acid probes that hybridize to a nucleic acid comprising Target Region 3, and that detect the presence of the G→T substitution depicted in Target Region 3 include:

[0172] Probe 5: 5'-ctaagctggcaactagatgg-3' (SEQ ID NO:37), or the complement thereof, e.g., 5'-ccatctagtgc-cagcttag-3' (SEQ ID NO:38);

[0173] Probe 6: 5'-gatctaagctggcaactagat-3' (SEQ ID NO:39), or the complement thereof, e.g., 5'-atcattgtgccagcttagatc-3' (SEQ ID NO:40); and

[0174] Probe 7: 5'-gatctaagctggcaactagatgg-3' (SEQ ID NO:41), or the complement thereof, e.g., 5'-ccatctagtgc-cagcttagatc-3' (SEQ ID NO:42).

[0175] Any of Probes 5-7, above, or any of the complements thereof, will in some embodiments include additional 5' and/or 3' sequences. The additional 5' and/or 3' sequences will in some embodiments provide for hybridization with the target region. In other embodiments, the additional 5' and/or 3' sequences will provide restriction endonuclease recognition sites, for cloning the probe into a vector.

Probes for Detecting a 2060 C→T Substitution

[0176] Non-limiting examples of nucleic acid probes that hybridize to a nucleic acid comprising Target Region 4, and that detect the presence of the C→T substitution depicted in Target Region 4 include:

[0177] Probe 8: 5'-ccatcccagccgtggagtaaatgg (SEQ ID NO:43), or the complement thereof, e.g., 5'-ccattactccacg-gctgggatgg-3' (SEQ ID NO:44);

[0178] Probe 9: 5'-cccagccgtggagtaaatgg-3' (SEQ ID NO:45), or the complement thereof, e.g., 5'-ccattactccacg-gctggg-3' (SEQ ID NO:46); and

[0179] Probe 10: 5'-gtccatcccagccgtggagtaaatggactg-3' (SEQ ID NO:47), or the complement thereof, e.g., 5'-cagtc-cattactccacgctgggatggac-3' (SEQ ID NO:48).

[0180] Any of Probes 8-10, above, or any of the complements thereof, will in some embodiments include additional 5' and/or 3' sequences. The additional 5' and/or 3' sequences will in some embodiments provide for hybridization with the target region. In other embodiments, the additional 5' and/or 3' sequences will provide restriction endonuclease recognition sites, for cloning the probe into a vector.

[0181] Probes for Detecting Abnormal Expression of a Palladin Nucleic Acid

[0182] In some embodiments, a suitable nucleic acid probe hybridizes to and provides for detection of a palladin nucleic acid that is abnormally expressed (e.g., over-expressed or under-expressed) in a cell (e.g., a cell from a tissue suspected of comprising a cancer cell; or an indicator cell such as a white blood cell). In these embodiments, suitable target nucleic acids include a palladin mRNA; a cDNA copy of all or a portion of (e.g., a subsequence of) a palladin mRNA; a nucleic acid amplification product generated by amplifying a cDNA copy of a palladin mRNA; and the like. In some

embodiments, a target nucleic acid will include all or a part of an exon (e.g., exon 2, exon 9, exon 10) of the palladin gene, and may or may not include a nucleotide substitution that is associated with pancreatic cancer. In some of these embodiments, a nucleic acid probe will comprise a variable nucleotide that is associated with pancreatic cancer (e.g., will comprise one or more of: a) a nucleotide sequence that includes either a C or a T at a position corresponding to 715 of SEQ ID NO:1; b) a nucleotide sequence that includes either a G or a T at a position corresponding to 1671 of SEQ ID NO:1; and c) a nucleotide sequence that includes either a C or a T at a position corresponding to 2060 of the sequence set forth in FIGS. 13A and 13B and in NM_016801). In other embodiments, a nucleic acid probe will not include a nucleotide sequence that includes any of the aforementioned nucleotides (e.g., a) a nucleotide sequence that includes either a C or a T at a position corresponding to 715 of SEQ ID NO:1; b) a nucleotide sequence that includes either a G or a T at a position corresponding to 1671 of SEQ ID NO:1; and c) a nucleotide sequence that includes either a C or a T at a position corresponding to 2060 of the sequence set forth in FIGS. 13A and 13B and in NM_016801).

[0183] Exemplary, non-limiting examples of probes that are suitable for detecting a palladin nucleic acid that is abnormally expressed in a cell include any of Probes 1-10, or a complement thereof, as described above.

[0184] Further suitable probes for detecting a palladin nucleic acid that is abnormally expressed in a cell include, e.g.:

[0185] Probe 11: 5'-accgcttcg gccacggcca-3' (SEQ ID NO:49; corresponding to nucleotides 691-710 of SEQ ID NO:1), or the complement thereof;

[0186] Probe 12: 5'-gtcgcctgcc accgcttcg gccacggc-3' (SEQ ID NO:50; corresponding to nucleotides 681-708 of SEQ ID NO:1) or the complement thereof;

[0187] Probe 13: 5'-gccttctca gcctctgct-3' (SEQ ID NO:51; corresponding to nucleotides 721-740 of SEQ ID NO:1), or the complement thereof;

[0188] Probe 14: 5'-gccttctca gcctctgct gccctgcag-3' (SEQ ID NO:52; corresponding to nucleotides 721-750 of SEQ ID NO:1), or the complement thereof;

[0189] Probe 15: 5'-gcctctgct gccctgcag cggc-3' (SEQ ID NO:53; corresponding to nucleotides 731-755 of SEQ ID NO:1), or the complement thereof;

[0190] Probe 16: 5'-gccctgcag ccgcccgg cggcgtcaa-3' (SEQ ID NO:54; corresponding to nucleotides 741-770 of SEQ ID NO:1), or the complement thereof;

[0191] Probe 17: 5'-cagtgggta ccaacccag-3' (SEQ ID NO:55; corresponding to nucleotides 1641-1660 of SEQ ID NO:1), or the complement thereof;

[0192] Probe 18: 5'-ccaacccag atctaagctg-3' (SEQ ID NO:56; corresponding to nucleotides 1651-1670 of SEQ ID NO:1), or the complement thereof;

[0193] Probe 19: 5'-ggaagccc tacgacctga-3' (SEQ ID NO:57; corresponding to nucleotides 1681-1700 of SEQ ID NO:1), or the complement thereof;

[0194] Probe 20: 5'-cagtgetcac aagatgctgg-3' (SEQ ID NO:58; corresponding to nucleotides 1701-1720 of SEQ ID NO:1), or the complement thereof;

[0195] Probe 21: 5'-aatgctgctg agaggaaac (SEQ ID NO:59; corresponding to nucleotides 2021-2040 of the sequence set forth in FIGS. 13A and 13B (NM_016801), or the complement thereof;

[0196] Probe 22: 5'-agaggaaac gaacggagtc catcc-3' (SEQ ID NO:60; corresponding to nucleotides 2031-2055 of the sequence set forth in FIGS. 13A and 13B (NM_016801), or the complement thereof;

[0197] Probe 23: 5'-gtggagtaaa tggactgatt aacggcaaag-3' (SEQ ID NO:61; corresponding to nucleotides 2061-2090 of the sequence set forth in FIGS. 13A and 13B (NM_016801), or the complement thereof; and

[0198] Probe 24: 5'-gtaaa tggactgatt aacggc-3' (SEQ ID NO:62; corresponding to nucleotides 2066-2086 of the sequence set forth in FIGS. 13A and 13B (NM_016801), or the complement thereof.

[0199] Any of Probes 11-16, above, or any of the complements thereof, is suitable for detecting a palladin nucleic acid comprising all or a portion of Target Region 1, as described above. Any of Probes 17-20, above, or any of the complements thereof, is suitable for detecting a palladin nucleic acid comprising all or a portion of Target Region 3, as described above. Any of Probes 21-24, above, or any of the complements thereof, is suitable for detecting a palladin nucleic acid comprising all or a portion of Target Region 4, as described above.

[0200] Any of Probes 11-24, above, or any of the complements thereof, will in some embodiments include additional 5' and/or 3' sequences. The additional 5' and/or 3' sequences will in some embodiments provide for hybridization with the target region. In other embodiments, the additional 5' and/or 3' sequences will provide restriction endonuclease recognition sites, for cloning the probe into a vector.

Probes for Detecting a Cancer-Associated Nucleotide Sequence Insertions in a Palladin Nucleic Acid

[0201] In some embodiments, a suitable nucleic acid probe hybridizes to and provides for detection of a palladin nucleic acid (e.g., genomic DNA; mRNA; cDNA; amplified copies of any of the foregoing; etc.) comprising a nucleotide sequence insertion associated with cancer (e.g., pancreatic cancer, breast cancer, head and neck squamous cell cancer, etc.).

[0202] For example, in some embodiments, a suitable nucleic acid probe hybridizes to and provides for detection of a palladin nucleic acid (e.g., a genomic DNA) comprising a nucleotide sequence insertion in a palladin promoter. In a particular non-limiting embodiment, a suitable nucleic acid probe hybridizes to and provides for detection of a palladin nucleic acid (e.g., a genomic DNA) comprising a nucleotide sequence insertion in a promoter of a palladin gene encoding the 90 kDa isoform of palladin.

[0203] For example, in some embodiments, a suitable nucleic acid probe hybridizes to and provides for detection of a nucleotide sequence insertion in the nucleotide sequence set forth in SEQ ID NO:111. A 12-base pair insertion was found in a palladin gene promoter, and is shown in FIG. 19. In some embodiments, a suitable nucleic acid probe hybridizes to and provides for detection of this 12-base pair insertion.

[0204] Non-limiting examples of a suitable probe include the following:

[0205] Probe 25: 5'-GCGGAGCCCGCTGCAGCTCCCG-3' (SEQ ID NO:113), corresponding to nucleotides 700-721 of the nucleotide sequence depicted in FIG. 19 and set forth in SEQ ID NO:110.

[0206] Probe 26: 5'-CGGAGCCCGCTGCAGCTCCC-3' (SEQ ID NO:114), corresponding to nucleotides 701-720 of the nucleotide sequence depicted in FIG. 19 and set forth in SEQ ID NO:10.

[0207] Probes 25 and 26, above, or any of the complements thereof, are suitable for detecting a palladin nucleic acid comprising all or a portion of Target Region 5a, as described above.

Nucleic Acid Primers

[0208] As discussed above, a subject method (e.g., a method for detecting and diagnosing cancer in an individual; a method of identifying individuals at risk of developing cancer; a method of staging cancer) will in some embodiments involve amplification of a target palladin nucleic acid. For example, a subject method that involves detection of a 715 C→T substitution, a 1671 G→T, or a 2060 C→T substitution in a palladin nucleic acid, or detection of a nucleotide substitution in a palladin gene, will in some embodiments comprise amplifying a target palladin nucleic acid, using a nucleic acid primer pair, forming a nucleic acid amplification product; and detecting the 715 C→T substitution, the 1671 G→T, or the 2060 C→T substitution, or an insertion in a palladin gene promoter, in the amplification product. In other aspects, a subject method provides for detection of a level of a palladin mRNA in a cell. In some embodiments, methods of detecting a level of a palladin mRNA in a cell will involve amplifying a target palladin nucleic acid, using a pair of nucleic acid primers.

[0209] As noted above, the present invention provides methods that involve detecting a palladin nucleic acid (e.g., a palladin mRNA) that is over-expressed in an indicator cell which is a cancerous or pre-cancerous cell. A palladin nucleic acid that is over-expressed in a cancerous tissue is a palladin nucleic acid (e.g., a palladin mRNA) that is present in a cancerous cell at a level that is at least about 25%, at least about 50%, at least about 2-fold, at least about 2.5-fold, at least about 5-fold, at least about 7.5-fold, at least about 10-fold, at least about 15-fold, at least about 20-fold, at least about 25-fold, at least about 30-fold, at least about 35-fold, at least about 40-fold, at least about 50-fold, or more, higher than the level of palladin nucleic acid (e.g., palladin mRNA) present in a control cell, e.g., a non-cancerous cell of the same cell type.

[0210] For example, the present invention provides methods that involve detecting a palladin nucleic acid (e.g., a palladin mRNA) that is over-expressed in a cancerous or pre-cancerous pancreatic cell. A palladin nucleic acid that is over-expressed in a cancerous pancreatic tissue is a palladin nucleic acid (e.g., a palladin mRNA) that is present in a cancerous pancreatic cell at a level that is at least about 25%, at least about 50%, at least about 2-fold, at least about 2.5-fold, at least about 5-fold, at least about 7.5-fold, at least about 10-fold, at least about 15-fold, at least about 20-fold, at least about 25-fold, at least about 30-fold, at least about 35-fold, at least about 40-fold, at least about 50-fold, or more, higher than the level of palladin nucleic acid (e.g., palladin mRNA) present in a control cell, e.g., a non-cancerous pancreatic cell of the same cell type.

[0211] A palladin nucleic acid that is over-expressed in a pre-cancerous pancreatic tissue is a palladin nucleic acid (e.g., a palladin mRNA) that is present in a pre-cancerous pancreatic cell at a level that is at least about 25%, at least about 50%, at least about 2-fold, at least about 2.5-fold, at least about 5-fold, at least about 7.5-fold, at least about 10-fold, at least about 15-fold, at least about 20-fold, at least about 25-fold, at least about 30-fold, at least about 35-fold, at least about 40-fold, at least about 50-fold, or more, higher

than the level of palladin nucleic acid (e.g., palladin mRNA) present in a control cell, e.g., a non-cancerous pancreatic cell of the same cell type.

[0212] As noted above, in some embodiments, the methods provide for detection of a cancer in an individual, where a palladin target nucleic acid is detected in an indicator cell in a biological sample obtained from the individual, and where a level of palladin target nucleic acid that is lower than normal indicates the presence of cancer in the individual. A palladin target nucleic acid that is expressed at lower than normal levels is expressed at a level that is less than about 80%, less than about 70%, less than about 60%, less than about 50%, less than about 40%, less than about 30%, less than about 20%, or less than about 10% of the level present in an indicator cell from an individual not having cancer.

[0213] In general, primers provide for amplification of a palladin target nucleic acid to produce a palladin target nucleic acid amplification product (also referred to as an "amplicon"). Primers will in some embodiments be used in conjunction with a nucleic acid probe. 5' primers generally bind to a region to provide for amplification of the target nucleic, and in many embodiments bind to a 5' portion of the target sequence, as shown in FIGS. 10 and 14-16. 3' primers generally bind to a sequence that is complementary to a 3' portion of the nucleic acid generated by extension from the 5' primer, as exemplified in FIGS. 10 and 14-16.

[0214] Target nucleotide sequences to which 5' and 3' primers hybridize will be separated from one another by from about 10 nucleotides to about 650 nucleotides, e.g., from about 10 nucleotides to about 20 nucleotides, from about 20 nucleotides to about 30 nucleotides, from about 30 nucleotides to about 40 nucleotides, from about 40 nucleotides to about 50 nucleotides, from about 50 nucleotides to about 60 nucleotides, from about 60 nucleotides to about 70 nucleotides, from about 70 nucleotides to about 100 nucleotides, from about 100 nucleotides to about 150 nucleotides, from about 150 nucleotides to about 200 nucleotides, from about 200 nucleotides to about 250 nucleotides, from about 250 nucleotides to about 300 nucleotides, from about 300 nucleotides to about 400 nucleotides, from about 400 nucleotides to about 500 nucleotides, or from about 500 nucleotides to about 650 nucleotides.

[0215] The amplification product that is generated will have a length of from about 30 nucleotides to about 670 nucleotides, e.g., from about 30 nucleotides to about 50 nucleotides, from about 50 nucleotides to about 100 nucleotides, from about 100 nucleotides to about 150 nucleotides, from about 150 nucleotides to about 200 nucleotides, from about 200 nucleotides to about 250 nucleotides, from about 250 nucleotides to about 300 nucleotides, from about 300 nucleotides to about 400 nucleotides, from about 400 nucleotides to about 500 nucleotides, or from about 500 nucleotides to about 670 nucleotides.

[0216] In certain embodiments, primers are designed so as to have a sequence complementary to one or more variant nucleotides within a target region sequence and/or to have a 3' end adjacent a variant nucleotide of a sequence of a target region. In some embodiments involving amplification-based detection, probes are designed so as to have a sequence complementary to a sequence flanked by the sequence(s) complementary to one or more primers used for amplification.

[0217] The amplification product will in many embodiments have a length in a range of from about 30 nucleotides

(or base pairs, bp) to about 1000 nucleotides (or base pairs), e.g., from about 30 bp to about 50 bp, from about 50 bp to about 60 bp, from about 60 bp to about 70 bp, from about 70 bp to about 80 bp, from about 80 bp to about 90 bp, from about 90 bp to about 100 bp, from about 100 bp to about 150 bp, from about 150 bp to about 200 bp, from about 200 bp to about 250 bp, from about 250 bp to about 300 bp, from about 300 bp to about 350 bp, from about 350 bp to about 400 bp, from about 400 bp to about 500 bp, from about 500 bp to about 600 bp, from about 700 bp, from about 700 bp to about 800 bp, from about 800 bp to about 900 bp, or from about 900 bp to about 1000 bp (e.g., about 1 kb).

[0218] In some embodiments, the primer sequences are in the range of between 10-75 nucleotides in length, such as 10 to 70 nucleotides, 12 to 65 nucleotides, 15 to 60 nucleotides, 20 to 55 nucleotides, 25 to 50 nucleotides, 30 to 45 nucleotides, and the like. In some embodiments, primers are in the range of between 18 to 40, 19 to 35, 20 to 30, 21 to 29, 22 to 28, 23 to 27, 24-25 nucleotides long, and any length between the stated ranges. Primers of about 20 to 22 nucleotides in length are of particular interest in some embodiments.

[0219] In some embodiments, the first and/or the second primer comprises a detectable label. Suitable labels include fluorochromes, e.g. fluorescein isothiocyanate (FITC), rhodamine, Texas Red, phycoerythrin, allophycocyanin, 6-carboxyfluorescein (6-FAM), 2',7'-dimethoxy-4',5'-dichloro-6-carboxyfluorescein (JOE), 6-carboxy-X-rhodamine (ROX), 6-carboxy-2',4',7',4,7-hexachlorofluorescein (HEX), 5-carboxyfluorescein (5-FAM) or N,N,N',N'-tetramethyl-6-carboxyrhodamine (TAMRA); radioactive labels, e.g. ³²P, ³⁵S, ³H; etc. The label may be a two stage system, where the amplified DNA is conjugated to biotin, haptens, etc. having a high affinity binding partner, e.g. avidin, specific antibodies, etc., where the binding partner is conjugated to a detectable label. The label may be conjugated to one or both of the primers. Alternatively, the pool of nucleotides used in the amplification is labeled, so as to incorporate the label into the amplification product.

[0220] In one non-limiting exemplary embodiment, where the target nucleic acid comprises Target Region 1d, where Target Region 1d is:

[0221] 5'-ggacagggcgt cccactgctc gtcgctgcc accgcttcg gccacggcca gacgtccgcg gcttctca gcgctgct gacctcgcg ccgccgcccg cgccgctcaa gcacctggg-3' (SEQ ID NO:63; corresponding to nucleotides 621-780 of SEQ ID NO:1), or the complement thereof, a suitable primer pair is:

[0222] Primer pair A:

[0223] primer 1: 5'-ggacagggcgtccactgctc-3' (SEQ ID NO:64); and

[0224] primer 2: 5'-ccccagggcgtgacggccc-3' (SEQ ID NO:65).

[0225] Target region 1d, and a suitable primer pair for amplifying a target nucleic acid comprising Target region 1d, are depicted in FIG. 10.

[0226] Primer pair A is suitable for use in methods of detecting a nucleotide substitution that is associated with pancreatic cancer. Primer pair A will in some embodiments be used to amplify a palladin nucleic acid comprising Target Region 1d. In some embodiments, the amplification product generated using Primer pair A and a nucleic acid target comprising Target Region 1d will then be analyzed using a nucleic acid probe that hybridizes to and detects a nucleotide substitution that is associated with pancreatic cancer, as described above. For example, any of probes 1-4 are suitable for use.

[0227] Primer pair A is also suitable for use in a method of detecting a palladin nucleic acid that is overexpressed in an indicator cell that is a cancerous or precancerous cell (e.g., a cancerous or precancerous pancreatic cell) or that is under-expressed in an indicator cell that is a white blood cell.

[0228] In other non-limiting exemplary embodiments, where the target nucleic acid comprises Target Region 1, where Target Region 1 is:

[0229] 5'-ggacctctgaagctccagcaactcca-gaaccaaatccgactggagcaggaggccgctcgg cagcctccgccagc-cccgcgagcgcgcccctcgcceccctcccgccgcccgccttcc cggagctcgcgctgacgccgc-ccgctccccggagcccatgagcgcgctggctccccgctc cgccccgcatg-cagtcctccgctctcaactacgcgcceccaagcagttcatcgccgcg cagaacctcggcccgcgctcgggccacg-gcacgcggcctccagcccagctcgtccagcctcc cgtegccatgctc-cccagcggcggagcagttcggccgcccccgctccgcccctcgcgcagcc ctccgctcgcgctgacggaggc-cccgtgggctcctcctcgcctcgccecccgccccgccacc ccggtct-tcagccccagcgcgtccttcccggctc-cgacgtgttccactgcccggcccaccac cgccgctcccagcccggacaggcgtc-ccactgctcgtcgcctgccaccgcttcggccacgg ccagacgTccgcgacct tctcagcgcctcgtcctcgcagccgcgcccggcgccgtaac gccctgggctgcccagggtgtcaccgcc-3' (SEQ ID NO:6; corresponding to nucleotides 132-803 of SEQ ID NO:1, but including the C→T substitution, shown in bold and underlined), or the complement thereof, suitable primer pairs include:

[0230] Primer Pair B:

[0231] 5'-GGACCCTCTGAAGCTCCAGC-3' (SEQ ID NO:66); and

[0232] 5'-GCGGGGTGACACCCTTGGG-3' (SEQ ID NO:67), where

[0233] Primer 1 of pair B corresponds to nucleotides 132-151 of SEQ ID NO:1; and Primer 2 of pair B corresponds to the complement of nucleotides 784-803 of SEQ ID NO:1.

[0234] Primer Pair C:

[0235] 5'-GGACCCTCTGAAGCTCCAGC-3' (SEQ ID NO:68); and

[0236] Primer 2: 5'-CCGTGGCCGAAGCGGGTGCAGG-3' (SEQ ID NO:69), where

[0237] Primer 1 of pair C corresponds to nucleotides 132-151 of SEQ ID NO:1; and Primer 2 of pair C corresponds to the complement of nucleotides 685-707 of SEQ ID NO:1.

[0238] Primer Pair D:

[0239] 5'-CCTCGCCGTCGCCCCGCC-3' (SEQ ID NO:70); and

[0240] Primer 2: 5'-CCGTGGCCGAAGCGGGTGCAGG-3' (SEQ ID NO:71), where

[0241] Primer 1 of pair D corresponds to nucleotides 551 to 575 of SEQ ID NO:1; and Primer 2 of pair D corresponds to the complement of nucleotides 685-707 of SEQ ID NO:1.

[0242] Primer Pair E:

[0243] 5'-CCTCGCCGTCGCCCCGCC-3' (SEQ ID NO:72); and

[0244] 5'-CCGTGGCCGAAGCGGGTGGCAGG-3' (SEQ ID NO:73),

[0245] where Primer 1 of pair E corresponds to nucleotides 551 to 575 of SEQ ID NO:1; and

[0246] Primer 2 of pair E corresponds to the complement of nucleotides 685-707 of SEQ ID NO:1. Primer pair B is suitable for use in methods of detecting a nucleotide substitution that is associated with cancer (e.g., pancreatic cancer). Primer

pair B will in some embodiments be used to amplify a palladin nucleic acid comprising Target Region 1. In some embodiments, the amplification product generated using Primer pair B and a nucleic acid target comprising Target Region 1 will then be analyzed using a nucleic acid probe that hybridizes to and detects a nucleotide substitution that is associated with pancreatic cancer, as described above. For example, any of probes 1-4 are suitable for use.

[0247] Primer pair B is also suitable for use in a method of detecting a palladin nucleic acid that is overexpressed in a cancerous or precancerous pancreatic cell. Primer pairs C, D, and E are suitable for use in a method of detecting a palladin nucleic acid that is overexpressed in a cancerous or precancerous pancreatic cell.

[0248] Primer Pair F:

[0249] 5'-AGGTGTCACTTCTCTTTTCCCCC-3' (SEQ ID NO:74); and

[0250] Primer 2: 5'-GGGGAGGGAAGTGGAGGACCGCGG-3' (SEQ ID NO:75),

[0251] where Primer 1 of pair F corresponds to nucleotides 139671 to 139695 of the genomic sequence depicted in FIG. 16; and where Primer 2 of pair F corresponds to the complement of nucleotides 140401 to 140425 of the sequence depicted in FIG. 16.

[0252] Primer pair F is suitable for use in methods of detecting a nucleotide substitution that is associated with cancer. Primer pair F will in some embodiments be used to amplify a genomic palladin nucleic acid comprising exon 2 (e.g., as shown in bold text in FIG. 16). In some embodiments, the amplification product generated using Primer pair F and a nucleic acid target comprising palladin exon 2 will then be analyzed using a nucleic acid probe that hybridizes to and detects a nucleotide substitution that is associated with cancer, as described above.

[0253] Primer Pair G:

[0254] 5'-cagtgggta ccaacccag-3' (SEQ ID NO:76); and

[0255] 5'-gtgcacccgttctcagca-3' (SEQ ID NO:77).

[0256] Primer 1 of pair G corresponds to nucleotides 1641-1660 of SEQ ID NO:1; and Primer 2 of pair G corresponds to the complement of nucleotides 1721-1740 of SEQ ID NO:1.

[0257] Primer pair G is suitable for use in methods of detecting a nucleotide substitution that is associated with cancer. Primer pair G will in some embodiments be used to amplify a palladin nucleic acid comprising Target Region 3. In some embodiments, the amplification product generated using Primer pair G and a nucleic acid target comprising Target Region 3 will then be analyzed using a nucleic acid probe that hybridizes to and detects a nucleotide substitution that is associated with cancer, as described above.

[0258] Primer pair G is also suitable for use in a method of detecting a palladin nucleic acid that is overexpressed in an indicator cells that is a cancerous or precancerous cell (e.g., a cancerous or precancerous pancreatic cell), or that is overexpressed in an indicator cell that is a white blood cell

[0259] Primer Pair H

[0260] 5' TCTTGACTACTGAAGGAGGAATTTATGC (SEQ ID NO:78); and

[0261] 5' TTTTCCGTATTGGTTAGTAATGTA-GAATTAG (SEQ ID NO:79),

[0262] where Primer 1 of primer pair H corresponds to nucleotides 183428-18356 of BAC RP11-635L1 (GenBank Accession No. AC080188; as depicted in FIG. 14), and Primer 2 of primer pair H corresponds to the complement of nucleotides 183798-183828 of BAC RP11-635L1 (GenBank

Accession No. AC080188; as depicted in FIG. 14). Primers 1 and 2 of primer pair H are underlined in the sequence set forth in FIG. 14.

[0263] Primer pair H is suitable for use in methods of detecting a nucleotide substitution that is associated with cancer. Primer pair H will in some embodiments be used to amplify a palladin nucleic acid comprising palladin exon 9 (as shown in bold text in FIG. 14). In some embodiments, the amplification product generated using Primer pair H and a nucleic acid target comprising palladin exon 9 will then be analyzed using a nucleic acid probe that hybridizes to and detects a nucleotide substitution that is associated with cancer, as described above.

[0264] Primer Pair I:

[0265] 5'-gggaaac gaacggagtc catccc-3' (SEQ ID NO:80); and

[0266] 5'-agcaggacagctggtgtgg-3' (SEQ ID NO:81).

[0267] Primer 1 of pair I corresponds to nucleotides 2034-2056 of the sequence depicted in FIGS. 13A and 13B (NM_016081); and Primer 2 of pair I corresponds to the complement of nucleotides 2111-2130 of the sequence depicted in FIGS. 13A and 13B (NM_016081).

[0268] Primer pair I is suitable for use in methods of detecting a nucleotide substitution that is associated with cancer. Primer pair G will in some embodiments be used to amplify a palladin nucleic acid comprising Target Region 4. In some embodiments, the amplification product generated using Primer pair G and a nucleic acid target comprising Target Region 4 will then be analyzed using a nucleic acid probe that hybridizes to and detects a nucleotide substitution that is associated with cancer, as described above. For example, any of probes 8-10 are suitable for use.

[0269] Primer pair I is also suitable for use in a method of detecting a palladin nucleic acid that is overexpressed in a cancerous or precancerous cell.

[0270] Primer Pair J

[0271] 5' CACAACACAGGGATTCTCAGAAGA (SEQ ID NO:82)

[0272] 5' AGCTGGTGTGGAAGAGATT (SEQ ID NO:83),

[0273] where primer 1 of pair J corresponds to nucleotides 26927-26950 of the genomic DNA sequence set forth in BAC RP11-592K15 (GenBank Accession No. AC084353; as depicted in FIG. 15), and Primer 2 of primer pair J corresponds to the complement of nucleotides 27294-27313 of BAC RP11-592K15 (GenBank Accession No. AC084353; as depicted in FIG. 15). Primers 1 and 2 of primer pair J are underlined in the sequence set forth in FIG. 15.

[0274] Primer pair J is suitable for use in methods of detecting a nucleotide substitution that is associated with cancer. Primer pair J will in some embodiments be used to amplify a palladin nucleic acid comprising at least a portion of palladin exon 10. In some embodiments, the amplification product generated using Primer pair J and a nucleic acid target comprising at least a portion of palladin exon 10 will then be analyzed using a nucleic acid probe that hybridizes to and detects a nucleotide substitution that is associated with cancer, as described above.

[0275] Primer Pair K

[0276] 5'-CCCCGCGCCCGGTCGCGGA-3' (SEQ ID NO:115), corresponding to nucleotides 685-704 of the nucleotide sequence depicted in FIG. 19 and set forth in SEQ ID NO:110; and

[0277] 5'-CGCGTCCGGAGCGAGCGGGA-3' (SEQ ID NO:116), corresponding to the complement of nucleotides 717-736 of the nucleotide sequence depicted in FIG. 19 and set forth in SEQ ID NO:110.

[0278] Primer pair K flanks the 12-base pair insertion depicted in FIG. 19; and is suitable for amplifying a target palladin nucleic acid that includes a nucleotide sequence insertion in a palladin promoter.

Internal Control Nucleic Acids

[0279] In certain embodiments, an internal control (IC) or an internal standard is added to serve as a control to show that any negative result is not due to failure of the assay. The use of the IC permits the control of the separation process, the amplification process, and the detection system, and permits the monitoring of assay performance and quantification for the sample(s). The IC can be included at any suitable point, for example, in the lysis buffer. In one embodiment, the IC comprises phage nucleic acid. Where a solid support is used in the assay, the solid support may additionally include probes specific to the internal standard (IC probe), thereby facilitating capture when using the IC probe. The IC probe can optionally be coupled with a detectable label that is different from the detectable label for the target sequence. In embodiments where the detectable label is a fluorophore, the IC can be quantified spectrophotometrically and by limit of detection studies.

[0280] In another embodiment, an IC, as described herein, is combined with RNA isolated from the sample according to standard techniques known to those of skill in the art, and described herein. The RNA is then reverse-transcribed using a reverse transcriptase to provide copy DNA. The cDNA sequences can be optionally amplified (e.g., by PCR) using labeled primers. The amplification products are separated, typically by electrophoresis, and the amount of radioactivity (proportional to the amount of amplified product) is determined. The amount of mRNA in the sample can then be calculated where desired by comparison with the signal produced by the known standards.

Synthesis of Primers and Probes

[0281] Primers and probes described above are designed based on the sequences disclosed herein and are readily synthesized by standard techniques, e.g., solid phase synthesis via phosphoramidite chemistry, as disclosed in U.S. Pat. Nos. 4,458,066 and 4,415,732, incorporated herein by reference; Beaucage et al. (1992) *Tetrahedron* 48:2223-2311; and Applied Biosystems User Bulletin No. 13 (1 Apr. 1987). Other chemical synthesis methods include, for example, the phosphotriester method described by Narang et al., *Meth. Enzymol.* (1979) 68:90 and the phosphodiester method disclosed by Brown et al., *Meth. Enzymol.* (1979) 68:109. Poly (A) or poly(C), or other non-complementary nucleotide extensions may be incorporated into probes using these same methods. Hexaethylene oxide extensions may be coupled to probes by methods known in the art. Cload et al. (1991) *J. Am. Chem. Soc.* 113:6324-6326; U.S. Pat. No. 4,914,210 to Levenson et al.; Durand et al. (1990) *Nucleic Acids Res.* 18:6353-6359; and Horn et al. (1986) *Tet. Lett.* 27:4705-4708.

Compositions

[0282] The present invention further provides compositions comprising a nucleic acid suitable for use in a subject

method (e.g., a nucleic acid probe as described above; a primer pair as described above). Compositions comprising a subject nucleic acid will in some embodiments include one or more of: a salt, e.g., NaCl, MgCl, KCl, MgSO₄, etc.; a buffering agent, e.g., a Tris buffer, N-(2-Hydroxyethyl)piperazine-N'-(2-ethanesulfonic acid) (HEPES), 2-(N-Morpholino) ethanesulfonic acid (MES), 2-(N-Morpholino) ethanesulfonic acid sodium salt (MES), 3-(N-Morpholino) propanesulfonic acid (MOPS), N-tris[Hydroxymethyl] methyl-3-aminopropanesulfonic acid (TAPS), etc.; a solubilizing agent; a detergent, e.g., a non-ionic detergent such as Tween-20, etc.; a nuclease inhibitor; and the like. In some embodiments, a subject nucleic acid is lyophilized.

Antibodies

[0283] Detection of a precancerous or cancerous cell (e.g., a cancerous or precancerous pancreatic cell, breast cancer cell, head and neck squamous cell carcinoma, etc.) can also be accomplished by detecting a palladin protein that is encoded by a palladin nucleic acid comprising a mutation associated with cancer, or by detecting a palladin protein that is present in an indicator cell at an abnormal level. For example, where the palladin gene mutation associated with cancer (e.g., pancreatic cancer, breast cancer, head and neck squamous cell carcinoma, etc.) results in a mutated palladin polypeptide, e.g., a palladin polypeptide that differs in amino acid sequence from "wild-type" palladin, the mutated palladin polypeptide can be detected. Detection of a mutated palladin polypeptide in a biological sample indicates the presence in the biological sample, or in a cell or tissue from which the biological sample was derived, of a precancerous or cancerous cell (e.g., a precancerous or cancerous pancreatic cell, precancerous or cancerous breast cell, head and neck squamous cell carcinoma, etc.).

[0284] As another example, the presence or absence of cancer in a subject is assessed by detecting an abnormal level of palladin polypeptide in an indicator cell (e.g., an elevated level in a pancreatic cell, breast cell, upper aerodigestive tract cell, axillary lymph node cell, etc., relative to a normal (non-cancerous) palladin polypeptide level; or decreased in a white blood cell relative to a normal white blood cell palladin polypeptide level in white blood cells of a non-cancerous subject). In this example, detection of a level of palladin polypeptide provides for detection of a cancerous or precancerous cell in the subject from whom the indicator cell was obtained.

[0285] The present invention provides antibody reagents, including isolated antibodies, which bind specifically to a palladin polypeptide, e.g., a palladin polypeptide comprising an epitope that includes an amino acid substitution encoded by a palladin gene mutation associated with pancreatic cancer. A palladin-specific antibody is useful for detecting a palladin polypeptide (including a mutant palladin polypeptide), and therefore finds use in certain embodiments in diagnostic methods, e.g., methods involving detection of pancreatic cancer. In certain embodiments, a palladin-specific antibody is isolated, e.g., is in an environment other than its naturally-occurring environment. Suitable anti-palladin antibodies include antibodies of any isotype; single-chain Fv; Fab; Fab₂; Fv; F(ab')₂; artificial antibodies; and the like. In some embodiments, the antibody is other than the 1E6 antibody described in Rachlin and Otey (2006) *J. Cell Sci.* 119: 995-1004.

[0286] Suitable antibodies are obtained by immunizing a host animal with peptides comprising all or a portion of a palladin protein. Suitable host animals include mouse, rat, sheep, goat, hamster, rabbit, etc. The host animal will generally be from a different species than the immunogen where the immunogen is from a naturally occurring source, e.g., a human sample, where representative host animals include, but are not limited to, e.g., rabbits, goats, mice, etc.

[0287] The immunogen may comprise the complete protein, or fragments and derivatives thereof. Generally, immunogens comprise all or a part of the protein, where these residues contain any post-translation modifications found on the native target protein. Immunogens are produced in a variety of ways known in the art, e.g., expression of cloned genes using conventional recombinant methods, preparation of fragments of a subject deacylase protein using well-known methods, etc.

[0288] For preparation of polyclonal antibodies, the first step is immunization of the host animal with the target protein, where the target protein can be in substantially pure form, comprising less than about 1% contaminant. The immunogen may comprise the complete palladin protein, fragments or derivatives thereof. To increase the immune response of the host animal, the palladin protein may be combined with an adjuvant, where suitable adjuvants include alum, dextran, sulfate, large polymeric anions, and oil-and-water emulsions, e.g. Freund's adjuvant, Freund's complete adjuvant, and the like. The palladin protein may also be conjugated to synthetic carrier proteins or synthetic antigens. A variety of hosts may be immunized to produce the polyclonal antibodies. Such hosts include rabbits, guinea pigs, rodents, e.g. mice, rats, sheep, goats, and the like. The palladin protein is administered to the host, e.g., intradermally or intramuscularly, with an initial dosage followed by one or more, usually at least two, additional booster dosages. Following immunization, the blood from the host will be collected, followed by separation of the serum from the blood cells. The Ig present in the resultant antiserum may be further fractionated using known methods, such as ammonium salt fractionation, DEAE chromatography, and the like.

[0289] Monoclonal antibodies are produced by conventional techniques. Generally, the spleen and/or lymph nodes of an immunized host animal provide a source of plasma cells. The plasma cells are immortalized by fusion with myeloma cells to produce hybridoma cells. Culture supernatant from individual hybridomas is screened using standard techniques to identify those producing antibodies with the desired specificity. Suitable animals for production of monoclonal antibodies include mouse, rat, hamster, guinea pig, rabbit, etc. The antibody may be purified from the hybridoma cell supernatants or ascites fluid by conventional techniques, e.g. affinity chromatography using protein bound to an insoluble support, protein A sepharose, etc.

[0290] The antibody may be produced as a single chain, instead of the normal multimeric structure. Single chain antibodies are described in Jost et al. (1994) J.B.C. 269:26267-73, and others. DNA sequences encoding the variable region of the heavy chain and the variable region of the light chain are ligated to a spacer encoding at least about 4 amino acids of small neutral amino acids, including glycine and/or serine. The protein encoded by this fusion allows assembly of a functional variable region that retains the specificity and affinity of the original antibody.

[0291] Anti-palladin antibodies also include "artificial" antibodies, e.g., antibodies and antibody fragments produced and selected *in vitro*. In some embodiments, such antibodies are displayed on the surface of a bacteriophage or other viral particle. In many embodiments, such artificial antibodies are present as fusion proteins with a viral or bacteriophage structural protein, including, but not limited to, M13 gene III protein. Methods of producing such artificial antibodies are well known in the art. See, e.g., U.S. Pat. Nos. 5,516,637; 5,223,409; 5,658,727; 5,667,988; 5,498,538; 5,403,484; 5,571,698; and 5,625,033.

[0292] Antibody fragments, such as Fv, F(ab')₂ and Fab may be prepared by cleavage of the intact protein, e.g. by protease or chemical cleavage. Alternatively, a truncated gene is designed. For example, a chimeric gene encoding a portion of the F(ab')₂ fragment would include DNA sequences encoding the CH1 domain and hinge region of the H chain, followed by a translational stop codon to yield the truncated molecule.

[0293] Expression vectors include plasmids, retroviruses, YACs, EBV derived episomes, and the like. A convenient vector is one that encodes a functionally complete human CH or CL immunoglobulin sequence, with appropriate restriction sites engineered so that any VH or VL sequence can be easily inserted and expressed. In such vectors, splicing usually occurs between the splice donor site in the inserted J region and the splice acceptor site preceding the human C region, and also at the splice regions that occur within the human CH exons. Polyadenylation and transcription termination occur at native chromosomal sites downstream of the coding regions. The resulting chimeric antibody may be joined to any strong promoter, including retroviral LTRs, e.g. SV-40 early promoter, (Okayama et al. (1983) Mol. Cell. Bio. 3:280), Rous sarcoma virus LTR (Gorman et al. (1982) P.N.A.S. 79:6777), and moloney murine leukemia virus LTR (Grosschedl et al. (1985) Cell 41:885); native Ig promoters, etc.

[0294] An anti-palladin antibody will in some embodiments be detectably labeled, e.g., with a radioisotope, an enzyme which generates a detectable product, a fluorescent protein, a chromogenic protein, and the like. An anti-palladin antibody may be further conjugated to other moieties, such as members of specific binding pairs, e.g., biotin (member of biotin-avidin specific binding pair), and the like. An anti-palladin antibody may also be bound to a solid support, including, but not limited to, polystyrene plates or beads, magnetic beads, test strips, membranes, and the like.

[0295] In some embodiments, the anti-palladin antibody is detectably labeled, either directly or indirectly. Direct labels include radioisotopes (e.g., ¹²⁵I, ³⁵S, and the like); enzymes whose products are detectable (e.g., luciferase, β -galactosidase, horse radish peroxidase, alkaline phosphatase, and the like); fluorescent labels (e.g., fluorescein isothiocyanate, rhodamine, phycoerythrin, and the like); fluorescence emitting metals, e.g., ¹⁵²Eu, or others of the lanthanide series, attached to the antibody through metal chelating groups such as EDTA; chemiluminescent compounds, e.g., luminol, isoluminol, acridinium salts, and the like; bioluminescent compounds, e.g., luciferin; fluorescent proteins; and the like. Indirect labels include second antibodies specific for palladin-specific antibodies, wherein the second antibody is labeled as described above; and members of specific binding pairs, e.g., biotin-avidin, and the like.

[0296] In some embodiments, an anti-palladin antibody comprises, covalently linked to the antibody, a protein that

provides for a detectable signal. Suitable proteins include, but are not limited to, fluorescent proteins and enzymes (e.g., β -galactosidase, luciferase, horse radish peroxidase, alkaline phosphatase, etc.). Suitable fluorescent proteins include, but are not limited to, a green fluorescent protein (GFP), including, but not limited to, a GFP derived from *Aequoria Victoria* or a derivative thereof, a number of which are commercially available; a GFP from a species such as *Renilla reniformis*, *Renilla mulleri*, or *Ptilosarcus guernei*, as described in, e.g., WO 99/49019 and Peelle et al. (2001) *J. Protein Chem.* 20:507-519; any of a variety of fluorescent and colored proteins from Anthozoan species, as described in, e.g., Matz et al. (1999) *Nature Biotechnol.* 17:969-973, U.S. Patent Publication No. 2002/0197676, or U.S. Patent Publication No. 2005/0032085; and the like.

[0297] In some embodiments, an anti-palladin antibody is specific for a particular isoform of palladin. For example, in some embodiments, an anti-palladin antibody is specific for the 90 kD isoform of palladin. In other embodiments, an anti-palladin antibody is specific for the 140 kD isoform of palladin. In some embodiments, an anti-palladin antibody is specific for wild-type palladin, e.g., where "wild-type" palladin has an amino acid sequence that does not include mutations associated with pancreatic cancer.

[0298] In a particular embodiment, an anti-palladin antibody suitable for use in a diagnostic or detection method binds specifically to an epitope comprising amino acid 239 of the amino acid sequence depicted in FIG. 11. A diagnostic method could involve use of a first anti-palladin antibody specific for an epitope comprising amino acid 239 of the amino acid sequence depicted in FIG. 11, where amino acid 239 is Pro; and a second anti-palladin antibody specific for an epitope comprising amino acid 239 of the amino acid sequence depicted in FIG. 11, where amino acid 239 is Ser.

[0299] In another embodiment of interest, an anti-palladin antibody suitable for use in a diagnostic or detection method binds specifically to an epitope comprising amino acid 557 of the amino acid sequence depicted in FIG. 11. A diagnostic method could involve use of a first anti-palladin antibody specific for an epitope comprising amino acid 557 of the amino acid sequence depicted in FIG. 11, where amino acid 557 is Trp; and a second anti-palladin antibody specific for an epitope comprising amino acid 557 of the amino acid sequence depicted in FIG. 11, where amino acid 557 is Cys.

[0300] In another embodiment of interest, an anti-palladin antibody suitable for use in a diagnostic or detection method binds specifically to an epitope comprising amino acid 617 of the amino acid sequence depicted in FIG. 12. A diagnostic method could involve use of a first anti-palladin antibody specific for an epitope comprising amino acid 617 of the amino acid sequence depicted in FIG. 12, where amino acid 617 is Arg; and a second anti-palladin antibody specific for an epitope comprising amino acid 557 of the amino acid sequence depicted in FIG. 12, where amino acid 617 is Cys.

Diagnostic and Detection Methods

[0301] The present invention provides various diagnostic and detection methods relating to cancer (e.g., pancreatic cancer, breast cancer, head and neck squamous cell cancer, etc.). The invention provides methods for detection of the presence or absence of a pre-cancerous cell in a subject; methods for detection of the presence or absence of a cancerous cell in a subject; methods for identification of individuals at risk of developing cancer; methods for staging of cancer

cells; and methods for assessing response to a treatment for cancer. The presence or absence of a cancerous or pre-cancerous cell can be detected either directly (e.g., by assessing a cell suspected of being cancerous for a palladin mutation or aberrant palladin expression level) or indirectly, by assessing a palladin expression level in a white blood cell.

[0302] For example, the invention provides methods for detection of a pre-cancerous pancreatic cell; methods for detection of a cancerous pancreatic cell; methods for identification of individuals at risk of developing pancreatic cancer; methods for staging of pancreatic cancer cells; and methods for assessing response to a treatment for pancreatic cancer. Nucleic acid probes and/or primers as discussed above can be used in a subject diagnostic or detection method. Similarly, anti-palladin antibodies are discussed above, and can be used in a subject diagnostic or detection method. These methods are described in detail below.

Detection and/or Diagnostic Methods Involving Assessing a Palladin mRNA Expression Level

[0303] The present invention provides methods of diagnosing a cancerous or precancerous cancer (e.g., pancreatic cancer, breast cancer, head and neck squamous cell cancer, etc.) condition by detecting abnormal levels of palladin mRNA in a biological sample. The present invention provides methods of detecting a cancerous or pre-cancerous cell in a biological sample.

[0304] A subject method of diagnosing cancer, or detecting a cancerous or pre-cancerous cell in a biological sample, generally involves detecting the level of palladin mRNA in a cell in a biological sample, where a level of palladin mRNA in the cell that indicates that the palladin mRNA is overexpressed in the cell indicates that the cell is cancerous or pre-cancerous. In some embodiments, detection of a cancerous or precancerous cell indicates that an individual has cancer; and thus provides a diagnosis of cancer. In other embodiments, a subject method of diagnosing cancer involves detecting a level of a palladin mRNA in an indicator cell obtained from an individual, where a level of palladin mRNA in the indicator cell that is abnormally low indicates the presence in the individual of cancer.

[0305] For example, a subject method of diagnosing pancreatic cancer, or detecting a cancerous or pre-cancerous pancreatic cell in a biological sample, generally involves detecting the level of palladin mRNA in a pancreatic cell in a biological sample, where a level of palladin mRNA in the cell that indicates that the palladin mRNA is overexpressed in the cell indicates that the cell is cancerous or pre-cancerous. In some embodiments, detection of a cancerous or precancerous pancreatic cell indicates that an individual has pancreatic cancer; and thus provides a diagnosis of pancreatic cancer. In other embodiments, a subject method of diagnosing pancreatic cancer involves detecting a level of a palladin mRNA in an indicator cell obtained from an individual, where a level of palladin mRNA in the indicator cell that is abnormally low indicates the presence in the individual of pancreatic cancer.

[0306] In some embodiments, mRNA is isolated from pancreatic tissue or other biological sample obtained from an individual; the mRNA is reverse transcribed to generate a sample comprising a cDNA copy of a palladin mRNA; and a subject method involves detecting the level of palladin cDNA in the sample. In other embodiments, mRNA is isolated from pancreatic tissue obtained from an individual; the mRNA is reverse transcribed to generate a sample comprising a cDNA copy of a palladin mRNA; the palladin cDNA is amplified

using a nucleic acid primer pair, generating a palladin amplification product; and a subject method involves detecting the level of palladin amplification product in the sample. In other embodiments, a probe that binds to a palladin nucleic acid that is overexpressed in pancreatic cancer is used, where a level of probe binding is correlated with a level of palladin expression.

[0307] Where a subject detection method detects the presence of a cancerous or precancerous pancreatic cell in an individual, in some embodiments the individual will undergo one or more confirmatory tests for pancreatic cancer. Where a subject detection method detects the presence of a cancerous or precancerous pancreatic cell in an individual, in some embodiments the individual will be treated for pancreatic cancer.

[0308] Treatments for pancreatic cancer include: surgery (e.g., pancreaticoduodenectomy; Whipple procedure; total pancreatectomy; partial pancreatectomy; distal pancreatectomy; etc.); radiation; chemotherapy; and combinations of two or more of the foregoing. Chemotherapy treatments include, but are not limited to, gemcitabine (Gemzar®) monotherapy; gemcitabine+erlotinib (Tarceva®; an epidermal growth factor receptor tyrosine kinase inhibitor); 5-fluorouracil; gemcitabine+erbitux; gemcitabine+bevacizumab; gemcitabine+oxaliplatin; gemcitabine+cisplatin; etc. Gemcitabine is 2'-deoxy-2',2'-difluorocytidine; see, e.g., U.S. Pat. Nos. 5,464,826 and 6,555,518.

[0309] Where a subject detection method detects the presence of a cancerous or precancerous breast cell in an individual, in some embodiments the individual will undergo one or more confirmatory tests for breast cancer. Similarly, where a subject detection method detects the presence of a cancerous or precancerous cell in the upper aerodigestive tract in an individual, in some embodiments the individual will undergo one or more confirmatory tests for head and neck cancer.

[0310] Where a subject detection method detects the presence of a cancerous or precancerous breast cell in an individual, in some embodiments the individual will be treated for breast cancer. Similarly, where a subject detection method detects the presence of a cancerous or precancerous cell in the upper aerodigestive tract in an individual, in some embodiments the individual will be treated for head and neck squamous cell carcinoma.

[0311] Detection methods can also be performed to monitor progression of a cancer. For example, a biological sample is taken from the individual and tested at a frequency of once per week, twice weekly, once per month, bimonthly, once every three months, once every four months, once every 6 months, or once a year, depending on various factors. The biological sample is tested for the level of palladin mRNA. The rate of increase in the level of palladin mRNA is an indication of the rate of progression of the cancer.

[0312] As an example, detection methods can also be performed to monitor progression of pancreatic cancer. For example, a biological sample is taken from the individual and tested at a frequency of once per week, twice weekly, once per month, bimonthly, once every three months, once every four months, once every 6 months, or once a year, depending on various factors. The biological sample is tested for the level of palladin mRNA. The rate of increase in the level of palladin mRNA is an indication of the rate of progression of the disease (e.g., pancreatic cancer).

[0313] Detection methods can also be performed to assess response to therapy, e.g., therapy for pancreatic cancer,

therapy for breast cancer, therapy for head and neck squamous cell carcinoma, etc. For example, where the pancreatic cancer is associated with an elevated level of palladin expression, the levels of palladin expression during the course of and/or following therapy can be assessed, thus providing an indicator of response to therapy. Measurements of palladin mRNA levels can be used to determine whether a patient is responding to treatment. In some embodiments, palladin mRNA levels are measured before and after a treatment, e.g. surgery or a drug treatment, to determine if the treatment is efficacious. In other embodiments, palladin mRNA levels are determined during the course of the treatment, to determine whether the treatment slows the progression of the disease, and to what extent the treatment slows the progression of the disease. For example, a reduction of at least about 10%, at least about 20%, at least about 25%, at least about 30%, or at least about 40% or more, in the rate of increase in the level of palladin mRNA in response to a given treatment indicates that the treatment is efficacious in treating pancreatic cancer.

Detection and/or Diagnostic Methods Involving Assessing a Palladin Polypeptide Level and/or Detecting Mutant Palladin Polypeptide

[0314] Methods of diagnosing cancer, and methods of detecting a cancerous or precancerous cell, as described above, can also be carried out by detecting palladin polypeptide levels. Thus, the present invention provides methods of diagnosing cancer, and methods of detecting a cancerous or precancerous cell, where the methods generally involve detecting a level of palladin polypeptide in a biological sample. An abnormal level of palladin polypeptide in an indicator cell of a biological sample from an individual indicates the presence or absence of a cancerous or precancerous cell in the individual. For example, an elevated level of palladin polypeptide above normal levels in a cell suspected of being cancerous or pre-cancerous can indicate that the individual has cancer. In another example, a decreased level of palladin polypeptide in a white blood cell of an individual relative to a normal level of palladin in a white blood cell of an unaffected subject can indicate that the individual has cancer. The source of palladin protein can be the same as the source of palladin nucleic acid. Sources of palladin nucleic acids are discussed above.

[0315] For example, methods of diagnosing pancreatic cancer, and methods of detecting a cancerous or precancerous pancreatic cell, as described above, can also be carried out by detecting palladin polypeptide levels. Thus, the present invention provides methods of diagnosing pancreatic cancer, and methods of detecting a cancerous or precancerous pancreatic cell, where the methods generally involve detecting a level of palladin polypeptide in a biological sample. An abnormal level of palladin polypeptide in a biological sample from an individual indicates the presence in the biological sample, and/or in the individual, of a cancerous or precancerous pancreatic cell. For example, an elevated level of palladin polypeptide above normal levels may indicate that the individual has pancreatic cancer.

[0316] Methods of diagnosing cancer, and methods of detecting a cancerous or precancerous cell, as described above, can also be carried out by detecting mutant palladin polypeptide. Thus, the present invention provides methods of diagnosing cancer, and methods of detecting a cancerous or precancerous cell, where the methods generally involve detecting mutant palladin polypeptide in a biological sample suspected of containing a cancerous or pre-cancerous cell,

where the mutant palladin polypeptide is encoded by a palladin nucleic acid comprising one or more mutations associated with cancer. Non-limiting examples of mutant palladin polypeptides that provide for diagnosis of cancer and/or detection of a cancerous or precancerous cell include: a) a palladin polypeptide comprising a Pro→Ser mutation at a position corresponding to amino acid 239 of the sequence depicted in FIG. 11 (GenBank BAA76836.1); b) a palladin polypeptide comprising a Trp→Cys mutation at a position corresponding to amino acid 557 of the sequence depicted in FIG. 11 (GenBank BAA76836.1); and c) an Arg→Cys mutation at a position corresponding to amino acid 617 of the sequence depicted in FIG. 12 (GenBank NP_057165.3).

[0317] For example, methods of diagnosing pancreatic cancer, and methods of detecting a cancerous or precancerous pancreatic cell, as described above, can also be carried out by detecting mutant palladin polypeptide. Thus, the present invention provides methods of diagnosing pancreatic cancer, and methods of detecting a cancerous or precancerous pancreatic cell, where the methods generally involve detecting mutant palladin polypeptide in a biological sample, where the mutant palladin polypeptide is encoded by a palladin nucleic acid comprising one or more mutations associated with pancreatic cancer. Non-limiting examples of mutant palladin polypeptides that provide for diagnosis of pancreatic cancer and/or detection of a cancerous or precancerous pancreatic cell include: a) a palladin polypeptide comprising a Pro→Ser mutation at a position corresponding to amino acid 239 of the sequence depicted in FIG. 11 (GenBank BAA76836.1); b) a palladin polypeptide comprising a Trp→Cys mutation at a position corresponding to amino acid 557 of the sequence depicted in FIG. 11 (GenBank BAA76836.1); and c) an Arg→Cys mutation at a position corresponding to amino acid 617 of the sequence depicted in FIG. 12 (GenBank NP_057165.3).

[0318] The palladin polypeptide that is being detected may be isolated, but need not be. In some embodiments, palladin polypeptide is isolated from a biological sample obtained from an individual. In other embodiments, palladin polypeptide is detected in crude cell lysates, in whole cells, or in partially purified preparations (e.g., total protein isolated from a biological sample). The biological sample is contacted with an anti-palladin antibody (e.g., an antibody that specifically binds palladin); and binding of the anti-palladin antibody to a protein in the sample is detected. Binding of the anti-palladin antibody to protein in the biological sample indicates the presence in the sample of palladin protein.

[0319] In some embodiments, the anti-palladin antibody is specific for the 90 kD isoform of palladin. In some of these embodiments, the anti-palladin antibody detects a level of palladin 90 kD isoform in the biological sample. In other embodiments, the anti-palladin antibody is specific for the 140 kD isoform of palladin. In some of these embodiments, the anti-palladin antibody detects a level of palladin 145 kD isoform in the biological sample.

[0320] Where a subject method involves detecting palladin polypeptide levels, palladin polypeptide levels can be measured using any standard method, including, but not limited to, immunoprecipitation, Western blotting, enzyme-linked immunosorbent assays (ELISA), radioimmunoassay, etc., where antibody specific for palladin polypeptide is employed. The anti-palladin antibody can include a detectable label, as described above. In some embodiments, a subject method that provides for detection of a palladin polypep-

tide level detects a level of wild-type palladin. Alternatively, the polypeptide can be detected using proteomic methods, such as mass spectroscopy. In other embodiments, a subject method that provides for detection of a palladin polypeptide level detects a level of palladin polypeptide containing one or more mutations associated with pancreatic cancer. In other embodiments, a subject method that provides for detection of a palladin polypeptide level detects both a level of wild-type and a level of mutated palladin.

[0321] Where a subject method involves detecting mutated palladin (e.g., palladin polypeptide comprising one or more mutations associated with cancer), the method will in some embodiments involve contacting a biological sample with one or more of: a) an anti-palladin antibody that binds specifically to an epitope comprising amino acid 239 of the amino acid sequence depicted in FIG. 11; b) an anti-palladin antibody that binds specifically to an epitope comprising amino acid 557 of the amino acid sequence depicted in FIG. 11; and c) an anti-palladin antibody that binds specifically to an epitope comprising amino acid 617 of the amino acid sequence depicted in FIG. 12.

[0322] In some embodiments, a subject method will involve detecting both wild-type and mutant palladin polypeptide. For example, a detection or diagnostic method will in some embodiments involve contacting a biological sample with: a) a first anti-palladin antibody specific for an epitope comprising amino acid 239 of the amino acid sequence depicted in FIG. 11, where amino acid 239 is Pro; and a second anti-palladin antibody specific for an epitope comprising amino acid 239 of the amino acid sequence depicted in FIG. 11, where amino acid 239 is Ser; b) a first anti-palladin antibody specific for an epitope comprising amino acid 557 of the amino acid sequence depicted in FIG. 11, where amino acid 557 is Trp; and a second anti-palladin antibody specific for an epitope comprising amino acid 557 of the amino acid sequence depicted in FIG. 11, where amino acid 557 is Cys; or c) a first anti-palladin antibody specific for an epitope comprising amino acid 617 of the amino acid sequence depicted in FIG. 12, where amino acid 617 is Arg; and a second anti-palladin antibody specific for an epitope comprising amino acid 557 of the amino acid sequence depicted in FIG. 12, where amino acid 617 is Cys.

[0323] Where a subject detection method detects the presence of a cancerous or precancerous cell in an individual, in some embodiments the individual will undergo one or more confirmatory tests for the cancer. Where a subject detection method detects the presence of a cancerous or precancerous cell in an individual, in some embodiments the individual will be treated for the cancer.

[0324] For example, where a subject detection method detects the presence of a cancerous or precancerous pancreatic cell in an individual, in some embodiments the individual will undergo one or more confirmatory tests for pancreatic cancer. Where a subject detection method detects the presence of a cancerous or precancerous pancreatic cell in an individual, in some embodiments the individual will be treated for pancreatic cancer.

[0325] Detection methods can also be performed to monitor progression of pancreatic cancer. For example, a biological sample is taken from the individual and tested at a frequency of once per week, twice weekly, once per month, bimonthly, once every three months, once every four months, once every 6 months, or once a year, depending on various factors. The biological sample is tested for the level of palla-

din polypeptide, e.g., the level of mutant palladin polypeptide. The rate of increase in the level of palladin polypeptide is an indication of the rate of progression of the disease (e.g., pancreatic cancer).

[0326] Detection methods can also be performed to assess response to therapy. For example, where the pancreatic cancer is associated with an elevated level of palladin expression, the levels of palladin expression during the course of and/or following therapy can be assessed, thus providing an indicator of response to therapy. Measurements of palladin polypeptide levels can be used to determine whether a patient is responding to treatment. In some embodiments, palladin polypeptide levels are measured before and after a treatment, e.g. surgery or a drug treatment, to determine if the treatment is efficacious. In other embodiments, palladin polypeptide levels are determined during the course of the treatment, to determine whether the treatment slows the progression of the disease, and to what extent the treatment slows the progression of the disease. For example, a reduction of at least about 10%, at least about 20%, at least about 25%, at least about 30%, or at least about 40% or more, in the rate of increase in the level of palladin polypeptide in response to a given treatment indicates that the treatment is efficacious in treating the cancer (e.g., pancreatic cancer, breast cancer, head and neck squamous cell carcinoma, etc.).

Diagnostic Methods Involving Detection of Palladin Nucleic Acid Having a Cancer-Associated Nucleotide Substitution

[0327] The present invention provides methods of determining the likelihood that an individual has or will develop cancer, the methods generally involving detecting a nucleotide substitution associated with cancer in a palladin target nucleic acid of the individual, e.g., a C→T nucleotide substitution at position 715 in a palladin nucleic acid, a G→T nucleotide substitution at position 1617 in a palladin nucleic acid, or a C→T nucleotide substitution at position 2060 in a palladin nucleic acid, as described above. The mutations in palladin, however, are not limited to the ones listed herein.

[0328] For example, the present invention provides methods of determining the likelihood that an individual has or will develop pancreatic cancer, the methods generally involving detecting a nucleotide substitution associated with pancreatic cancer in a palladin target nucleic acid of a pancreatic cell of the individual, e.g., a C→T nucleotide substitution at position 715 in a palladin nucleic acid, a G→T nucleotide substitution at position 1617 in a palladin nucleic acid, or a C→T nucleotide substitution at position 2060 in a palladin nucleic acid, as described above.

[0329] Detecting a nucleotide substitution associated with cancer in a palladin target nucleic acid can involve contacting the palladin target nucleic acid with a subject palladin probe under stringent hybridization conditions, where the subject palladin probe hybridizes only with a target palladin nucleic acid that comprises a nucleotide substitution associated with cancer; and detecting hybridization between the probe and the target. Hybridization between the probe and the target indicates that the individual has a palladin nucleic acid that includes a nucleotide substitution associated with cancer; and that the individual either has cancer, or has an increased risk of developing cancer compared to an individual who does not have a palladin nucleic acid that includes the nucleotide substitution.

[0330] Detection of a nucleotide substitution associated with cancer in a palladin target nucleic acid can involve use of

a primer pair (which can be used with or without a probe), where the primer pair provides for amplification of a target nucleic acid such that the amplification product is produced at levels above background when a target palladin nucleic acid having the nucleotide substitution is present.

[0331] Detection of a nucleotide substitution associated with cancer in a palladin target nucleic acid can also be accomplished using a combination of primers and probes. For example, primer and probes described herein can be adapted for use in a detection system that provides for amplification and detection of the nucleotide substitution, e.g., in a real-time PCR assay. Exemplary systems, discussed in more detail herein, include Taqman™ systems, Amplifluor™ hairpin primer-based systems, Scorpion systems (e.g., bi-functional molecules containing a PCR primer element covalently linked to a probe element), Sunrise™ primer-based systems, Molecular Beacons, and a Light Upon Extension or LUX™-based systems.

[0332] Substitution can also be detected using any test that detects a change in the palladin sequence, including, but not limited to, restriction enzyme digestion or lack of digestion at the location of a substitution in palladin DNA or cDNA. For example, a nucleotide substitution can provide a sequence that is cut by a restriction endonuclease such as HindIII or PvuII, while the non-mutated sequence is not cut by the restriction endonuclease.

[0333] In some embodiments, the target palladin nucleic acid is palladin genomic DNA. In other embodiments, the target palladin nucleic acid is palladin mRNA, a cDNA copy of palladin mRNA, or an amplification product of palladin genomic DNA, palladin mRNA, or palladin cDNA.

[0334] Where the diagnostic method indicates that the individual has an increased risk of developing cancer, the risk that the individual will develop cancer is at least about 25%, at least about 50%, at least about 2-fold, at least about 2.5-fold, at least about 5-fold, at least about 10-fold, or at least about 20-fold, or more, higher than the risk that an individual without the nucleotide substitution associated with cancer.

[0335] For example, where the diagnostic method indicates that the individual has an increased risk of developing pancreatic cancer, the risk that the individual will develop pancreatic cancer is at least about 25%, at least about 50%, at least about 2-fold, at least about 2.5-fold, at least about 5-fold, at least about 10-fold, or at least about 20-fold, or more, higher than the risk that an individual without the nucleotide substitution associated with pancreatic cancer.

[0336] Where the diagnostic method indicates that the individual has an increased risk of developing cancer, in some embodiments the individual will be tested for the presence of cancer. Where the diagnostic method indicates that the individual has an increased risk of developing cancer, in some embodiments the individual will be monitored on a regular basis (e.g., monthly, bimonthly, twice a year, once a year, etc.) for cancer.

[0337] For example, where the diagnostic method indicates that the individual has an increased risk of developing pancreatic cancer, in some embodiments the individual will be tested for the presence of pancreatic cancer. Where the diagnostic method indicates that the individual has an increased risk of developing pancreatic cancer, in some embodiments the individual will be monitored on a regular basis (e.g., monthly, bimonthly, twice a year, once a year, etc.) for pancreatic cancer.

Diagnostic Methods Involving Detection of an Altered Palladin Gene Promoter

[0338] The present invention provides methods of detecting a cancerous or precancerous cell (e.g., a cancerous or

precancerous pancreatic cell), the methods generally involving detecting an alteration in a palladin gene promoter. In some embodiments, the alteration being detected is in the 90 kDa isoform palladin gene promoter.

[0339] Alterations include, but are not limited to, insertions of nucleotide sequences, duplications of nucleotide sequences; deletions of nucleotide sequences; inversions of nucleotide sequences; and alterations in the methylation status of a promoter.

[0340] For example, the present invention provides methods of determining the likelihood that an individual has or will develop pancreatic cancer, the methods generally involving detecting a nucleotide sequence insertion associated with pancreatic cancer in a palladin target nucleic acid of a pancreatic cell of the individual, where the nucleotide sequence insertion is in a palladin gene promoter, as described above. In some embodiments, insertion of a 12-bp sequence in the 90 kDa isoform palladin promoter provides for detection of a cancerous or precancerous cell.

Methylation Status

[0341] The methylation status of a palladin gene promoter provides for detection of a cancerous or precancerous cell. For example, the methylation status of the 90 kDa isoform palladin promoter provides for detection of a cancerous or precancerous cell. Hypomethylation of the 90 kDa isoform palladin promoter indicates a cancerous or precancerous cell.

[0342] Any of a variety of methods can be used to detect DNA methylation status, where suitable methods include, but are not limited to, methylation-specific polymerase chain reaction (MSP; see, e.g., Herman et al. (1996) *Proc. Natl. Acad. Sci. USA* 93:9821-9826); MethylLight (Eads et al. (2000) *Nucl. Acids Res.* 28:E32; and U.S. Pat. No. 6,331,393); HeavyMethyl (Cottrell et al. (2004) *Nucl. Acids Res.* 32:e10); MethylQuant (Thomassin et al. (2004) *Nucl. Acids Res.* 32:e168; and the like.

[0343] A number of methods involve treatment of a target DNA with a bisulfite reagent, which converts unmethylated cytosines to uracils, leaving only methylated cytosines unchanged (see, e.g., WO 05/038051). Following bisulfite treatment, individual cytosine positions can be detected by a primer extension reaction (Gonzalzo and Jones (1997) *Nucleic Acids Res.* 25:2529-31; and WO 95/00669) or by enzymatic digestion (Xiong and Laird (1997) *Nucleic Acids Res.* 25: 2535-4). Alternatively, following bisulfite treatment, a methylation-specific polymerase chain reaction (PCR) can be carried out, using primers that bind either to methylated or unmethylated DNA only and that therefore selectively amplify only DNA with a defined methylation. MethylLight is a variation of MSP, and involves use of a methylation-specific real-time detection probe (MethylLight), which makes the assay both homogenous and quantitative. HeavyMethyl is also a variation on MSP. In the HeavyMethyl method, the priming is methylation specific, but non-extendable oligonucleotide blockers provide this specificity instead of the primers themselves. The blockers bind to bisulfite-treated DNA in a methylation-specific manner, and their binding sites overlap the primer binding sites. When the blocker is bound, the primer cannot bind and therefore the amplicon is not generated. When the blocker is not bound, the primer-binding site is accessible and the amplicon is generated. HeavyMethyl in combination with real-time detection with methylation-specific fluorogenic probes provides sensitive and specific detection of DNA methylation.

[0344] In some embodiments, a methylation-sensitive restriction endonuclease (e.g., a restriction endonuclease that recognizes and cleaves a nucleic acid having a particular nucleotide sequence only when the sequence is unmethylated), and/or a methylation-insensitive restriction endonuclease (e.g., a restriction endonuclease that recognizes and cleaves a nucleic acid having a particular nucleotide sequence, regardless of the methylation status of the nucleotide sequence); are used. The term "methylation-sensitive enzyme" refers to a restriction enzymes that does not cleave DNA (or cleaves DNA poorly) if one or more nucleotides in its recognition site are methylated. Suitable methylation-sensitive and methylation-insensitive restriction endonucleases that are suitable for use include, but are not limited to, MboI, DpnII, HpaII, BsmBI, Sau3A, and ClaI.

[0345] Various methods can be used to determine the methylation status of a target DNA. For example, indirect methods for DNA methylation pattern determinations at specific loci that have been developed rely on techniques that alter the genomic DNA in a methylation-dependent manner before an amplification event. Two exemplary methods that can be used to achieve this methylation-dependent DNA alteration include: 1) digestion by a restriction enzyme that is affected in its activity by 5-methylcytosine in a CpG sequence context. The cleavage, or lack of it, can subsequently be revealed by Southern blotting or by PCR; and 2) the treatment of genomic DNA with sodium bisulfite. Sodium bisulfite treatment converts all unmethylated cytosines in the DNA to uracil by deamination, but leaves the methylated cytosine residues intact. Subsequent PCR amplification replaces the uracil residues with thymines and the 5-methylcytosine residues with cytosines. The resulting sequence difference can be detected using standard DNA sequence detection techniques, e.g., PCR.

[0346] An exemplary method involves use of a bisulfite treatment-based method followed by a PCR reaction to analyze a target nucleic acid. There are two principally different ways in which the sequence difference generated by the sodium bisulfite treatment can be revealed. The first is to design PCR primers that uniquely anneal with either methylated or unmethylated converted DNA. This technique is referred to as "methylation specific PCR" or "MSP". See, e.g., U.S. Pat. No. 5,786,146. The method used by all other bisulfite-based techniques (such as bisulfite genomic sequencing, COBRA and Ms-SNuPE) is to amplify the bisulfite-converted DNA using primers that anneal at locations that lack CpG dinucleotides in the original genomic sequence. In this way, the PCR primers can amplify the sequence in between the two primers, regardless of the DNA methylation status of that sequence in the original genomic DNA. This results in a pool of different PCR products, all with the same length and differing in their sequence only at the sites of potential DNA methylation at CpGs located in between the two primers. The difference between these methods of processing the bisulfite-converted sequence is that in MSP, the methylation information is derived from the occurrence or lack of occurrence of a PCR product, whereas in the other techniques a mix of products is always generated and the mixture is subsequently analyzed to yield quantitative information on the relative occurrence of the different methylation states. A method such as described in U.S. Pat. No. 7,186,512 is also suitable for use.

[0347] In some embodiments, the methods involve contacting a genomic sample of DNA with a modifying agent that

modifies unmethylated cytosine (e.g., sodium bisulfite), to produce a converted nucleic acid; (b) amplifying the converted nucleic acid by means of oligonucleotide primers in the presence of one or a plurality of specific oligonucleotide probes, where the one or the plurality of the oligonucleotide primers or the specific probe(s) is/are capable of distinguishing between unmethylated and methylated nucleic acid (e.g., a CpG-specific probe capable of distinguishing between unmethylated and methylated nucleic acid); and (c) detecting, in real-time during the amplification, the methylated nucleic acid based on amplification-mediated probe displacement. See, e.g., U.S. Pat. No. 7,112,404. Amplification and detection can occur simultaneously as measured by fluorescence-based real-time quantitative PCR ("RT-PCR") using specific, dual label TaqMan® oligonucleotide probes. The displaceable probes can be specifically designed to distinguish between methylated and unmethylated CpG sites present in the original, unmodified nucleic acid sample. Sodium-bisulfite readily reacts with the 5,6-double bond of cytosine, but not with methylated cytosine, to produce a sulfonated cytosine intermediate that undergoes deamination under alkaline conditions to produce uracil. Because Taq polymerase recognizes uracil as thymine and 5-methylcytidine (m5C) as cytidine, the sequential combination of sodium bisulfite treatment and PCR amplification results in the ultimate conversion of unmethylated cytosine residues to thymine (C→U→T) and methylated cytosine residues ("mC") to cytosine (mC→mC→C). Thus, sodium-bisulfite treatment of genomic DNA creates methylation-dependent sequence differences by converting unmethylated cytosines to uracil, and upon PCR the resultant product contains cytosine only at positions where methylated cytosine occurs in the unmodified nucleic acid.

[0348] In some embodiments, the specific primers are designed to be substantially complementary to each strand of the genomic locus of interest. Typically, one primer is complementary to the negative, (-) strand of the locus (the "lower" strand of a horizontally situated double-stranded DNA molecule) and the other is complementary to the positive (+) strand ("upper" strand). In some embodiments, the primers are designed to overlap potential sites of DNA methylation (CpG nucleotides) and specifically distinguish modified unmethylated from methylated DNA. This sequence discrimination can be based upon the differential annealing temperatures of perfectly matched, versus mismatched oligonucleotides. In some embodiments, primers are typically designed to overlap from one to several CpG sequences. In other embodiments, e.g., in a quantitative embodiment, the primers do not overlap any CpG sequences.

Staging Methods Involving Analysis of Palladin Expression Levels

[0349] The present invention provides methods of staging cancer, the methods generally involving determining a level of palladin mRNA in a cell; and, based on the level of palladin mRNA, assigning the cell to a cancerous stage.

[0350] For example, in some embodiments, where the level of palladin mRNA in a cell is from about 2-fold higher to about 5-fold higher than the level of palladin mRNA in a normal, non-cancerous cell of the same cell type, the cell is designated stage I.

[0351] In some embodiments, where the level of palladin mRNA in a cell is from about 5-fold higher to about 10-fold

higher than the level of palladin mRNA in a normal, non-cancerous cell of the same cell type, the cell is designated stage 11.

[0352] In some embodiments, where the level of palladin mRNA in a cell is from about 10-fold higher to about 50-fold higher than the level of palladin mRNA in a normal, non-cancerous cell of the same cell type, the cell is designated stage III.

[0353] These stage designations can be applied to any cancer associated with abnormal expression of palladin, e.g., pancreatic cancer, breast cancer, head and neck cancer, etc.

[0354] For example, the present invention provides methods of staging pancreatic cancer, the methods generally involving determining a level of palladin mRNA in a pancreatic cell; and, based on the level of palladin mRNA, assigning the pancreatic cell to a cancerous stage.

[0355] For example, in some embodiments, where the level of palladin mRNA in a pancreatic cell is from about 2-fold higher to about 5-fold higher than the level of palladin mRNA in a normal, non-cancerous pancreatic cell of the same cell type, the pancreatic cell is designated stage I.

[0356] In some embodiments, where the level of palladin mRNA in a pancreatic cell is from about 5-fold higher to about 10-fold higher than the level of palladin mRNA in a normal, non-cancerous pancreatic cell of the same cell type, the pancreatic cell is designated stage 11.

[0357] In some embodiments, where the level of palladin mRNA in a pancreatic cell is from about 10-fold higher to about 50-fold higher than the level of palladin mRNA in a normal, non-cancerous pancreatic cell of the same cell type, the pancreatic cell is designated stage III.

[0358] Stage I disease could be considered local cancer (within the organ that contains the cancer). Stage 11 disease could be considered regional (cancer extending beyond the organ of origin into the adjacent structures and lymph nodes), Stage III disease could be considered widespread or metastatic disease (spread of the cancer to areas remote from the organ of origin). Thus, the present methods provide for staging cancer metastasis.

Hybridization and Amplification Methods for Use in a Detection or Diagnostic Method

[0359] The diagnostic, detection, staging methods described above will in many embodiments involve nucleic acid hybridization with a nucleic acid probe, nucleic acid amplification with a nucleic acid primer pair, or both. Nucleic acid hybridization and nucleic acid amplification methods are known to those skilled in the art. Exemplary nucleic acid hybridization and nucleic acid amplification methods are discussed in detail below. The following provides detail of exemplary nucleic acid-based methods for detection, and examples of how such can be adapted for use in the methods of the invention.

Detection and Diagnostic Methods Using Hybridization with a Nucleic Acid Probe

[0360] As discussed above, where a subject detection or diagnostic method involves detecting a nucleotide sequence alteration (e.g., a C→T nucleotide substitution at position 715 in a palladin nucleic acid), the method will in some embodiments comprise hybridizing a target palladin nucleic acid with a nucleic acid probe. Also as discussed above, where a subject method involves detecting a level of a palladin nucleic acid in a cell, the method will in some embodiments comprise hybridizing a target palladin nucleic acid with a nucleic acid

probe. Nucleic acid hybridization methods that are suitable for use in a subject method are described below.

[0361] In some embodiments, a subject method for detecting a nucleotide sequence alteration associated with cancer involves contacting, under stringent hybridization conditions, a subject nucleic acid probe with a target palladin nucleic acid, where the subject probe hybridizes only to a target palladin nucleic acid that comprises a nucleotide sequence alteration associated with cancer (e.g., a nucleotide substitution as described above; a nucleotide insertion as described above); and detecting hybridization between the probe and the target. Hybridization of the palladin nucleic probe to the target palladin nucleic acid indicates that the target palladin nucleic acid comprises a nucleotide sequence alteration that is associated with cancer.

[0362] For example, in some embodiments, a subject method for detecting a nucleotide sequence alteration associated with pancreatic cancer involves contacting, under stringent hybridization conditions, a subject nucleic acid probe with a target palladin nucleic acid, where the subject probe hybridizes only to a target palladin nucleic acid that comprises a nucleotide sequence alteration associated with pancreatic cancer (e.g., a nucleotide sequence alteration as described above); and detecting hybridization between the probe and the target. Hybridization of the palladin nucleic probe to the target palladin nucleic acid indicates that the target palladin nucleic acid comprises a nucleotide sequence alteration that is associated with pancreatic cancer.

[0363] In some embodiments, a subject method for detecting a cancerous or pre-cancerous cell involves contacting, under stringent hybridization conditions, a subject nucleic acid probe with a target palladin nucleic acid in a sample; and detecting the level of palladin mRNA in the sample. Where the detected level of palladin mRNA indicates that palladin mRNA is overexpressed in the cell, the cell is considered cancerous or precancerous. In some of these embodiments, the palladin target nucleic acid is first amplified using a subject primer pair.

[0364] For example, in some embodiments, a subject method for detecting a cancerous or pre-cancerous pancreatic cell involves contacting, under stringent hybridization conditions, a subject nucleic acid probe with a target palladin nucleic acid in a sample; and detecting the level of palladin mRNA in the sample. Where the detected level of palladin mRNA indicates that palladin mRNA is overexpressed in the cell, the cell is considered cancerous or precancerous. In some of these embodiments, the palladin target nucleic acid is first amplified using a subject primer pair.

[0365] A number of methods are available for analyzing nucleic acids for the presence and/or level of a specific nucleic acid in a cell. The mRNA may be assayed directly or reverse transcribed into cDNA for analysis. The nucleic acid may be amplified by conventional techniques, such as the polymerase chain reaction (PCR), to provide sufficient amounts for analysis. The use of the polymerase chain reaction is described in Saiki, et al. (1985), *Science* 239:487, and a review of techniques may be found in Sambrook, et al. *Molecular Cloning: A Laboratory Manual*, CSH Press 1989, pp. 14.2-14.33.

[0366] In some embodiments, the method involves contacting the sample under stringent hybridization conditions with a subject palladin nucleic acid probe and detecting binding, if any, of the probe to a nucleic acid in the sample. A variety of nucleic acid hybridization methods are well known to those

skilled in the art, and any known method can be used. In many embodiments, the palladin nucleic acid probe will be detectably labeled.

Detection and Diagnostic Methods Using Amplification with Nucleic Acid Primer Pairs

[0367] As discussed above, where a subject detection or diagnostic method involves detecting a nucleotide sequence alteration, which is associated with cancer, in a palladin nucleic acid, the method will in some embodiments include amplification of a target palladin nucleic acid, forming a palladin amplification product; and hybridizing the palladin amplification product with a nucleic acid probe that detects the nucleotide sequence alteration. Also as discussed above, where a subject method involves detecting a level of a palladin nucleic acid in a cell, the method will in some embodiments include amplification of a target palladin nucleic acid, forming a palladin amplification product (and may further include a step of hybridizing the palladin amplification product with a nucleic acid probe).

[0368] In some embodiments, the method involves contacting a sample (e.g., under stringent hybridization conditions) with a subject nucleic acid primer pair, where the primer pair, under conditions that permit primer-initiated nucleic acid amplification, amplifies any target palladin nucleic acid present in the sample, generating an amplification product (where amplification product is generated when target palladin nucleic acid present in the sample).

[0369] Conditions that permit primer-initiated nucleic acid amplification and catalytic nucleic acid activity are well known to those skilled in the art, and include the presence of a DNA polymerase; deoxynucleotide triphosphates; and magnesium ions. Suitable reaction conditions are well known to those skilled in the art of nucleic acid amplification. Exemplary, non-limiting reaction conditions are described in the Examples. The DNA polymerase is generally one that has high affinity for binding at the 3'-end of an oligonucleotide hybridized to a nucleic acid strand. The DNA polymerase is generally one that has little or no 5'→3' exonuclease activity so as to minimize degradation of primer, termination or primer extension polynucleotides. The DNA polymerase is generally one that has little to no proofreading activity. In many embodiments, the DNA polymerase is thermostable, e.g., is catalytically active at temperatures in excess of about 75° C. DNA polymerases that are suitable for use in a subject method include, but are not limited to, DNA polymerases discussed in U.S. Pat. Nos. 5,648,211 and 5,744,312, which include *exo⁻ Vent* (New England Biolabs), *exo⁻ Deep Vent* (New England Biolabs), *Bst* (BioRad), *exo-Pfu* (Stratagene), *Bca* (Panvera), sequencing grade *Taq* (Promega); thermostable DNA polymerases from *Thermoanaerobacter thermo-hydrosulfuricus*; and the like. In some embodiments, the reaction mixture includes an RNase H.

[0370] Magnesium ions are typically present in the reaction mix in a concentration of from about 1 mM to about 100 mM, e.g., from about 1 mM to about 3 mM, from about 3 mM to about 5 mM, from about 5 mM to about 10 mM, from about 10 mM to about 25 mM, from about 25 mM to about 50 mM, from about 50 mM to about 75 mM, or from about 75 mM to about 100 mM.

[0371] Usually the reaction mixture will comprise four different types of dNTPs corresponding to the four naturally occurring bases are present, i.e. dATP, dTTP, dCTP and dGTP. In the subject methods, each dNTP will typically be present at a final concentration in the reaction, ranging from

about 10 μM to 5000 μM , e.g., from about 10 μM to about 50 μM , from about 50 μM to about 100 μM , from about 100 μM to about 200 μM , from about 200 μM to about 500 μM , from about 500 μM to about 1000 μM , from about 1000 μM to about 2000 μM , from about 2000 μM to about 3000 μM , from about 3000 μM to about 4000 μM , or from about 4000 μM to about 5000 μM . In some embodiments, each dNTP will be present at a final concentration in the reaction of from about 20 μM to 1000 μM , from about 100 μM to about 200 μM , or from about 50 μM to about 200 μM .

[0372] The amplification reaction mixture typically includes an aqueous buffer medium that includes a source of monovalent ions, a source of divalent cations and a buffering agent. Any convenient source of monovalent ions, such as KCl, K-acetate, NH_4 -acetate, K-glutamate, NH_4Cl , ammonium sulfate, and the like may be employed. The divalent cation may be magnesium, manganese, zinc and the like, where the cation will typically be magnesium. Any convenient source of magnesium cation may be employed, including MgCl_2 , Mg-acetate, and the like. Representative buffering agents or salts that may be present in the buffer include Tris, Tricine, HEPES, MOPS and the like, where the amount of buffering agent will typically range from about 5 to 150 mM, usually from about 10 to 100 mM, and more usually from about 20 to 50 mM, where in certain embodiments the buffering agent will be present in an amount sufficient to provide a pH ranging from about 6.0 to 9.5, e.g., pH 7.3 at 72° C. Other agents which may be present in the buffer medium include chelating agents, such as EDTA, EGTA and the like.

[0373] Each primer nucleic acid is present in the reaction mixture at a concentration of from about 50 nM to about 900 nM, e.g., the 3' primer and the 5' primer nucleic acid are each independently present at a concentration of from about 50 nM to about 75 nM, from about 75 nM to about 100 nM, from about 100 nM to about 150 nM, from about 150 nM to about 200 nM, from about 200 nM to about 250 nM, from about 250 nM to about 300 nM, from about 300 nM to about 400 nM, from about 400 nM to about 500 nM, from about 500 nM to about 600 nM, from about 600 nM to about 700 nM, from about 700 nM to about 800 nM, or from about 800 nM to about 900 nM.

[0374] A detectable label may be included in an amplification reaction. Suitable labels include fluorochromes, e.g. fluorescein isothiocyanate (FITC), rhodamine, Texas Red, phycoerythrin, allophycocyanin, 6-carboxyfluorescein (6-FAM), 2',7'-dimethoxy-4',5'-dichloro-6-carboxyfluorescein (JOE), 6-carboxy-X-rhodamine (ROX), 6-carboxy-2',4',7',4'-hexachlorofluorescein (HEX), 5-carboxyfluorescein (5-FAM) or N,N,N',N'-tetramethyl-6-carboxyrhodamine (TAMRA), radioactive labels, e.g. ^{32}P , ^{35}S , ^3H ; etc. The label may be a two stage system, where the amplified DNA is conjugated to biotin, haptens, etc. having a high affinity binding partner, e.g. avidin, specific antibodies, etc., where the binding partner is conjugated to a detectable label. The label may be conjugated to one or both of the primers. Alternatively, the pool of nucleotides used in the amplification is labeled, so as to incorporate the label into the amplification product.

[0375] In one embodiment, a subject method involves amplifying nucleic acids from a sample, which amplifying step follows a reverse transcription step to provide a cDNA template for amplification. If a diagnostic nucleic acid is obtained, the presence or absence of a nucleotide substitution associated with cancer (e.g., pancreatic cancer, breast cancer,

head and neck cancer, etc.) can be indicated, where the presence of a nucleotide substitution associated with pancreatic cancer indicates an increased likelihood that the individual from whom the nucleic acid was obtained has or will develop cancer (e.g., pancreatic cancer, breast cancer, head and neck cancer, etc.). In other embodiments, the level of a palladin mRNA can be indicated, where overexpression of a palladin mRNA indicates a cancerous or precancerous cell. In general, amplification-based methods involve reverse transcription of mRNA in a sample and amplifying the resulting cDNA from the sample using a primer and at least one other primer, as described above, and assessing the amplified nucleic acids.

[0376] As is known in the art, an amplified nucleic acid may be assessed by a number of methods, including, for example, determining the presence or absence of the nucleic acid, determining the size of the nucleic acid or determining the abundance of a nucleic acid in relation to another amplified nucleic acid. In most embodiments, an amplified nucleic acid is assessed using gel electrophoresis, nucleic acid hybridization, sequencing, and/or detection of a signal from a label bound to the amplified nucleic acid. Methods of amplifying (e.g., by polymerase chain reaction) nucleic acid, methods of performing primers extension, and methods of assessing nucleic acids are generally well known in the art (e.g., see Ausubel, et al, Short Protocols in Molecular Biology, 3rd ed., Wiley & Sons, 1995 and Sambrook, et al, Molecular Cloning: A Laboratory Manual, Third Edition, (2001) Cold Spring Harbor, N.Y.) and need not be described in any great detail.

[0377] For example, primers and probes described above may be used in polymerase chain reaction (PCR)-based techniques to detect palladin target nucleic acid (e.g., to detect a nucleotide substitution associated with pancreatic cancer; to detect a level of palladin mRNA; etc.) in biological samples. PCR is a technique for amplifying a desired target nucleic acid sequence contained in a nucleic acid molecule or mixture of molecules. In PCR, a pair of primers is employed in excess to hybridize to the complementary strands of the target nucleic acid. The primers are each extended by a polymerase using the target nucleic acid as a template. The extension products become target sequences themselves after dissociation from the original target strand. New primers are then hybridized and extended by a polymerase, and the cycle is repeated to geometrically increase the number of target sequence molecules. The PCR method for amplifying target nucleic acid sequences in a sample is well known in the art and has been described in, e.g., Innis et al. (eds.) PCR Protocols (Academic Press, NY 1990); Taylor (1991) Polymerase chain reaction: basic principles and automation, in PCR: A Practical Approach, McPherson et al. (eds.) IRL Press, Oxford; Saiki et al. (1986) *Nature* 324:163; as well as in U.S. Pat. Nos. 4,683,195, 4,683,202 and 4,889,818, all incorporated herein by reference in their entireties.

[0378] In particular, PCR uses relatively short oligonucleotide primers which flank the target nucleotide sequence to be amplified, oriented such that their 3' ends face each other, each primer extending toward the other. The polynucleotide sample is extracted and denatured, e.g. by heat, and hybridized with first and second primers which are present in molar excess. Polymerization is catalyzed in the presence of the four deoxyribonucleotide triphosphates (dNTPs-dATP, dGTP, dCTP and dTTP) using a primer- and template-dependent polynucleotide polymerizing agent, such as any enzyme capable of producing primer extension products, for example, *E. coli* DNA polymerase I, Klenow fragment of DNA poly-

merase I, T4 DNA polymerase, thermostable DNA polymerases isolated from *Thermus aquaticus* (Taq), available from a variety of sources (for example, Perkin Elmer), *Thermus thermophilus* (United States Biochemicals), *Bacillus stearothermophilus* (Bio-Rad), or *Thermococcus litoralis* ("Vent" polymerase, New England Biolabs). This results in two "long products" which contain the respective primers at their 5' ends covalently linked to the newly synthesized complements of the original strands.

[0379] The reaction mixture is then returned to polymerizing conditions, e.g., by lowering the temperature, inactivating a denaturing agent, or adding more polymerase, and a second cycle is initiated. The second cycle provides the two original strands, the two long products from the first cycle, two new long products replicated from the original strands, and two "short products" replicated from the long products. The short products have the sequence of the target sequence with a primer at each end. On each additional cycle, an additional two long products are produced, and a number of short products equal to the number of long and short products remaining at the end of the previous cycle. Thus, the number of short products containing the target sequence grows exponentially with each cycle. PCR is typically carried out with a commercially available thermal cycler, e.g., Perkin Elmer.

[0380] RNAs encoding a deacylase of interest can be amplified by reverse transcribing the mRNA into cDNA, and then performing PCR (RT-PCR), as described above. Alternatively, a single enzyme may be used for both steps as described in U.S. Pat. No. 5,322,770. mRNA may also be reverse transcribed into cDNA, followed by asymmetric gap ligase chain reaction (RT-AGLCR) as described by Marshall et al. (1994) PCR Meth. App. 4:80-84.

[0381] The fluorogenic 5' nuclease assay, known as the TAQMAN™ assay (Perkin-Elmer), is a powerful and versatile PCR-based detection system for nucleic acid targets. For a detailed description of the TAQMAN™ assay, reagents and conditions for use therein, see, e.g., Holland et al., Proc. Natl. Acad. Sci., U.S.A. (1991) 88:7276-7280; U.S. Pat. Nos. 5,538,848, 5,723,591, and 5,876,930, all incorporated herein by reference in their entireties. Hence, primers and probes derived from regions of a palladin nucleic acid as described herein can be used in TAQMAN™ analyses to detect the presence of a nucleotide substitution associated with pancreatic cancer and/or to detect a level of palladin mRNA in a biological sample. Analysis is performed in conjunction with thermal cycling by monitoring the generation of fluorescence signals. The assay system dispenses with the need for gel electrophoretic analysis, and has the capability to generate quantitative data allowing the determination of, for example, the level of palladin mRNA (e.g., to detect the presence of a cancerous pancreatic cell; to detect the presence of a pre-cancerous pancreatic cell; to stage a pancreatic tumor; etc.).

[0382] The fluorogenic 5' nuclease assay is conveniently performed using, for example, AMPLITAQ GOLD™ DNA polymerase, which has endogenous 5' nuclease activity, to digest an internal oligonucleotide probe labeled with both a fluorescent reporter dye and a quencher (see, Holland et al., Proc. Natl. Acad. Sci. USA (1991) 88:7276-7280; and Lee et al., Nucl. Acids Res. (1993) 21:3761-3766). Assay results are detected by measuring changes in fluorescence that occur during the amplification cycle as the fluorescent probe is digested, uncoupling the dye and quencher labels and causing an increase in the fluorescent signal that is proportional to the amplification of target nucleic acid.

[0383] The amplification products can be detected in solution or using solid supports. In this method, the TAQMAN™ probe is designed to hybridize to a target sequence within the desired PCR product. The 5' end of the TAQMAN™ probe contains a fluorescent reporter dye. The 3' end of the probe is blocked to prevent probe extension and contains a dye that will quench the fluorescence of the 5' fluorophore. During subsequent amplification, the 5' fluorescent label is cleaved off if a polymerase with 5' exonuclease activity is present in the reaction. Excision of the 5' fluorophore results in an increase in fluorescence which can be detected.

[0384] In particular, the oligonucleotide probe is constructed such that the probe exists in at least one single-stranded conformation when unhybridized where the quencher molecule is near enough to the reporter molecule to quench the fluorescence of the reporter molecule. The oligonucleotide probe also exists in at least one conformation when hybridized to a target polynucleotide such that the quencher molecule is not positioned close enough to the reporter molecule to quench the fluorescence of the reporter molecule. By adopting these hybridized and unhybridized conformations, the reporter molecule and quencher molecule on the probe exhibit different fluorescence signal intensities when the probe is hybridized and unhybridized. As a result, it is possible to determine whether the probe is hybridized or unhybridized based on a change in the fluorescence intensity of the reporter molecule, the quencher molecule, or a combination thereof. In addition, because the probe can be designed such that the quencher molecule quenches the reporter molecule when the probe is not hybridized, the probe can be designed such that the reporter molecule exhibits limited fluorescence unless the probe is either hybridized or digested.

[0385] Accordingly, the present invention provides methods for amplifying a target palladin nucleotide sequence using a nucleic acid polymerase having 5' to 3' nuclease activity, one or more primers capable of hybridizing to the target sequence or its extension product, and an oligonucleotide probe capable of hybridizing to the target sequence 3' relative to the primer. During amplification, the polymerase digests the oligonucleotide probe when it is hybridized to the target sequence, thereby separating the reporter molecule from the quencher molecule. As the amplification is conducted, the fluorescence of the reporter molecule is monitored, with fluorescence corresponding to the occurrence of nucleic acid amplification. The reporter molecule is in some embodiments a fluorescein dye and the quencher molecule is in some embodiments a rhodamine dye.

[0386] The palladin target nucleic acids described herein may also be used as a basis for transcription-mediated amplification (TMA) assays. TMA provides a method of identifying target nucleic acids present in very small amounts in a biological sample. Such nucleic acids may be difficult or impossible to detect using direct assay methods. In particular, TMA is an isothermal, autocatalytic nucleic acid target amplification system that can provide more than a billion RNA copies of a target sequence. The assay can be done qualitatively, to accurately detect the presence or absence of the target sequence in a biological sample. The assay can also provide a quantitative measure of the amount of target sequence over a concentration range of several orders of magnitude. TMA provides a method for autocatalytically synthesizing multiple copies of a target nucleic acid sequence without repetitive manipulation of reaction conditions such as temperature, ionic strength and pH.

[0387] Generally, TMA includes the following steps: (a) isolating nucleic acid from the biological sample of interest (e.g., pancreatic tissue; pancreatic cell(s); tissue adjacent pancreatic tissue; etc.); and (b) combining into a reaction mixture (i) the isolated nucleic acid, (ii) first and second oligonucleotide primers, the first primer having a complexing sequence sufficiently complementary to the 3' terminal portion of an RNA target sequence, if present (for example the (+) strand), to complex therewith, and the second primer having a complexing sequence sufficiently complementary to the 3' terminal portion of the target sequence of its complement (for example, the (-) strand) to complex therewith, wherein the first oligonucleotide further comprises a sequence 5' to the complexing sequence which includes a promoter, (iii) a reverse transcriptase or RNA and DNA dependent DNA polymerases, (iv) an enzyme activity which selectively degrades the RNA strand of an RNA-DNA complex (such as an RNase H) and (v) an RNA polymerase which recognizes the promoter.

[0388] The components of the reaction mixture may be combined stepwise or at once. The reaction mixture is incubated under conditions whereby an oligonucleotide/target sequence is formed, including DNA priming and nucleic acid synthesizing conditions (including ribonucleotide triphosphates and deoxyribonucleotide triphosphates) for a period of time sufficient to provide multiple copies of the target sequence. The reaction advantageously takes place under conditions suitable for maintaining the stability of reaction components such as the component enzymes and without requiring modification or manipulation of reaction conditions during the course of the amplification reaction. Accordingly, the reaction may take place under conditions that are substantially isothermal and include substantially constant ionic strength and pH. The reaction conveniently does not require a denaturation step to separate the RNA-DNA complex produced by the first DNA extension reaction.

[0389] Suitable DNA polymerases include reverse transcriptases, such as avian myeloblastosis virus (AMV) reverse transcriptase (available from, e.g., Seikagaku America, Inc.) and Moloney murine leukemia virus (MMLV) reverse transcriptase (available from, e.g., Bethesda Research Laboratories).

[0390] Promoters or promoter sequences suitable for incorporation in the primers are nucleic acid sequences (either naturally occurring, produced synthetically or a product of a restriction digest) that are specifically recognized by an RNA polymerase that recognizes and binds to that sequence and initiates the process of transcription whereby RNA transcripts are produced. The sequence may optionally include nucleotide bases extending beyond the actual recognition site for the RNA polymerase which may impart added stability or susceptibility to degradation processes or increased transcription efficiency. Examples of useful promoters include those which are recognized by certain bacteriophage polymerases such as those from bacteriophage T3, T7 or SP6, or a promoter from *E. coli*. These RNA polymerases are readily available from commercial sources, such as New England Biolabs and Epicentre.

[0391] Some of the reverse transcriptases suitable for use in the methods herein have an RNase H activity, such as AMV reverse transcriptase. In some embodiments, an exogenous RNase H, such as *E. coli* RNase H, is added, even when AMV reverse transcriptase is used. RNase H is readily available from, e.g., Bethesda Research Laboratories.

[0392] The RNA transcripts produced by these methods may serve as templates to produce additional copies of the target sequence through the above-described mechanisms. The system is autocatalytic and amplification occurs autocatalytically without the need for repeatedly modifying or changing reaction conditions such as temperature, pH, ionic strength or the like.

[0393] Another method of detection involves use of target sequence-specific oligonucleotide probes, which contain a region of complementarity to the target sequence described above. The probes may be used in hybridization protection assays (HPA). In this embodiment, the probes are conveniently labeled with acridinium ester (AE), a highly chemiluminescent molecule. See, e.g., Nelson et al. (1995) "Detection of Acridinium Esters by Chemiluminescence" in *Nonisotopic Probing, Blotting and Sequencing*, Kricka L. J. (ed) Academic Press, San Diego, Calif.; Nelson et al. (1994) "Application of the Hybridization Protection Assay (HPA) to PCR" in *The Polymerase Chain Reaction*, Mullis et al. (eds.) Birkhauser, Boston, Mass.; Weeks et al., *Clin. Chem.* (1983) 29:1474-1479; Berry et al., *Clin. Chem.* (1988) 34:2087-2090. One AE molecule is directly attached to the probe using a non-nucleotide-based linker arm chemistry that allows placement of the label at any location within the probe. See, e.g., U.S. Pat. Nos. 5,585,481 and 5,185,439. Chemiluminescence is triggered by reaction with alkaline hydrogen peroxide which yields an excited N-methyl acridone that subsequently collapses to ground state with the emission of a photon. Additionally, AE causes ester hydrolysis which yields the nonchemiluminescent-methyl acridinium carboxylic acid.

[0394] When the AE molecule is covalently attached to a nucleic acid probe, hydrolysis is rapid under mildly alkaline conditions. When the AE-labeled probe is exactly complementary to the target nucleic acid, the rate of AE hydrolysis is greatly reduced. Thus, hybridized and unhybridized AE-labeled probe can be detected directly in solution, without the need for physical separation.

[0395] HPA generally consists of the following steps: (a) the AE-labeled probe is hybridized with the target nucleic acid in solution for about 15 to about 30 minutes. A mild alkaline solution is then added and AE coupled to the unhybridized probe is hydrolyzed. This reaction takes approximately 5 to 10 minutes. The remaining hybrid-associated AE is detected as a measure of the amount of target present. This step takes approximately 2 to 5 seconds. In some embodiments, the differential hydrolysis step is conducted at the same temperature as the hybridization step, typically at 50 to 70 degrees Celsius. Alternatively, a second differential hydrolysis step may be conducted at room temperature. This allows elevated pHs to be used, for example in the range of 10-11, which yields larger differences in the rate of hydrolysis between hybridized and unhybridized AE-labeled probe. HPA is described in detail in, e.g., U.S. Pat. Nos. 6,004,745; 5,948,899; and 5,283,174, the disclosures of which are incorporated by reference herein in their entireties.

[0396] TMA is described in detail in, e.g., U.S. Pat. No. 5,399,491, the disclosure of which is incorporated herein by reference in its entirety. In one example of a typical assay, an isolated nucleic acid sample, suspected of containing a deacylase-encoding nucleic acid as described herein, is mixed with a buffer concentrate containing the buffer, salts, magnesium, nucleotide triphosphates, primers, dithiothreitol, and spermidine. The reaction is optionally incubated at about 100° C. for

approximately two minutes to denature any secondary structure. After cooling to room temperature, reverse transcriptase, RNA polymerase, and RNase H are added and the mixture is incubated for two to four hours at 37° C. The reaction can then be assayed by denaturing the product, adding a probe solution, incubating 20 minutes at 60° C., adding a solution to selectively hydrolyze the unhybridized probe, incubating the reaction six minutes at 60° C., and measuring the remaining chemiluminescence in a luminometer.

[0397] Oligonucleotides will in some embodiments be used in nucleic acid sequence-based amplification (NASBA). This method is a promoter-directed, enzymatic process that induces in vitro continuous, homogeneous and isothermal amplification of a specific nucleic acid to provide RNA copies of the nucleic acid. The reagents for conducting NASBA include a first DNA primer with a 5' tail comprising a promoter, a second DNA primer, reverse transcriptase, RNase-H, T7 RNA polymerase, NTP's and dNTP's. Using NASBA, large amounts of single-stranded RNA are generated from either single-stranded RNA or DNA, or double-stranded DNA. When RNA is to be amplified, the ssRNA serves as a template for the synthesis of a first DNA strand by elongation of a first primer containing an RNA polymerase recognition site. This DNA strand in turn serves as the template for the synthesis of a second, complementary, DNA strand by elongation of a second primer, resulting in a double-stranded active RNA-polymerase promoter site, and the second DNA strand serves as a template for the synthesis of large amounts of the first template, the ssRNA, with the aid of a RNA polymerase. The NASBA technique is known in the art and described in, e.g., European Patent 329,822, International Patent Application No. WO 91/02814, and U.S. Pat. Nos. 6,063,603, 5,554,517 and 5,409,818, all of which are incorporated herein in their entireties.

[0398] The target palladin nucleic acids described herein are also useful in nucleic acid hybridization and amplification techniques that utilize branched DNA molecules. In a basic nucleic acid hybridization assay, single-stranded analyte nucleic acid is hybridized to a labeled single-stranded nucleic acid probe and resulting labeled duplexes are detected. Variations of this basic scheme have been developed to facilitate separation of the duplexes to be detected from extraneous materials and/or to amplify the signal that is detected. One method for amplifying the signal uses amplification multimers that are polynucleotides with a first segment that hybridizes specifically to the analyte nucleic acid or a strand of nucleic acid bound to the analyte and iterations of a second segment that hybridizes specifically to a labeled probe. The amplification is theoretically proportional to the number of iterations of the second segment. The multimers may be either linear or branched. Two general types of branched multimers are useful in these techniques: forked and combed. Methods for making and using branched nucleic acid molecules are known in the art and described in, e.g., U.S. Pat. No. 5,849,481, incorporated herein by reference in its entirety.

[0399] As is readily apparent, design of the assays described herein is subject to a great deal of variation, and many formats are known in the art. The above descriptions are merely provided as guidance and one of skill in the art can readily modify the described protocols, using techniques well known in the art.

Kits

[0400] The invention further provides a kit for use in a subject method. Kits provide components for detection of palladin nucleic acid and/or palladin polypeptide.

Kits for Detecting Palladin Nucleic Acid

[0401] A subject kit comprises a pair of nucleic acids (primer pairs), one or more nucleic acid probes, or both, where the primer pairs and probes are suitable for use in a subject method, as described above. The nucleic acids will in some embodiments be present in a suitable storage medium, e.g., buffered solution, typically in a suitable container. The kit includes the primers and/or probes, and may further include a buffer; reagents (e.g., for polymerase chain reaction (e.g., deoxynucleotide triphosphates (dATP, dTTP, dCTP, and dGTP), a thermostable DNA polymerase, a buffer suitable for polymerase chain reaction, a solution containing Mg²⁺ ions (e.g., MgCl₂), and other components well known to those skilled in the art for carrying out a polymerase chain reaction)). The kit may further include instructions for use of the kit, which instructions may be provided in a variety of forms, e.g., as printed information, on a compact disc, and the like. The kit may further include reagents necessary for extraction of DNA (or mRNA) from a biological sample (e.g., pancreatic tissue; pancreatic cells; ductal pancreatic epithelial cells; lymphocytes; peripheral white blood cells; etc.) from an individual. The kit may further include reagents necessary for reverse transcription of an mRNA, to make a cDNA copy of the mRNA.

[0402] The kit may further include positive and negative controls. An example of a positive control is a palladin nucleic acid that includes a region that will be amplified by primer pairs included in the kit. An example of a negative control is a nucleic acid (e.g., an albumin-encoding nucleic acid) that will not be amplified by nucleic acid primers included in the kit. The kits are useful in diagnostic applications, as described in detail above. For example, in some embodiments, a subject kit is useful to determine whether a given DNA sample (or an mRNA sample) obtained from an individual comprises a nucleotide substitution associated with pancreatic cancer. In other embodiments, a subject kit is useful to determine whether a palladin mRNA is present at higher or lower than normal levels in a pancreatic cell or other cell.

[0403] A kit will in some embodiments provide a standard for normalization of a level of a palladin polynucleotide to a standard, e.g., a level of a glucose-6-phosphate dehydrogenase polynucleotide (e.g., a G6PDH mRNA or cDNA copy of a G6PDH mRNA).

[0404] Exemplary kits include at least one primer, at least two primers (a 5' and a 3' primer), or at least two primers and a probe, as described above. Also included in the subject kits may be buffers, dNTPs, and controls, (e.g., positive and negative control nucleic acids) for performing the subject methods. Primers in the subject kits may be detectably labeled or unlabeled).

Kits for Detecting Palladin Polypeptide

[0405] A subject kit for detecting palladin polypeptide comprises one or more anti-palladin antibody reagents. For example, a subject kit will include one or more of: a) an anti-palladin antibody specific for the 90 kD isoform of palladin; b) an anti-palladin antibody specific for the 140 kD isoform of palladin; c) an anti-palladin antibody that binds

specifically to an epitope comprising amino acid 239 of the amino acid sequence depicted in FIG. 11; d) an anti-palladin antibody that binds specifically to an epitope comprising amino acid 557 of the amino acid sequence depicted in FIG. 11; and e) an anti-palladin antibody that binds specifically to an epitope comprising amino acid 617 of the amino acid sequence depicted in FIG. 12. In some embodiments, the anti-palladin antibody will comprise a detectable label. In some embodiments, the anti-palladin antibody will be bound to an insoluble support, e.g., a bead (e.g., a polystyrene bead, a magnetic bead, etc.); a plastic surface (e.g., the well of an ELISA plate); a membrane (e.g., a test strip; a polyvinylpyrrolidone membrane; a nitrocellulose membrane; etc.); and the like.

[0406] A subject kit can further include, if desired, one or more of various conventional components, such as, for example, containers with one or more buffers (e.g., wash buffers), detection reagents or antibodies. Printed instructions, either as inserts or as labels, indicating quantities of the components to be used and guidelines for their use, can also be included in the kit. In the present disclosure it should be understood that the specified materials and conditions are important in practicing the invention but that unspecified materials and conditions are not excluded so long as they do not prevent the benefits of the invention from being realized.

[0407] A kit will in some embodiments provide a standard for normalization of a level of a palladin polypeptide to a standard, e.g., a level of an actin polypeptide, a level of a GAPDH polypeptide, etc. A kit will in some embodiments further include negative controls, e.g., antibodies specific for a non-palladin polypeptide; and the like.

[0408] Kits may also include components for conducting western blots (e.g., pre-made gels, membranes, transfer systems, etc.); components for carrying out ELISAs (e.g., 96-well plates; plates containing wells in multiples of 96); components for carrying out immunoprecipitation (e.g. protein A); columns, especially spin columns, for affinity or size separation of palladin polypeptide from other (non-palladin) polypeptides or separation of a first palladin isoform from a second palladin isoform (e.g. gel filtration columns, antibody columns, size exclusion columns, membrane cut-off spin columns etc.).

EXAMPLES

[0409] The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the present invention, and are not intended to limit the scope of what the inventors regard as their invention nor are they intended to represent that the experiments below are all or the only experiments performed. Efforts have been made to ensure accuracy with respect to numbers used (e.g. amounts, temperature, etc.) but some experimental errors and deviations should be accounted for. Unless indicated otherwise, parts are parts by weight, molecular weight is weight average molecular weight, temperature is in degrees Celsius, and pressure is at or near atmospheric. Standard abbreviations may be used, e.g., bp, base pair(s); kb, kilobase(s); pl, picoliter(s); U, units; s or sec, second(s); min, minute(s); h or hr, hour(s); aa, amino acid(s); kb, kilobase(s); bp, base pair(s); nt, nucleotide(s); i.m., intramuscular(ly); i.p., intraperitoneal(ly); s.c., subcutaneous(ly); and the like.

Example 1

Identification and Characterization of a Palladin Gene Mutation

Summary

[0410] A large kindred (Family X) was identified, which inherits pancreatic cancer in an autosomal dominant fashion with high penetrance. In order to identify living affected familial members prior to the onset of cancer, an endoscopic surveillance program was developed that assists in the detection of pancreatic pre-cancer (also called pancreatic intraepithelial neoplasia or PanIN). Family X includes 18 cases in 4 generations of either cancer (n=9) or histologically proven pre-cancer (n=9) (FIG. 2). In FIG. 2, nine members of this family were diagnosed with pancreatic cancer, and nine with pancreatic precancer (five with carcinoma in situ (PanIN 3); and four with low-grade dysplasia (PanIN 2)).

[0411] A pancreatic cancer susceptibility locus was mapped to 4q32-34. Identification of the gene of interest in this region represented a considerable challenge because of the size (16 MB) and the number of genes localized to this region (approximately 250). To narrow the region, a custom microarray of the genes and expressed sequence tags located at 4q32-34 was created. Palladin, one of the candidate genes that showed the highest up-regulation in the array analysis, was assessed further and a C to T base pair substitution was identified in the highly conserved region of palladin in all of the affected family members. The Palladin gene is highly conserved through evolution; it is a cytoskeletal protein and a component of actin-containing microfilaments that control cell shape, adhesion, and motility.

Methods

Tissues and RNAs

[0412] Pancreatic adenocarcinoma tissue samples were collected from the University of Washington, Cleveland Clinic, and University of Pittsburgh with authorized IRB approval. Details of tissue collection, RNA preparation and RNA labeling for microarrays were previously described (Pogue-Geile et al. (2004) *Cancer Genomics and Proteomics* 1:371-386). Normal pancreas samples were obtained commercially from Stratagene, Clontech, Biochain, Chemicon, and Ambion and from normal pancreatic surgical specimens obtained through the University of Washington. In case of the surgical specimens, the histologic normalcy of the tissue was ascertained by pathologist Dr. Mary Bronner and for tissues obtained at the University of Pittsburgh by pathologist at the University of Pittsburgh.

Microarray Construction Hybridization and Analysis

[0413] Unigenes were identified using the UCSC genome maps (available on the internet at genome.ucsc.edu; Kent et al) and NCBI maps. 243 unigenes clones mapping between markers D4S413-D4S299 were identified. Seventeen clones were included as housekeeping genes and 25 additional clones were included because of their location in a region of the genome that is frequently lost in pancreatic cancer. Clones were acquired from Research Genetics, (now Invitrogen) and from RZPD (German Resource Center for Genome Research) and from University of Pittsburgh Genomics and Proteomics Core Laboratories. Identification of all clones and accurate location on chromosome 4q was confirmed by

sequence analysis using Big Dye. Clones were amplified, purified and arrayed onto glass slides as previously described (Pogue-Geile et al. (2004) *Cancer Genomics Proteomics* 1:371).

[0414] Slides were scanned with the GMS 418 scanner (Genetic Microsystems). The Cy-5 and Cy-3 images were overlaid, and raw data was generated for both channels using the ImaGene program United (Bioinformatics Inc, Calgary AB).

QRT-PCR

[0415] QRT-PCR was carried out using the 5' nuclease assay and an Applied Biosystems 7700 Sequence Detection Instrument (TaqMan). Tissues used for expression analysis utilized whole tissues. Palladin expression was measured relative to the endogenous control, GAPDH, using the comparative C_T method described previously (Godfrey et al. 2000). cDNAs were generated at 2 different RNA input concentrations (10 ng/ μ l and 4 ng/ μ l) and TaqMan reactions with the endogenous control were run in duplicate from both RT reactions. Palladin TaqMan reactions were carried out in triplicate using RT reactions with 10 ng/ μ l reactions. A calibrator RNA, composed of 50 ng/ μ l universal reference RNA (Stratagene, cat no 40000-41) and 50 ng/ μ l colon RNA (Ambion cat no 7986) was included on every amplification plate to allow comparison of samples run at different times. RT-negative controls were run on each plate to ensure that no amplification occurred in the absence of cDNA.

Statistical Analysis

[0416] Microarray analysis was performed using the Gene Expression Differential Analysis tool (caGEDA) a web application specifically developed for cancer microarray data analysis at [http://bioinformatics\(dot\)upmc\(dot\)edu/GE2/GEDA\(dot\)html](http://bioinformatics(dot)upmc(dot)edu/GE2/GEDA(dot)html) (Patel and Lyons-Weiller). The J5 test was used; and the data were normalized with log₂ and Z transformation.

Sequence Analysis

[0417] A restriction enzyme assay can be used to detect the specific family X mutation. First, PCR was performed with Fast Start Taq from Roche using 30 μ g genomic DNA, with 200 μ M dNTPs, 20% GC-rich solution, 0.04 U/ μ L Taq polymerase, 1 \times Buffer, 200 nM forward primer, 200 nM reverse primer. Forward primer: 5'-CCGACGTGTTCCCACTGC (SEQ ID NO:84; corresponding to nucleotides 140200 to 140217 of *Homo sapiens* BAC clone RP11-635L1; see FIG. 16) and reverse primer: 5'-CGCACGGAGAGAAATGTGTG (SEQ ID NO:85; corresponding to the complement of nucleotides 140510 to 140529 of *Homo sapiens* BAC clone RP11-635L1; see FIG. 16 (SEQ ID NO:100)). Conditions were as follows: 5-min denaturation step at 95 $^{\circ}$ C., 40 cycles of 30s at 95 $^{\circ}$ C., 30s of annealing at 54 $^{\circ}$ C., and 30s extension at 72 $^{\circ}$ C., followed by a final extension for 7 min at 72 $^{\circ}$ C. The 330 bp PCR products were separated on 1.5% agarose gels and visualized with syber green and UV light. Ten μ L of each PCR product was digested using AatII from NEB in 1 \times digestion buffer for 1 hour at 37 $^{\circ}$ C. Amplicons containing the C to T mutation cut once giving 102 bp and 228 bp bands, while the amplicons from wild type DNA will not be cut by this restriction enzyme. This 330 bp fragment can also be sequenced using the PCR primers listed above.

Constructs and Transfection

[0418] The human wild-type (WT) palladin construct was made by PCR-cloning the entire coding sequence from a

human palladin cDNA clone (human cDNA clone hk07554) into the sites of EcoRI and BamHI of pHRGFP IIN vector (Stratagene), downstream of, and in frame with, the green fluorescent protein (GFP) tag. To create the Family X mutant construct (FX), Quick-Change Multi Site-Directed Mutagenesis (Stratagene) was used, with primers centered at the Family X mutation (P239S), according to manufacturer's protocol. Briefly, (1) 100 ng of WT construct was added as a template to a PCR cocktail containing 2.5 μ l of 10 \times mutagenesis buffer, 100 ng of mutagenesis primer containing the Family X mutation, 250 μ M each dNTP, 1 μ l QuickChange enzyme blend and 1 μ l QuickSolution. The PCR cycling parameters were one cycle of: 1 min at 95 $^{\circ}$ C., followed by 30 cycles of 1 min at 95 $^{\circ}$ C., and 15 min at 65 $^{\circ}$ C.; (2) the parental template DNA was treated with DpnI (10 U) at 37 $^{\circ}$ C. for 60 min; (3) 4 μ l of this reaction was transformed into XL10 Gold Ultracompetent cells (Stratagene); and (4) several clones were chosen for PCR and/sequencing to confirm the incorporation of the mutation.

HeLa Cell Transfection

[0419] Transfection was performed using a Fugene kit (Roche Diagnostics) on human cancer epithelial cell lines (HeLa cells) according to the manufacturer's protocol. Briefly, (1) one day before the transfection, the cells were distributed into a six-well plate so that they would be approximately 70% confluent the next day; (2) 3-6 μ l of transfection reagent was mixed with 94-97 μ l of DMEM and left for 5 min; (3) then 1 μ g of construct (either WT or FX mutant construct) was added to the complex and left for 10 min; (4) the complex was then added to the cells in a dropwise fashion; and (5) the expression of the GFP construct was observed the next day. Fluorescence staining was performed on cells fixed in 3.7% formaldehyde and permeabilized with 0.1% Triton X-100. The cells were then stained with 50 μ g/ml of TRITC-phalloidin (Sigma) for 40 min, followed by washing with PBS. Finally, DAPI (Sigma) was added at 10 μ g/ml to stain the nucleus, and antifade reagent (Invitrogen) was used to prevent auto-color bleaching. The construct/transfection experiments were run in triplicate and assessed blindly.

Analysis of Protein Levels

[0420] Size fractionation (SDS-PAGE) was performed on 20 μ g of protein from each pancreatic cell line sample. The samples were individually loaded onto a gel, separated through electrophoresis, and then blotted onto a nitrocellulose membrane according to manufacturer's protocol (Amersham Biosciences, Piscataway, N.J., United States). The polyclonal palladin antibody (ab 621; Otey et al. (2005) *Int Rev Cytol.* 246:31-58) was used. A 1:2,000 dilution was used for Western blotting. The ECL plus kit (Amersham Biosciences) was used to detect protein in the Western blot.

Results

[0421] It was hypothesized that a mutated gene is best detected in very early pre-cancerous tissue, where the mutated gene has initiated neoplastic progression but before the genetic chaos of cancer has occurred. Using RNA from the pre-cancerous pancreatic tissue from a Family X member, the gene expression in the 4q32-34 region was profiled on a custom microarray. The custom array was created using 243 sequence-verified Unigenes clones located between markers D4S2976 to D4S415, a region slightly larger than the region identified as the pancreatic cancer susceptibility locus (between markers D4S413-D4S2991). Unigenes represent a non-redundant set of gene-oriented clusters of cDNA clones

assigned to a unique gene and genomic location. In addition to the Family X pre-cancer RNA, 10 different sporadic pancreatic cancers and 2 normal donor pancreata were also profiled. Samples were spotted in sextuplicates (6x) (FIGS. 3A-C). Comparative expression in Family X and the 10 pancreatic cancer samples were rank ordered by their degree of over or under-expression. Two Palladin clones were shown to be the most highly differentially expressed in sporadic pancreatic cancers and were also over expressed in Family X, using a J5 analysis which was used previously. Patel et al. (2004) *Appl. Bioinf.* 3:49-62. The J5 test compares the difference in mean expression for a given gene to the magnitude (absolute value) of mean difference in all of the genes on an array. Palladin clones were also shown to be among the most differentially expressed genes when Family X gene expression was analyzed by N-fold analysis.

[0422] To validate the findings on the custom microarray, commercially available primers directed toward exons 9 and 10 were used to measure gene expression of the Palladin gene (FIGS. 3A-C). Expression analysis via quantitative RT-PCR (qRT-PCR) was performed in a) 16 sporadic pancreatic cancers, b) 4 pre-cancerous tissues (2 from Family X and 2 from other familial pancreatic cancer families) c) 9 histologically normal appearing tissues adjacent to sporadic pancreatic cancers (normal adjacent) and d) 6 normal pancreas samples. Although the sample size was small, a significant difference was detected between all cancer and pre-cancer groups and normal pancreata (One way-4 group ANOVA $F(3,35)=5.86$, $p=0.003$). All possible pair-wise tests using Least Significant Difference test ($p<0.05$) showed significant differences between all cancer and pre-cancer groups (cancer, normal adjacent, PanIN) compared with normal. Palladin was over-expressed in most of the cancerous and precancerous pancreatic tissue, as well as in the histologically normal appearing tissue adjacent to the cancers, when compared to normal. This expression analysis indicates that Palladin is likely over-expressed very early in the development of pancreatic cancer—including both Family X and sporadic forms of the disease.

[0423] FIGS. 3A-C. Identification of 4q32-34 Genes Differentially Expressed in Sporadic Pancreatic Cancer and In Family X Pre-Cancer. A). The Gene Discovery Array, a custom 4q32-34 microarray was interrogated using RNA derived from 10 sporadic pancreatic cancers, 1 pre-cancerous pancreas from an Family X individual and 2 normal donor pancreas samples. cDNAs were spotted six times. B). The top 2 most differentially expressed genes among 10 sporadic pancreatic cancers and the corresponding differential expression in Family X are shown. C). Average expression of Palladin using commercially available probes to Palladin exons 9 and 10. Samples tested included 6 normal pancreas samples, two from Stratagene and one each from Ambion, Biochain, Chemicon and Clontech, 4 PanIN III tissues (2 Family X individuals and 2 non-Family X individuals), 9 histologically normal tissue adjacent to cancer and 16 pancreatic cancer tissues. Error bars indicate 1 standard deviation above and below the average.

A Base Pair Substitution in Palladin Tracks with the Affected Members in Family X and Causes an Amino Acid Change in the Actinin Binding Site.

[0424] Palladin is an extremely large gene spanning 432 Kb, with a total of 31 exons, up to 9 probable alternative promoters, and at least 8 alternatively spliced transcripts. The published literature has described at least 3 major different isoforms of Palladin ranging in molecular weight from 140 kD, 120 kD and 90 kD. The smallest isoform (90 kD) is a constituent of the larger isoforms.

[0425] Twenty-nine individuals from Family X were available for mutational testing of Palladin: 9 unaffected, 12 with pancreatic cancer or pre-cancer, 9 known to be unaffected, and 8 with unknown status. Sequence analysis of Palladin identified a C to T base pair substitution in exon 2 of all 12 affected family members and none of the 9 unaffected family members. The C to T transition causes a proline (hydrophobic) to serine (hydrophilic) amino acid change at position 715 in Palladin RNA (FIG. 4).

[0426] FIG. 4. Location and identification of the Family X mutation. Top black bar indicates the genomic location of the Family X mutation between microsatellite markers, D4S413 and D4S299, on chromosome 4 (Eberle et al. 2002) *Am. J. Hum. Genet.* 70:1044). Center line designates the location of the palladin transcript, AB023209, with vertical boxes and lines indicating exons. At the bottom a small portion of the DNA sequence is shown with a C to T transition at position 715 (indicated with an arrow) which causes a proline (hydrophobic) to serine (hydrophilic) amino acid change at amino acid 236.

[0427] The mutation discovered in Family X is contained in the cDNA clone AB023209 (FIGS. 1A and B), which encodes a 4,349 nucleotide transcript, containing 12 exons and encoding a 772 amino acid protein (90 kD isoform).

[0428] The mutation occurs in exon 2 of this Palladin isoform (peptide beginning MSALA, 126 amino acids downstream 90 kD start site).

[0429] This base pair substitution is not a known single nucleotide polymorphism, nor was it detected in the blood of 294 of 295 non-pancreatic cancer controls (589 of 590 alleles). One control blood sample revealed the mutation; unfortunately, the medical history of this individual is unknown. The Family X mutation occurs in a region that is highly conserved across species (FIG. 5).

[0430] FIG. 5. The binding site of Actinin to Palladin is highly conserved across species. The mutation in Family X causes a proline (hydrophobic) to serine (hydrophilic) change in the amino acid sequence.

The 90 Kda Isoform of Palladin is the Major Isoform Expressed in Human Pancreatic Epithelium.

[0431] As shown in FIG. 6, at least three major isoforms of palladin are known, with molecular weights of 200 kDa, 140 kDa, and 90 kDa. The smallest isoform (90 kDa) is a component of the larger isoforms. Western blot analysis suggests that the 90 kD is the major isoform expressed in human pancreas (FIG. 6).

[0432] Western blotting and a polyclonal antibody (ab 621) were used to assess the protein isoform expressed in pancreatic ductal epithelium. The results indicated that the 90 kDa isoform is the predominant form of palladin in pancreatic ductal epithelium and in pancreatic cancer cell lines.

[0433] FIG. 6. The 90 kD palladin is the major isoform expressed in human pancreas. Shown here is the western blot of the proteins extracted from various human cells detected with polyclonal antibody against palladin raised in rabbit (Ab 621). The major isoform expressed in human pancreas tissue and cultured epithelial cells (HPDE: human pancreatic ductal epithelial cell line) is the 90 kD isoform (e.g., arrow in "HPDE" lane). In the control sample (MSC: human mesenchymal stem cells), besides the major 90 kD isoform, the 140 kD isoform is also detectable (arrow in "MSC" lane). Palladin does not Fit the Model of a Classical Tumor Suppressor Gene.

[0434] Loss of heterozygosity (LOH) studies have shown that a region in 4q32-35 is lost in many different cancers, including cancers of the pancreas, breast, colon, liver. Com-

parative genomic hybridization of pancreatic cancers using cDNA microarrays defined 3 different regions of LOH located between 4q32-35, suggesting that tumor suppressor gene might be localized to this region. Tumor suppressor genes lose activity when both alleles are inactivated: in many classical familial syndromes, one allele is inactivated through an inherited mutation and the second allele subsequently becomes lost, initiating neoplastic progression. To determine whether Palladin has genetic characteristics consistent with a classical tumor suppressor gene, a search for loss of a Palladin allele was conducted. A custom-made copy number chip (Nimblegen) was used to measure the number of alleles in pre-cancerous (PanIN II and III) pancreatic tissues from Family X members. The entire Palladin gene was interrogated and the copy number was 2 in every instance. In addition, two Palladin SNP probes with a minor allele frequency greater than or equal to 0.47 were used in TaqMan SNP analysis to test 21 sporadic pancreatic cancers for evidence of LOH. Ten sporadic pancreatic cancer tissue samples were heterozygous and thus, informative. None of these 10 samples showed loss of either allele in the tumor. These findings suggest that mutated Palladin is probably not a tumor suppressor gene, but rather may act as an oncogene.

Palladin is Increasingly Over-Expressed In Ductal Epithelial Cells with Neoplastic Progression.

[0435] Pancreatic cancer tumors are a mixture of tumor cells and a variety of other cell types due to the strong desmoplastic change common to pancreatic cancers. To determine whether the Palladin gene is expressed in the epithelial cells, the expression in primary human ductal pancreatic epithelial (HPDE) cells was examined with qRT-PCR. Epithelial cells were isolated from pancreatic tissue and grown in primary culture. Little to no expression could be detected in cells from a normal pancreatic ductal epithelium; in contrast, increasing expression was seen in epithelial cells derived from 2 different affected Family X individuals who had pre-cancer, but not cancer. The greatest expression was seen in ductal epithelial cells taken from an individual with a case of sporadic pancreatic cancer. The overexpression of Palladin increased as the epithelial cells became increasingly neoplastic (Pan I < PanIN III < Cancer) (FIG. 7).

[0436] FIG. 7. Palladin expression in human ductal pancreatic epithelial (HPDE) primary cultures increases with neoplastic progression from normal (HPDE Norm) to PanIN I to PanIN III to cancer (HPDE PC). Each bar represents HPDE epithelial cultures from one person. The PanIN I and PanIN III lesions are from affected members of Family X. The pancreatic cancer epithelial cells are from a sporadic cancer.

[0437] Palladin is an actin-binding protein that controls cytoskeletal formation and cell movement. As depicted schematically in FIG. 8A, palladin binds other key proteins including actinin (the site of the Family X mutation), and ezrin. The pro→ser mutation identified in Family X occurs directly in the binding site for actinin (codons 243-258). FIG. 8B presents a schematic depiction of various palladin isoforms (90 kDa, 140 kDa, and 200 kDa). Proline-rich domains are designated PR1 and PR2, and the immunoglobulin-like (Ig-like) domains are numbered 1-5. The arrow marks the location of the Family X mutation.

Palladin and Alpha-Actinin Proteins are Abnormally Expressed in Sporadic Pancreatic Cancer Cell Lines.

[0438] Protein expression analysis of sporadic pancreatic cancer cell lines was performed, using protein (“western”) blotting. The sporadic pancreatic cancer cell lines analyzed were PaTu1, PaTu2, Panc-1 (American Type Culture Collection (ATCC) accession number CRL-1469), FA6 (Morgan et

al. ((2006) *Mol. Cancer* 5:1), MiaPaca2, and HPAF-II (ATCC CRL-1997). PaTu1 and PaTu2 are described in, e.g., Missiaglia et al. ((2004) *Int'l. J. Cancer* 112:100-112); FA6 is described in, e.g., Morgan et al. ((2006) *Mol. Cancer* 5:1). Abnormal palladin protein levels were detected in five of the six cell lines.

[0439] The 90 kDa palladin isoform co-localizes in stress fibers with several cytoskeletal components, such as actin and alpha-actinin. Protein expression of these cytoskeletal components in pancreatic cancer cell lines was examined using Western blotting and a polyclonal antibody (ab 621) to palladin. As shown in FIG. 9, of seven sporadic pancreatic cancer cell lines (FA6, HPAF, IMIMPC2, SUIT2, HS766T, PANC-1, and PATU2), five showed clear protein overexpression of palladin. Cell lines FA6, HPAF, IMIMPC2, SUIT2, and PATU2 overexpress palladin protein compared to the minimal expression in normal HPDE; alpha-actinin was also abnormally expressed in some of the pancreatic cancer cell lines, especially in PANC-1, a line that did not show overexpression changes in palladin protein.

[0440] These data suggest that abnormal expression of palladin or its binding partner, alpha-actinin, is present in six of the seven pancreatic cancer cell lines tested.

[0441] FIG. 9. Abnormal protein expression of palladin and α -actinin proteins in sporadic pancreatic cancer cell lines. Human mesenchymal cell lysate (hMCL) was used as a positive control for palladin overexpression. FA6, HPAF, IMIMPC2, SUIT2, and PATU2 are sporadic pancreatic cancer cell lines that overexpress palladin protein compared to the minimal expression evident in normal HPDE. Alpha-actinin is also abnormally expressed in some of the pancreatic cancer cell lines compared to normal pancreas—especially PANC-1, one of the few pancreatic cell lines that does not show expression changes in palladin protein.

Overexpression of Mutant Palladin Results in Different Cellular Cytoskeletal Phenotypes

[0442] It has been previously shown that palladin (90 kD isoform) colocalizes in the stress fiber with several cytoskeletal components, such as actin, actinin, ezrin, and Ena/VASP^{5,7,8}. To investigate if the mutant form of palladin exhibits abnormality in this function, the human 90 kD isoform sequence was cloned into vector with GFP downstream, designated as wildtype construct. Site-directed mutagenesis was used to create a mutation of C to T at position 715 bp, resembling the Family X mutation, designated as FX construct. These constructs were transfected into HeLa cells to study the overexpression phenotypes. Cells transfected with wildtype construct display well organized actin bundle, while GFP-palladin (wildtype) completely co-localize in the stress fiber with actin. When cells were transfected with FX construct, cells often exhibit almost normal actin bundle, however, GFP-mutant palladin frequently forms aggregates in these cells. Perhaps the most distinct phenotype for FX construct transfection is the presence of cells with multiple small blobs that do not colocalize with actin. In contrast, cells transfected with GFP empty vector exhibited green in the whole cells. These results suggest that mutant palladin may lose its ability to participate in regulating the actin networks, thus may impair the organization of normal cytoskeleton given the important role of palladin in the cytoskeleton.

Example 2

Additional Palladin Gene Mutations

[0443] Additional palladin gene mutations were identified: one in a pancreatic cancer cell line, a second in a family,

designated Family U (or Utah Family 1); and a third in two other kindreds (Utah Family 2 and Seattle Family 2).

Materials and Methods

Mutant Palladin Construct

[0444] The human wild-type (WT) palladin construct was made by PCR-cloning the entire coding sequence from a human palladin cDNA clone (human cDNA clone hk07554) into the sites of EcoRI and BamHI of pHRGFP IIN vector (Stratagene), downstream of, and in frame with, the green fluorescent protein (GFP) tag. To create a construct with the G→T mutation at position 1671, Quick-Change Multi Site-Directed Mutagenesis (Stratagene) was used with primers centered at the G→T mutation at position 1671, according to manufacturer's protocol. Briefly, (1) 100 ng of WT construct was added as a template to a PCR cocktail containing 2.5 μl of 10× mutagenesis buffer, 100 ng of mutagenesis primer containing the G→T mutation at position 1671, 250 μM each dNTP, 11 QuickChange enzyme blend and 1 μl QuickSolution. The PCR cycling parameters were one cycle of: 1 min at 95° C., followed by 30 cycles of 1 min at 95° C., and 15 min at 65° C.; (2) the parental template DNA was treated with DpnI (10 U) at 37° C. for 60 min; (3) 4 μl of this reaction was transformed into XL10 Gold Ultracompetent cells (Stratagene); and (4) several clones were chosen for PCR and sequencing to confirm the incorporation of the mutation.

[0445] Transfection of HeLa cells was performed as described in Example 1. The transwell migration assay was performed essentially as described in Example 1.

Analysis of Protein Levels

[0446] Size fractionation (SDS-PAGE) was performed on 20 μg of protein from each pancreatic cell line sample. The samples were individually loaded onto a gel, separated through electrophoresis, and then blotted onto a nitrocellulose membrane according to manufacturer's protocol (Amersham Biosciences, Piscataway, N.J., United States). The polyclonal palladin antibody (ab 621; Otey et al. (2005) *Int Rev Cytol.* 246:31-58) was used. A 1:2,000 dilution was used for Western blotting. The ECL plus kit (Amersham Biosciences) was used to detect protein in the Western blot.

Results

1) Pancreatic Cancer Cell Line

[0447] The palladin gene in cell line PaTu2 was analyzed.

The Palladin Gene in PaTu2 Contains a Mutation.

[0448] It was found that a mutation occurs in the following sequence:

(SEQ ID NO: 86)

GTCAAGTGGGTTACCAACCCAGATCTAAGCTGNCAACTAGATGGAAAGCC
CGTACGCCCTGACAGTGCTCACAAGATGCTGGTGCGTGAGAACGGGGTGC
ACTCTCTGATCATAGAGCCAGTCACGTACGTGATGCCGGCATCTACACA
TGATAGCTACCAACCCAGCAGGACAGAACTCATTACGCCCTGGAGCTTGT
GGTTGCTG,

[0449] where N=G/T at position 1671 of AB023209 in the ninth exon (see, e.g., GenBank AB023209; and FIGS. 1A and 1B). The wild-type sequence at position 1671 of AB023209 is G; the mutant has a T at position 1671. The G→T substitution

results in a change in codon sequence of TGG→TGT, leading to a Trp→Cys mutation at amino acid 557 of palladin (see GenBank Accession No. BAA76836.1; and FIG. 12; SEQ ID NO:87):

[0450] VSGLPTPDL~~S~~XL~~D~~GKPV~~R~~PDS~~A~~HK~~M~~-LVRENGVHSLIIEP (SEQ ID NO:88).

[0451] The Trp→Cys mutation occurs in a conserved region, the Ig2 domain of palladin:

Human	(SEQ ID NO:89)
VSGLPTPDL S XL D GKPV R PDS A HK M LVRENGVHSLIIEP	
Mouse	(SEQ ID NO:90)
VSGLPTPDL S XL D GKPI R PDS A HK M LVRENGVHSLIIEP	

[0452] Primers suitable for use in amplifying a sequence containing the G→T substitution at 1671 of AB023209 include the following:

Forward primer:
5' TCTTGTACTACTGAAGGAGGAATTTATGC; (SEQ ID NO:91)
and

Reverse primer:
5' TTTCCGTATTGGTTAGTAATGTAGAATTAG. (SEQ ID NO:92)

[0453] The position of the forward and reverse primers in genomic DNA is shown in FIG. 14, where the nucleotides corresponding to the primer sequences are underlined, exon 9 is shown in bold text, and the G→T substitution is underlined and in bold. The sequence amplified by these primers corresponds to nucleotides 183428 to 183828 of the *Homo sapiens* BAC clone RP11-635L1 as set forth in GenBank Accession No. AC080188.

[0454] The above forward and reverse primers result in an amplification product that contains either wild-type G at position 1671 of AB023209 or mutation T at position 1671 of AB023209. An amplification product containing the G→T mutation can be cut with HindIII; an amplification product containing the wild-type G at position 1671 of AB023209 is not cut with HindIII.

The W557C Mutation Alters Palladin Function.

[0455] An expression construct was made that includes a nucleotide sequence encoding the 90 kD isoform of palladin, and including the G→T mutation at position 1671, resulting in a Trp→Cys substitution at amino acid 557 (W557C). The expression construct included a green fluorescent protein (GFP)-coding sequence in-frame with the palladin-encoding nucleotide sequence. The construct was transfected into HeLa cells. HeLa cells expressing the mutant palladin/GFP protein exhibited cytoskeletal abnormalities. HeLa cells expressing the W557C mutant palladin/GFP protein exhibited a 200% increase in mobility in a transwell assay, compared to HeLa cells containing a control expression construct that included a nucleotide sequence encoding wild-type palladin 90 kD isoform.

2) Utah Family 1 Mutation

[0456] A family, referred to as "Utah Family 1," was analyzed for the presence of mutations in the palladin gene. A mutation occurs in the following palladin sequence:

(SEQ ID NO:93)
 CCAACACTGAAAACCTGTAGTTACGAGTCAATGGGAGAAATCCAACAATGAC
 CACTTCCAACACTTTCCACCTCCCCTCCAATCTGGAGACAAGTTCCTT
 GGAGTTGGCTTCAAAGAAACCATCTGAGATCCAGCAGGTGAACAACCTG
 AGTTAGGCTTGGAGAGGAGCCCTTCAAATGCAATTCATGCTGCTGAG
 AGGGAACGAAACGGAGTCCATCCAGCNGTGGAGTAAATGGACTGATTAA
 CGGCAAGACTAACAGTATAAATCTCTTCCAACACCAGCTGTCTGCTTT
 CACCCACTAAGGAGCCACCACCTCTGCTTGCCAAACCAAAACT,

[0457] where N+C/T at position 2060 of NM_016081 in the tenth exon (see GenBank NM_016081; and FIGS. 13A and 13B). The wild-type sequence has a C at position 2060 of NM_016081; and the mutant has a T at position 2060 of NM_016081. The C→T substitution results in a change in codon sequence of CGT→TGT, leading to an Arg→Cys substitution at amino acid 617 of palladin (see GenBank Accession No. NP_057165.3; FIG. 12 (SEQ ID NO:87); and FIGS. 13A and 13B (SEQ ID NO:20)):

(SEQ ID NO:94)
 NTENCYESMGESNNDHFQHFPPPPPILETSSLELASKKPSSEIQVNNPE
 LGLSRAALQMQFNAAERETNGVHPSXGVNGLINGKANSKSLPTPAVLLS
 PTKEPPPLLAKPK.

[0458] The Arg→Cys mutation occurs in a conserved region of palladin, the poly-Pro region:

Human: AERETNGVHPSRGVNGLINGKA (SEQ ID NO:95)

Mouse: AERETNGVHPSHGVNGLINGKA. (SEQ ID NO:96)

[0459] Primers suitable for use in amplifying a sequence containing the C→T substitution at position 2060 of NM_016081 include the following:

Forward primer:
 5' CACAACACAGGGATTCTCAGAAGA; (SEQ ID NO:97)
 and

Reverse primer:
 5' AGCTGGTGTGGAAGAGATT. (SEQ ID NO:98)

[0460] The position of the forward and reverse primers in genomic DNA is shown in FIG. 15, where the nucleotides corresponding to the primer sequences are underlined, and the C→T substitution is underlined and in bold. The sequence amplified by these primers corresponds to nucleotides 26927 to 27313 of the *Homo sapiens* BAC clone RP11-592K15 as set forth in GenBank Accession No. AC084353.

[0461] The above forward and reverse primers result in an amplification product that contains either wild-type c at position 2060 of NM_016081 or mutation T at position 2060 of NM_016081. An amplification product containing the C→T mutation can be cut with PvuII; an amplification product containing the wild-type C at position 2060 of NM_016081 is not cut with PvuII.

3) Utah Family 2 and Seattle Family 2 Mutation

[0462] A family, referred to as "Utah Family 2," was analyzed for the presence of mutations in the palladin gene. A 12-bp insertion was found in the putative promoter region of the 90 kD isoform of palladin. Utah Family 2 had three members with pancreatic cancer, all cousins, as well as a family member with esophageal cancer and a member with breast cancer.

[0463] The same 12-bp insertion was detected in one of nine Seattle FPC kindreds tested, but was absent from 198 palladin alleles from normal control DNA. FIG. 19 presents a nucleotide sequence showing the promoter of the palladin gene encoding the 90 kDa isoform of palladin (boxed sequence), with a 12-bp insertion (bold sequence).

[0464] Table 1, below, presents a summary of mutations associated with pancreatic cancer.

TABLE 1

Nt substitution (Codon)	Position	GenBank nt (GenBank protein)	Amino acid substitution	Position (Figure; SEQ ID)	Protein
C→T CCC→TCC	cDNA nt 715 (FIG. 1; SEQ ID NO: 1) Exon 2 Boundaries: <u>nt132-803</u> Genomic AC080188 Nt 140301 FIG. 16 (SEQ ID NO: 100)	cDNA AB023209 <u>(BAA76836.1)</u> genomic AC080188 Nt 139718- 140388 FIG. 16	Pro→Ser	aa 239; FIG. 11 (SEQ ID NO: 99)	90 kD isoform
G→T	cDNA nt 1671 (FIG. 1; SEQ ID NO: 1) exon 9	cDNA AB023209 <u>(BAA76836.1)</u> genomic Ac080188 Nucleotides	Trp→Cys	aa 557 FIG. 11 (SEQ ID NO: 99)	90 kD isoform

TABLE 1-continued

Nt substitution (Codon)	Position	GenBank nt (GenBank protein)	Amino acid substitution	Position (Figure; SEQ ID)	Protein
	boundaries: 1639-1846	183529-183736			
	genomic	FIG. 14			
	nt 183561	(SEQ ID			
	FIG. 14	NO: 101)			
	(SEQ ID				
	NO: 101)				
C→T	cDNA	cDNA	Arg→Cys	aa 617	140 kD isoform
	nt 2060	NM_016081		FIG. 12	
	(FIG. 13;	(NP_057165.3)		(SEQ ID	
	SEQ ID	genomic		NO: 87)	
	NO: 20)	AC084353			
	exon 10	FIG. 15			
	boundaries: 1833-2175	(SEQ ID			
	genomic	NO: 102)			
	nt 27251				
	FIG. 15				
	(SEQ ID				
	NO: 102)				

Example 3

Palladin Gene Expression in White Blood Cells

[0465] Palladin mRNA levels were assessed in white blood cells (WBC) from normal individuals (e.g., individuals without pancreatic cancer), and from individuals with pancreatic cancer. The levels were normalized to the level of glyceraldehydes-3-phosphate dehydrogenase (GAPDH) mRNA present in the cells and to a standard sample following the ddCt method described previously (Pogue-Geile et al. (2004) *Cancer Genomics and Proteomics* 1:371-386). The data shown in FIG. 17, show the level of palladin mRNA in WBC from normal individuals (Nor-M1, -F1, F2, F4, RS28, RS45, RS48, RS72, and RS98) were higher than that in WBC from individuals with pancreatic cancer. The level of normalized palladin mRNA in WBC from individuals with pancreatic cancer was from about 0.4 to about 1.2 log lower than that in individuals without cancer. Thus, the level of palladin mRNA in WBC from individuals with pancreatic cancer (PG6, PG8, PG9, PG10, PG14, PG18, PG19, PG21, PA17, and PA25) was abnormally low.

Example 4

Palladin Expression in Various Cancers

[0466] Various cancer tissues were tested for reactivity with 1E6, a monoclonal antibody specific for the 90 kDa isoform of palladin. Rachlin and Otey (2006) *J. Cell Sci.* 119:995-1004. The data, shown in Table 2, below, indicate that palladin is over-expressed in at least breast cancer and in head and neck squamous cell cancer. The "1E6 grade" indicates, on a scale of 1⁺-3⁺, the level of antibody binding and thus the level of palladin protein over-expression.

TABLE 2

Cases positive	1E6 grade	Cancer type
0/31	—	Lung cancer, both small cell and non-small cell types

TABLE 2-continued

Cases positive	1E6 grade	Cancer type
0/15	—	Renal cell
2/15	1 ⁺	Lobular breast cancer stroma
3/7	1 ⁺	Ductal breast cancer stroma
1/18	—	Colorectal cancer
0/18	—	Ovarian cancer
13/15	1 ⁺ to 2 ⁺	Head and neck squamous cell cancer stroma
8/12	1 ⁺ to 3 ⁺	Pancreatic ductal cancer stroma
0/15	—	Melanoma
0/15	—	Gastric cancer
1/15	—	Urothelial cancer (note: epithelial cells stained)

Example 5

Effect of Palladin Mutation on Mobility

Cell Migration Assay

[0467] HeLa cells were first transfected with constructs of WT, FX, and empty vector respectively for 24 h, and then sorted for GFP-positive cells. The sorted cells continued to grow for 2 d before they were seeded onto a transwell plate for the migration assay. Migration assays were performed on 24-well Transwell cell culture chambers (Corning Costar Corporation, Cambridge, Mass., United States) fitted with multiporous (8-µm pore size) polycarbonate membranes. The upper chambers of the membrane were coated with fibronectin (10 µg/ml in PBS) 24-48 h before the assay. The upper chamber was then filled with 40,000 cells in suspension with 4001 of medium (DMEM with 10% FCS), and the lower chamber was filled with 500 µl of the same medium. Plates were placed in a humidified CO₂ incubator at 37° C. for 17 h. Membrane inserts were then removed, fixed by immersing in ethanol five times with 1 s durations, and stained with 0.5% crystal violet dye (Sigma; #C3886) in 20% methanol for 30 min. After gentle rinsing with water, the nonmigratory cells on the upper surface of the membrane were removed with

cotton swabs, leaving only the migratory cells within the membrane. The membrane insert was left to dry overnight and then placed in a 96-well plate. The dye was extracted with 30% acetic acid and the plate was placed on a shaker for 10 min to allow the dye to dissolve completely. The calorimetric absorbance was assessed at 590 nm. A transwell without cells was used as a background control. To correct for the cell numbers seeded in each transwell experiment, the optical density of all the original cells in the transwell (migrated and nonmigrated) was measured for each transfected cell type sample. By comparing the optical density from each sample type, a correction factor was provided for any variation in starting cell number between samples. Each sample was done in triplicate.

[0468] Mobility is Increased in Cells that Express Mutant Palladin.

[0469] It was hypothesized that the cytoskeletal changes in the cells containing the mutated Family X construct might provide a cancer phenotype, specifically increased cell motility. The cytoskeleton is essential for cell movement, and cell motility is important for the invasive nature of cancer cells. HeLa cells transfected with one of three constructs of palladin

(FX, WT construct, or an empty vector) were individually plated onto a fibronectin-coated membrane in a transwell chamber, and a migration assay was performed in a standard fashion. Hu et al. (2006) *FASEB J.* 20:1892-1894.

[0470] The cells transfected with the FX construct encoding P239S mutant palladin outpaced the other cells at every time point. As shown in FIG. 18, on average, 33% more cells with the FX construct migrated through the transwell than the cells with WT construct and 40% more than the cells with the empty vector. Therefore, P239S palladin expression induces increased cell motility, consistent with a proposed oncogenic function.

[0471] While the present invention has been described with reference to the specific embodiments thereof, it should be understood by those skilled in the art that various changes may be made and equivalents may be substituted without departing from the true spirit and scope of the invention. In addition, many modifications may be made to adapt a particular situation, material, composition of matter, process, process step or steps, to the objective, spirit and scope of the present invention. All such modifications are intended to be within the scope of the claims appended hereto.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 116

<210> SEQ ID NO 1

<211> LENGTH: 4347

<212> TYPE: DNA

<213> ORGANISM: H. sapiens

<400> SEQUENCE: 1

```

cttgagtcac ccggcgggag aggtataaag cccgatacct gccccgcgcc eggtccgegg      60
agcccgcctg agtcccctgc cgtcccgac gcggaatcgg gcagcagcgg gaggcggccc      120
ggagagccga gggaccctct gaagctccag caactccaga accaaatccg actggagcag      180
gaggccggcg ctgcggcagc tccgccagcc ccgcgcagcg cgcgcgccctc gcccccttc      240
ccgcgcggcg ccgccttccc ctagctcgcg gcctgcacgc cgcgccgctc cccggagccc      300
atgagcgcgc tggcctcccg ctccgcccc gccatgcagt cctccggctc cttcaactac      360
gcgcgcccc agcagttcat cgcgcgcag aaactcgggc ccgcgtcggg ccacggcagc      420
ccggcctcca gccccagctc gtcagcctc cgtcgccca tgtccccgac gccgaggcag      480
ttcggccgcg ccccctgccc gcccttcgcg cagcccttcg gcgctgagcc cgaggccccg      540
tggggctcct cctgcgcctc gcccccgccc ccgccacccc cggctctcag ccccacggct      600
gccttcccgg tgcccagcgt gttcccactg ccgcgcgcc caccgccgct cccgagcccc      660
ggacaggcgt cccactgctc gtcgcctgcc acccgcttcg gccacggcca gacgcccgcg      720
gccttctca gcgctctgct gccctcgcag ccgccgcgg cggccgtcaa cgcctggggg      780
ctgcccagg gtgtcaccgc cgcagattt ccaaagaagg ccagtagaac tgctagaata      840
gcctccgatg aggaattca aggcacaaag gatgctgta ttaagacct ggaacgaaaa      900
cttcgcttca aggaggacct cctgaacaat gccagccga ggtaaacata cgaagaaaga      960
atggctcgtc gactgctagg tgctgacagt gcaactgtct ttaatatcca ggagccagaa     1020
gaggaacag ctaatcagga atacaaagtc tccagctgtg aacagagact catcagttaa     1080

```

-continued

atagagtaca	ggctagaaaag	gtctcctgtg	gatgaatcag	gtgatgaagt	tcagtatgga	1140
gatgtgcctg	tggaaaaatgg	aatggcacca	ttttttgaga	tgaagctgaa	acattacaag	1200
atctttgagg	gaatgccagt	aactttcaca	tgtagagtgg	ctggaaatcc	aaagccaaag	1260
atctattggt	ttaaagatgg	gaagcagatc	tctccaaaaga	gtgatcacta	caccattcaa	1320
agagatctcg	atgggacctg	ctccctccat	accacagcct	ccaccctaga	tgatgatggg	1380
aattatacaa	ttatggctgc	aaaccctcag	ggccgcatca	gttgtactgg	acggctaatag	1440
gtacaggctg	tcaaccaaag	aggtcgaagt	ccccggcttc	cctcaggcca	tcctcatgtc	1500
agaaggcctc	gttctagatc	aagggacagt	ggagacgaaa	atgaaccaat	tcaggagcga	1560
ttcttcagac	ctcacttctt	gcaggctcct	ggagatctga	ctgttcaaga	aggaaaactc	1620
tgcaaatgg	actgcaaagt	cagtggtgta	ccaacccag	atctaagctg	gcaactagat	1680
ggaaagcccc	tacgccctga	cagtgctcac	aagatgctgg	tgcgtgagaa	cggggtgcac	1740
tctctgatca	tagagccagt	cacgtcacgt	gatgccggca	tctacacatg	tatagctacc	1800
aaccgagcag	gacagaactc	attcagcctg	gagcttgtgg	ttgctgctaa	agaagcacac	1860
aaacccctg	tgtttattga	gaagctcaa	aacacaggag	ttgctgatgg	gtaccagtg	1920
cggctggaat	gtcgtgtatt	gggagtgcc	ccacctcaga	tattttgaa	gaaagaaaat	1980
gaatcactca	ctcacagcac	tgaccgagtg	agcatgcacc	aggacaacca	cggctacatc	2040
tgctctgctca	ttcagggagc	cacaaaagaa	gatgctgggt	ggtatactgt	gtcagccaag	2100
aatgaagcag	ggattgtgtc	ctgtactgcc	aggctggacg	tttacacca	gtggcatcag	2160
cagtacaga	gcaccaagcc	aaaaaaaaagta	cggccctcag	ccagtcgcta	tgcagcactt	2220
tccgaccagg	gactagacat	caaagcagcg	ttccaacctg	aggccaaccc	atctcacctg	2280
acactgaata	ctgccttggg	agaaagttag	gacctgtaat	ccagcattct	tgttaaagct	2340
gaaacactga	aacagccatt	gccttgacca	acatattcct	ttgtcacatt	atgtaaaagg	2400
cagaaacata	cctttgacta	taagaaatta	aaaaaaaaaca	ccaaaataat	atttttctta	2460
cttgatatac	caaacttagt	ttaagtagat	aatgctaata	caaatataca	cattgcacag	2520
aaaaatacaca	tttactgtcc	aatttaaaac	tttggaaatg	ctgtgattaa	agtgatcaaa	2580
atgcaaaaat	actaaaggaa	atcaattggt	cacaggtaac	tacaatttgt	attatctaca	2640
agtgccctta	aacacaagat	ataggtgctg	tgtagcctga	tagtgtgaaa	tgtttaatga	2700
gggagtgtga	ccacaacag	tactacaatg	attctgaagc	acagtgtatt	cagacagata	2760
cagtgaacca	agtgcaatat	gtaaggatga	aagaagaaga	gatgacaaag	aatccaagt	2820
aaatgccttg	tctttgcaaa	tgtttttata	ttaaatcata	aggaaggaac	tacttgccct	2880
aaatgttaat	atcaaaagag	ttttctaaca	aggtaatac	cttagttcct	aacatttttt	2940
ttctttatgt	gtagtgtttt	catgtacct	tggtaggaaa	cttatttaca	aaccatatta	3000
aaaggctaata	ttaaatataa	ataatataaa	gtgctctgaa	taaagcagaa	atatattaca	3060
gttcattcca	cagaaagcat	ccaaaccacc	caaatgacca	aggcatatat	agtatttgg	3120
ggaatcaggg	gtttggaagg	agtagggagg	agaatgaagg	aaaatgcaac	cagcatgatt	3180
atagtggttt	caattagata	aaagtagaag	gcacaggaga	ggtagcaaaag	gccaggcttt	3240
tctttggttt	tcttcaaaaca	taggtgaaaa	aaacactgcc	attcacaagt	caaggaaccc	3300
agggccagct	ggaagtgtgg	agcacacatg	ctgtggagca	cacatgctgt	ggagattgca	3360

-continued

```

gtgtgtctga ggtttgtgta gtagtggaag attttaggta tgtagagcaa gttgaaaatg 3420
gattgagact gcatgggtggc ataaatgaga aattgcctgt agcatctagt ctacttgaag 3480
gaagtggaga cataaggaga gacaaaaaca ggtttgtgcc ataaagtatt tttcaaaga 3540
caccaagatg tggtaaatga aaattattag ttcacttccc tgctgccatg aaactttgcc 3600
ttaagaaggt gctggattcc aaggtttgta aaggcatctc ggtaaagact gctttttgaa 3660
tgcatatgat tttgcatcag ctgactgag ttgattctga ccagactga tggttttaag 3720
tcggaaccga taaattttaa aaaggagaaa aaataatttg acctagtagt ataaaacatg 3780
aggctttaat ggtactttgc tatgaaaaga aaacactgta ttccttatgc aaaacacatg 3840
tatctttcat tatttataag tggcctctct tagctcagtt actcaattca tacgtagtat 3900
tttttaaaat aattttatat ctgtgtacca ccccatatat ttcattatc tgtttcacat 3960
gtacagcttt ctacttcttt gtaagaacac caaccaacca aggtttaagt gattaatagg 4020
cttgagcacc ggggtggcaga tgttctatgc agtgtggttc aagtttcttt gaccgcactt 4080
atatgcattg ctaatatgga atttaagata ccatacacag tctctcatgg acctatctct 4140
attgtagaat tatgacttat gtcttacttg gcaaattttt ctgaatgtga cctttttttg 4200
ctgatttgct gggtttggga ttaactagca ttattttgcc acctttatat tgtatttata 4260
aaaaaaaaagt actatcaatc aatcactacta ctttggattg ttgtgctggt gtaatgtgga 4320
tttaacatca ataaatattt gacaaat 4347

```

```

<210> SEQ ID NO 2
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: H. sapiens

```

```

<400> SEQUENCE: 2

```

```

gacgcccgcg 10

```

```

<210> SEQ ID NO 3
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: H. sapiens

```

```

<400> SEQUENCE: 3

```

```

gacgtccgcg 10

```

```

<210> SEQ ID NO 4
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: H. sapiens

```

```

<400> SEQUENCE: 4

```

```

gccacggcca gacgcccgcg gccttctca 30

```

```

<210> SEQ ID NO 5
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: H. sapiens

```

```

<400> SEQUENCE: 5

```

```

gccacggcca gacgtccgcg gccttctca 30

```

-continued

<210> SEQ ID NO 6
<211> LENGTH: 672
<212> TYPE: DNA
<213> ORGANISM: *H. sapiens*

<400> SEQUENCE: 6

```
ggaccctctg aagctccagc aactccagaa ccaaatccga ctggagcagg aggccggcgc      60
tcggcagcct ccgccagccc cgcgcagcgc gccgccctcg ccccccttcc cgcgcgcgcc      120
cgccttcccc gagctcgcgg cctgcacgcc gcccgcgctc ccggagccca tgagcgcgct      180
ggcctccccg tcgcgcccg ccatgcagtc ctccggctcc ttcaactacg cgcgccccaa      240
gcagttcate gccgcgcaga acctcggggc cgcgtcgggc cacggcacgc cggcctccag      300
ccccagctcg tcacgcctcc cgtcgcceat gtccccgacg ccgaggcagt tcggccgcgc      360
ccccgtgcgc ccttgcgcgc agcccttcgg cgctgagccc gaggccccgt gggggtcctc      420
ctcgcgctcg cccccgccc cgcaccccc ggtttcagc cccacggctg ccttcccggc      480
gcccgcagtg ttcccactgc cgcgcacc accgcccctc ccgagcccg gacaggcgtc      540
ccactgctcg tcgctgccca ccgcttcgg ccacggccag acgtccgcgg ccttctcag      600
cgctctgctg cctcgcagc cgcgcgcggc ggccgtcaac gccctggggc tgcccaaggg      660
tgtcaccccc gc                                          672
```

<210> SEQ ID NO 7
<211> LENGTH: 140
<212> TYPE: DNA
<213> ORGANISM: *H. sapiens*

<400> SEQUENCE: 7

```
caccgccgct cccgagcccg ggacaggcgt cccactgctc gtcgctgcc acccgcttcg      60
gccacggcca gacgtccgcg gccttctca ggcctctgct gccctcgcag ccgccgcgg      120
cggcgtcaa cgcctggggg                                          140
```

<210> SEQ ID NO 8
<211> LENGTH: 110
<212> TYPE: DNA
<213> ORGANISM: *H. sapiens*

<400> SEQUENCE: 8

```
gccttcccgg tgcccagcgt gttcccactg ccgcgcacc caccgccgct cccgagcccg      60
ggacaggcgt cccactgctc gtcgctgcc acccgcttcg gccacggcca                                          110
```

<210> SEQ ID NO 9
<211> LENGTH: 120
<212> TYPE: DNA
<213> ORGANISM: *H. sapiens*

<400> SEQUENCE: 9

```
tggggctcct cctcgcgcgc gccccgccc ccgccacccc cggttctcag ccccacggct      60
gccttcccgg tgcccagcgt gttcccactg ccgcgcacc caccgccgct cccgagcccg      120
```

<210> SEQ ID NO 10
<211> LENGTH: 120
<212> TYPE: DNA
<213> ORGANISM: *H. sapiens*

<400> SEQUENCE: 10

-continued

```

ggacaggcgt cccactgctc gtgcgctgcc acccgcttcg gccacggcca gacgtccgcg      60
gccttctcca gcgctctgct gccctgcgag ccgcccgcgg cggccgtcaa cgccttgggg      120

```

```

<210> SEQ ID NO 11
<211> LENGTH: 672
<212> TYPE: DNA
<213> ORGANISM: H. sapiens

```

```

<400> SEQUENCE: 11

```

```

ggaccctctg aagctccagc aactccagaa ccaaatccga ctggagcagg aggccggcgc      60
tcggcagcct ccgccagccc cgcgcagcgc gccgccctcg ccccccttc cgcgcgcgcc      120
cgcttcccc gagctcgcgg cctgcacgcc gcccgctcc cggagccca tgagcgcgct      180
ggctccccg tcgcccccg ccctgcagtc ctccggctcc tcaactacg cgcgccccaa      240
gcagttcctc gccgcgcaga acctcgggcc cgcgtcgggc cacggcacgc cggcctccag      300
ccccagctcg tcacgctcc cgtcgcctat gtccccgacg ccgaggcagt tcggccgcgc      360
ccccgtgccc ccttcgcgc agcccttcgg cgctgagccc gaggccccgt ggggctcctc      420
ctgcgcgctg cccccgcgcc cgcaccccc ggtcttcagc cccacggctg ccttccccgt      480
gcccgcagtg ttccactgc cgcgcacc accgcccctc ccgagcccg gacaggcgtc      540
ccactgctcg tcgctgccca ccgcttcgg ccacggccag acgcccgcgg ccttctcag      600
cgctctgctg cctcgcgagc cgcgcgcggc ggccgtcaac gccttggggc tgcccaaggg      660
tgtaaccccc gc                                          672

```

```

<210> SEQ ID NO 12
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: H. sapiens

```

```

<400> SEQUENCE: 12

```

```

atctaagctg gcaactagat                                          20

```

```

<210> SEQ ID NO 13
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: H. sapiens

```

```

<400> SEQUENCE: 13

```

```

atctaagctg tcaactagat                                          20

```

```

<210> SEQ ID NO 14
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: H. sapiens

```

```

<400> SEQUENCE: 14

```

```

ccaaccccag atctaagctg gcaactagat ggaaagcccg                    40

```

```

<210> SEQ ID NO 15
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: H. sapiens

```

```

<400> SEQUENCE: 15

```

```

ccaaccccag atctaagctg tcaactagat ggaaagcccg                    40

```

-continued

<210> SEQ ID NO 16
<211> LENGTH: 208
<212> TYPE: DNA
<213> ORGANISM: *H. sapiens*

<400> SEQUENCE: 16

gtcagtgggt taccaacccc agatctaagc tggcaactag atggaaagcc cgtacgccct 60
gacagtgctc acaagatgct ggtgctgag aacgggggtgc actctctgat catagagcca 120
gtcacgtcac gtgatgccg catctacaca tgtatagcta ccaaccgagc aggacagaac 180
tcattcagcc tggagcttgt ggttgctg 208

<210> SEQ ID NO 17
<211> LENGTH: 208
<212> TYPE: DNA
<213> ORGANISM: *H. sapiens*

<400> SEQUENCE: 17

gtcagtgggt taccaacccc agatctaagc tgtcaactag atggaaagcc cgtacgccct 60
gacagtgctc acaagatgct ggtgctgag aacgggggtgc actctctgat catagagcca 120
gtcacgtcac gtgatgccg catctacaca tgtatagcta ccaaccgagc aggacagaac 180
tcattcagcc tggagcttgt ggttgctg 208

<210> SEQ ID NO 18
<211> LENGTH: 122
<212> TYPE: DNA
<213> ORGANISM: *H. sapiens*

<400> SEQUENCE: 18

gtcagtgggt taccaacccc agatctaagc tggcaactag atggaaagcc cgtacgccct 60
gacagtgctc acaagatgct ggtgctgag aacgggggtgc actctctgat catagagcca 120
gt 122

<210> SEQ ID NO 19
<211> LENGTH: 122
<212> TYPE: DNA
<213> ORGANISM: *H. sapiens*

<400> SEQUENCE: 19

gtcagtgggt taccaacccc agatctaagc tgtcaactag atggaaagcc cgtacgccct 60
gacagtgctc acaagatgct ggtgctgag aacgggggtgc actctctgat catagagcca 120
gt 122

<210> SEQ ID NO 20
<211> LENGTH: 5809
<212> TYPE: DNA
<213> ORGANISM: *H. sapiens*

<400> SEQUENCE: 20

gtgaccacgg accaggcagt ctctaataa taggcaaggc cacaacctcc attctcccag 60
aaaagaaga atgctcatc gaaattcacc acctctctgg agtcttcaaa ctgaccaagc 120
attgaaaaga acacagtttc agaaaacagt ttccagtgcc tctggccttc ctactgaaag 180
cagacacaga gtgcatgaag accgttcaaa tatgtcaggg acctcctccc atgagtcctt 240

-continued

ctatgactcc	ctctcagaca	tgcaggaaga	aagcaagaat	actgacttct	tcccgggcct	300
ttctgcttct	ctcagccagg	aagagataaa	caagagtctt	gacctggccc	ggagagccat	360
agccgactcc	gaaacagaag	atthtgactc	ggaaaaggag	atctcgcaga	ttttcagtac	420
ttctcctgca	agcctctgtg	aacatccttc	ccataaggag	accaaattgg	gtgaacacgc	480
ctcaggagga	cctcaggata	acaggtcaac	acctgtccag	cctctggcag	agaacaaaac	540
taagagtatc	tcttcacctg	tttcaaagag	gaaacctgcc	atgtcacccc	tgctcaccag	600
gcccagctac	atccggagcc	tccgaaagcc	tgaaaagcgt	ggtgcaaaaa	ctcccagcac	660
aaacgtaaa	cccaaaacgc	cacatcaaag	aaaggggtggc	cccagagcc	agctgtgtga	720
caaggcagct	aatttaattg	aggagctaac	atccatattt	aaagccgcaa	agccaagaaa	780
cagaagccca	aatggggagt	cctcgtcacc	agacagtggg	tacctgtctc	ctaaaaatca	840
gccgtcagcc	ctgctgagtg	cctcagccag	ccagagccct	atggaagacc	aaggggagat	900
ggaaagagag	gtcaagtccc	ctggggccag	gcattgtctac	caggacaacc	aggacttggc	960
agtgccacac	aaccgcaagt	ctcaccaca	gccccacagc	gcctccact	tcccagctgc	1020
acctcgatc	atccaaaagc	tgaggagcca	agaagtagca	gaagggagcc	gagtttatct	1080
ggagtgtaga	gtcaactgaa	acccactcc	tcgagtcaga	tggttctgtg	aagggaaaga	1140
actgcacaac	actcctgata	ttcaaatcca	ctgtgagggg	ggggacctcc	ataccctgat	1200
catagcagag	gcctttgagg	acgacacagg	tcgctacacc	tgtttggcta	cgaatcccag	1260
cggtcagac	acaacatctg	ctgaggtgtt	cattgaaggt	gccagttcaa	cagattctga	1320
cagtgaaggt	ttagctttca	aatcaagagc	tggagctatg	ccacaagctc	aaaagaaaac	1380
aacttctgtt	tccttgacaa	taggatcctc	atctccaaag	acaggggtga	ccacagctgt	1440
gattcaacca	ctgtctgtcc	ctgtgcaaca	ggttcacagt	ccaacttcat	atctctgccg	1500
acctgatgga	accactactg	cctactttcc	tcctgttttt	acaaggaac	tgcaaaacac	1560
agccgtggcg	gaaggccagg	tgggtgttct	ggagtgccgg	gtccgtgggg	caccccctct	1620
gcaggtccag	tggtttcgce	aagggagtga	aatccaagac	tctccagatt	tccgaattct	1680
acagaaaaaa	cctagatcta	cagctgaacc	tgaggagatt	tgcaccctag	ttatcgtctga	1740
gactttccct	gaagatgcag	ggatctttac	atgttcagca	agaaatgatt	atggatcagc	1800
aaccagcact	gcccagctgg	ttgtcacctc	agccaacact	gaaaactgta	gttacgagtc	1860
aatgggagaa	tccaacaatg	accactttca	acactttcca	cctccccctc	caatcttggg	1920
gacaagttcc	ttggagttag	cttcaaagaa	accatctgag	atccagcagg	tgaacaaccc	1980
tgagttaggc	ctgagcaggg	cagcccttca	aatgcaattc	aatgctgctg	agagggaaac	2040
gaacggagtc	catcccagcc	gtggagtaaa	tggactgatt	aacggcaaag	ctaacagtaa	2100
taaatctctt	ccaacaccag	ctgtcctgct	ttcaccact	aaggagccac	cacctctgct	2160
tgccaaacca	aaactaggat	ttocaaagaa	ggccagtaga	actgctagaa	tagcctccga	2220
tgaggaaatt	caaggcacia	aggatgctgt	tattcaagac	ctggaacgaa	aacttcgctt	2280
caaggaggac	ctcctgaaca	atggccagcc	gaggttaaca	tacgaagaaa	gaatggctcg	2340
tcgactgcta	gggtctgaca	gtgcaactgt	ctttaatatt	caggagccag	aagaggaaac	2400
agctaatcag	gaatacaaa	tctccagctg	tgaacagaga	ctcatcagtg	aaatagagta	2460
caggctagaa	aggtctcctg	tggatgaatc	aggtgatgaa	gttcagtatg	gagatgtgcc	2520

-continued

tgtggaaaat ggaatggcac cattctttga gatgaagctg aaacattaca agatctttga 2580
gggaatgccca gtaactttca catgtagagt ggctggaaaat ccaaagccaa agatctattg 2640
gtttaaagat ggaagcaga tctctccaaa gagtgatcac tacaccattc aaagagatct 2700
cgatgggacc tgctccctcc ataccacagc ctccacccta gatgatgatg ggaattatac 2760
aattatggct gcaaaccctc agggccgcat cagttgtact ggacggctaa tggtagagge 2820
tgtcaaccaa agaggtcgaa gtccccggtc tccctcagge catcctcatg tcagaaggcc 2880
tcgttctaga tcaagggaca gtggagacga aaatgaacca attcaggagc gattcttcag 2940
acctcacttc ttgcaggctc ctggagatct gactgttcaa gaaggaaaac tctgcagaat 3000
ggactgcaaa gtcagtggtt taccaacccc agatctaagc tggcaactag atggaaagcc 3060
cgtacgccct gacagtgtc acaagatgct ggtgcgtgag aacggggtgc actctctgat 3120
catagagcca gtcacgtcac gtgatgccg catctacaca tgtatagcta ccaaccgagc 3180
aggacagaac tcattcagcc tggagcttgt ggttgctgct aaagaagcac acaaaccccc 3240
tgtgtttatt gagaagctcc aaaacacagc agttgctgat gggtaaccag tgcggctgga 3300
atgtcgtgta ttgggagtgc caccacctca gatattttgg aagaagaaa atgaatcact 3360
cactcacagc actgaccgag tgagcatgca ccaggacaac cacggctaca tctgcctgct 3420
cattcagga gccacaaaag aagatgctgg gtggtatact gtgtcagcca agaatgaagc 3480
agggattgtg tctgtactg ccaggctgga cgtttacatt tctcgacatt aatagtgaac 3540
cacaccagga gaacaaatac ccaaccctcaggatcagca gtcacagagc accaagccaa 3600
aaaaagtacg gccctcagcc agtcgctatg cagcactttc ggaccagga ctagacatca 3660
aagcagcgtt ccaacctgag gccaacccat ctcacctgac actgaatact gccttggtag 3720
aaagtgagga cctgtaatcc agcattcttg ttaaagctga aacctgaaa cagccattgc 3780
cttgaccaac atattccttt gtcacattat gtaaaaggca gaaacatacc tttgactata 3840
agaaattaaa aaaaaaac caaataata tttttcttac ttgatatacc aaacttagtt 3900
taagtagata atgctaatac aaatatacac attgcacaga aaatacacat ttactgtcca 3960
atttaaaact ttggaattgc tgtgattaaa gtgatcaaaa tgccaaaata ctaaaggaaa 4020
tcaattgttc acaggttaact acaatttgta ttatctacaa gtgcctttaa acacaagata 4080
taggtgctgt gtagcctgat agtgtgaaat gtttaatgag ggagttgtac cacaaacagt 4140
actacaatga ttctgaaagca cagtgtattc agacagatagc agtgaaccaa gtgcaatag 4200
taaggatgaa agaagaagag atgacaaaga aatccaagta aatgccttgt ctttgcaaat 4260
gtttttatat taaatcataa ggaaggaact acttgcctta aatgttaata tcaaaagagt 4320
tttotaacaa ggttaatacc ttagtctta acatttttt tctttatgtg tagtgtttc 4380
atgtacctt ggtaggaaac ttatttaca accatattaa aaggctaatt taaatataaa 4440
taatataaag tgctctgaat aaagcagaaa tatattacag ttattccac agaagcatc 4500
caaacacccc aatgaccaa ggcatatata gtatttgag gaatcagggg tttggaagga 4560
gtagggagga gaatgaagga aatgcaacc agcatgatta tagtgtgttc atttagataa 4620
aagtagaagg cacagagag gtagcaagg ccaggctttt ctttggtttt cttcaaacat 4680
aggtgaaaa aacctgccca ttcacaagtc aaggaacca gggccagctg gaagtgtgga 4740
gcacacatgc tgtggagcac acatgctgtg gagattgcag tgtgtctgag gtttgtgtag 4800

-continued

```

tagtgaaga ttttaggtat gtagagcaag ttgaaaatgg attgagactg catgggtggca 4860
taaatgagaa attgctgtga gcatctagtc tacttgaagg aagtggagac ataaggagag 4920
acaaaaacag gtttgtgcc aaaagtattt tttcaaagac accaagatgt ggtaaatgaa 4980
aattattagt tcaactccct gctgcatga aactttgect taagaagggtg ctggattcca 5040
aggtttgtaa aggcactctcg gtaaagactg ctttttgaat gcatatgatt ttgcatcagc 5100
tagactgagt tgattctgac cagacttgat ggttttaagt cggaaaccgat aaattttaaa 5160
aaggagaaaa aataatttga cctagtagta taaaacatga ggctttaatg gtactttgct 5220
atgaaaagaa aacactgtat tccttatgca aaacacatgt atctttcatt atttataagt 5280
ggcctctctt agctcagtta ctcaattcat acgtagtatt ttttaaata attttatctc 5340
tgtgtaccac cccatatatt tcatattact gtttcacatg tacagctttc tacttctttg 5400
taagaacacc aaccaaccaa ggtttaagtg attaataggc ttgagcaccg ggtggcagat 5460
gttctatgca gtgtggttca agtttctttg accgcactta tatgcattgc taatatggaa 5520
tttaagatac catacacagt ctctcatgga cctatctcta ttgtagaatt atgacttatg 5580
tcttacttgc caaatttttc tgaatgtgac ctttttttgc tgatttgctg ggtttgggat 5640
taactagcat tattttgcc actttatatt gtatttataa aaaaaaagt actatcaatc 5700
aatcactacta ctttgattg ttgtgctggt gtaatgtgga tttaacatca ataaatattt 5760
gacaaataat agttgcagtt ttgtgaagca aaataaatat tcagtttta 5809

```

```

<210> SEQ ID NO 21
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: H. sapiens

```

```

<400> SEQUENCE: 21

```

```

cccagcctg gagtaaat 18

```

```

<210> SEQ ID NO 22
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: H. sapiens

```

```

<400> SEQUENCE: 22

```

```

cccagctgtg gagtaaat 18

```

```

<210> SEQ ID NO 23
<211> LENGTH: 41
<212> TYPE: DNA
<213> ORGANISM: H. sapiens

```

```

<400> SEQUENCE: 23

```

```

ggagtccatc ccagcctgg agtaaatgga ctgattaacg g 41

```

```

<210> SEQ ID NO 24
<211> LENGTH: 41
<212> TYPE: DNA
<213> ORGANISM: H. sapiens

```

```

<400> SEQUENCE: 24

```

```

ggagtccatc ccagctgtgg agtaaatgga ctgattaacg g 41

```

-continued

```

<210> SEQ ID NO 25
<211> LENGTH: 343
<212> TYPE: DNA
<213> ORGANISM: H. sapiens

<400> SEQUENCE: 25

ccaacactga aaactgtagt tacgagtcaa tgggagaatc caacaatgac cacttccaac    60
actttccacc tccccctcca atcttgagaga caagttcctt ggagttggct tcaaagaaac    120
catctgagat ccagcagggtg aacaaccctg agttaggcct gagcagggca gcccttcaaa    180
tgcaattcaa tgctgctgag agggaaacga acggagtcca tcccagccgt ggagtaaatg    240
gactgattaa cggcaaagct aacagtaata aatctcttcc aacaccagct gtctctgctt    300
caccactaa ggagccacca cctctgcttg ccaaaccaaa act                          343

```

```

<210> SEQ ID NO 26
<211> LENGTH: 343
<212> TYPE: DNA
<213> ORGANISM: H. sapiens

<400> SEQUENCE: 26

ccaacactga aaactgtagt tacgagtcaa tgggagaatc caacaatgac cacttccaac    60
actttccacc tccccctcca atcttgagaga caagttcctt ggagttggct tcaaagaaac    120
catctgagat ccagcagggtg aacaaccctg agttaggcct gagcagggca gcccttcaaa    180
tgcaattcaa tgctgctgag agggaaacga acggagtcca tcccagctgt ggagtaaatg    240
gactgattaa cggcaaagct aacagtaata aatctcttcc aacaccagct gtctctgctt    300
caccactaa ggagccacca cctctgcttg ccaaaccaaa act                          343

```

```

<210> SEQ ID NO 27
<211> LENGTH: 93
<212> TYPE: DNA
<213> ORGANISM: H. sapiens

<400> SEQUENCE: 27

ttcaatgctg ctgagagggga aacgaacgga gtccatccca gccgtggagt aaatggactg    60
attaacggca aagctaacag taataaatct ctt                                  93

```

```

<210> SEQ ID NO 28
<211> LENGTH: 93
<212> TYPE: DNA
<213> ORGANISM: H. sapiens

<400> SEQUENCE: 28

ttcaatgctg ctgagagggga aacgaacgga gtccatccca gctgtggagt aaatggactg    60
attaacggca aagctaacag taataaatct ctt                                  93

```

```

<210> SEQ ID NO 29
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 29

cggccagacg tccgcggcct t                                          21

```

-continued

<210> SEQ ID NO 30
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 30

aaggccgagg acgtctggcc g 21

<210> SEQ ID NO 31
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 31

gccacggcca gacgtccgag gcctt 25

<210> SEQ ID NO 32
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 32

aaggccgagg acgtctggcc gtggc 25

<210> SEQ ID NO 33
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 33

cggccagagc tccgaggcct tcctc 25

<210> SEQ ID NO 34
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 34

gaggaaggcc gaggacgtct ggccgtggc 29

<210> SEQ ID NO 35
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 35

gccacggcca gacgtccgag gccttctc 29

<210> SEQ ID NO 36
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 36

gaggaaggcc gcggacgtct ggccgtggc 29

<210> SEQ ID NO 37
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 37

ctaagctggc aactagatgg 20

<210> SEQ ID NO 38
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 38

ccatctagtt gccagcttag 20

<210> SEQ ID NO 39
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 39

gatctaagct ggcaactaga t 21

<210> SEQ ID NO 40
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 40

atctagttgc cagcttagat c 21

<210> SEQ ID NO 41
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 41

gatctaagct ggcaactaga tgg 23

<210> SEQ ID NO 42
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 42

-continued

ccatctagtt gccagcttag atc 23

<210> SEQ ID NO 43
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 43

ccatcccagc cgtggagtaa atgg 24

<210> SEQ ID NO 44
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 44

ccatttactc cacggctggg atgg 24

<210> SEQ ID NO 45
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 45

cccagccgtg gagtaaatgg 20

<210> SEQ ID NO 46
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 46

ccatttactc cacggctggg 20

<210> SEQ ID NO 47
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 47

gtccatccca gccgtggagt aatggactg 30

<210> SEQ ID NO 48
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 48

cagtccattt actccacggc tgggatggac 30

-continued

<210> SEQ ID NO 49
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 49

accgcgttcg gccacggcca 20

<210> SEQ ID NO 50
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 50

gtcgcctgcc accgcgttcg gccacggc 28

<210> SEQ ID NO 51
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 51

gccttctca gcgctctgct 20

<210> SEQ ID NO 52
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 52

gccttctca gcgctctgct gcctcgcag 30

<210> SEQ ID NO 53
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 53

gcgctctgct gcctcgcag ccgcc 25

<210> SEQ ID NO 54
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 54

gcctcgcag ccgccccgg cggccgtcaa 30

<210> SEQ ID NO 55
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 55

cagtgggta ccaacccag 20

<210> SEQ ID NO 56
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 56

ccaacccag atctaagctg 20

<210> SEQ ID NO 57
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 57

ggaaagcccg tacgcctga 20

<210> SEQ ID NO 58
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 58

cagtgctcac aagatgctgg 20

<210> SEQ ID NO 59
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 59

aatgctgctg agagggaac 20

<210> SEQ ID NO 60
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 60

agagggaac gaacggagtc catcc 25

<210> SEQ ID NO 61
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 61

-continued

gtggagtaaa tggactgatt aacggcaaag 30

<210> SEQ ID NO 62
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 62

gtaaattggac tgattaacgg c 21

<210> SEQ ID NO 63
<211> LENGTH: 120
<212> TYPE: DNA
<213> ORGANISM: H. sapiens

<400> SEQUENCE: 63

ggacaggcgt cccactgctc gtcgctgcc acccgcttcg gccacggcca gacgtccgcg 60

gccttctca gcgctctgct gccctcgcag ccgcccgcgg cggccgtcaa cgccttgggg 120

<210> SEQ ID NO 64
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 64

ggacaggcgt cactgctc 19

<210> SEQ ID NO 65
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 65

ccccagggcg ttgacggcgg 20

<210> SEQ ID NO 66
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 66

ggaccctctg aagctccagc 20

<210> SEQ ID NO 67
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 67

gcgggggtga caccctggg 20

-continued

<210> SEQ ID NO 68
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 68

ggaccctctg aagctccagc 20

<210> SEQ ID NO 69
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 69

ccgtggccga agcgggtggc agg 23

<210> SEQ ID NO 70
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 70

cctcgccgtc gccccgcc 20

<210> SEQ ID NO 71
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 71

ccgtggccga agcgggtggc agg 23

<210> SEQ ID NO 72
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 72

cctcgccgtc gccccgcc 20

<210> SEQ ID NO 73
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 73

ccgtggccga agcgggtggc agg 23

<210> SEQ ID NO 74
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 74

aggtgtcact tctctttttc ccccc 25

<210> SEQ ID NO 75
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 75

ggggagggaa gtggaggacc gcgg 24

<210> SEQ ID NO 76
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 76

cagtgggta ccaacccag 20

<210> SEQ ID NO 77
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 77

gtgcaccccg ttctcagca 20

<210> SEQ ID NO 78
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 78

tcttgtaact ctgaaggagg aatttatgc 29

<210> SEQ ID NO 79
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 79

ttttccgtat tggttagtaa tgtagaatta g 31

<210> SEQ ID NO 80
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 80

-continued

gggaaacgaa cggagtccat ccc 23

<210> SEQ ID NO 81
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 81

agcaggacag ctggtgttgg 20

<210> SEQ ID NO 82
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 82

cacaacacag ggattctcag aaga 24

<210> SEQ ID NO 83
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 83

agctggtgtt ggaagagatt 20

<210> SEQ ID NO 84
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 84

ccgacgtgtt cccactgc 18

<210> SEQ ID NO 85
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 85

cgcacggaga gaaatgtgtg 20

<210> SEQ ID NO 86
<211> LENGTH: 208
<212> TYPE: DNA
<213> ORGANISM: H. sapiens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: 33
<223> OTHER INFORMATION: n = A,T,C or G

<400> SEQUENCE: 86

gtcagtgggt taccaacccc agatctaagc tgncaactag atggaaagcc cgtacgcct 60

-continued

```

gacagtgctc acaagatgct ggtgcgtgag aacgggggtgc actctctgat catagagcca 120
gtcacgtcac gtgatgccgg catctacaca tgtatagcta ccaaccgagc aggacagaac 180
tcattcagcc tggagcttgt ggttgetg 208

```

```

<210> SEQ ID NO 87
<211> LENGTH: 1106
<212> TYPE: PRT
<213> ORGANISM: H. sapiens

```

```

<400> SEQUENCE: 87

```

```

Met Ser Gly Thr Ser Ser His Glu Ser Phe Tyr Asp Ser Leu Ser Asp
1           5           10           15
Met Gln Glu Glu Ser Lys Asn Thr Asp Phe Phe Pro Gly Leu Ser Ala
20          25          30
Phe Leu Ser Gln Glu Glu Ile Asn Lys Ser Leu Asp Leu Ala Arg Arg
35          40          45
Ala Ile Ala Asp Ser Glu Thr Glu Asp Phe Asp Ser Glu Lys Glu Ile
50          55          60
Ser Gln Ile Phe Ser Thr Ser Pro Ala Ser Leu Cys Glu His Pro Ser
65          70          75          80
His Lys Glu Thr Lys Leu Gly Glu His Ala Ser Arg Arg Pro Gln Asp
85          90          95
Asn Arg Ser Thr Pro Val Gln Pro Leu Ala Glu Lys Gln Thr Lys Ser
100         105        110
Ile Ser Ser Pro Val Ser Lys Arg Lys Pro Ala Met Ser Pro Leu Leu
115        120        125
Thr Arg Pro Ser Tyr Ile Arg Ser Leu Arg Lys Ala Glu Lys Arg Gly
130        135        140
Ala Lys Thr Pro Ser Thr Asn Val Lys Pro Lys Thr Pro His Gln Arg
145        150        155        160
Lys Gly Gly Pro Gln Ser Gln Leu Cys Asp Lys Ala Ala Asn Leu Ile
165        170        175
Glu Glu Leu Thr Ser Ile Phe Lys Ala Ala Lys Pro Arg Asn Arg Ser
180        185        190
Pro Asn Gly Glu Ser Ser Ser Pro Asp Ser Gly Tyr Leu Ser Pro Lys
195        200        205
Asn Gln Pro Ser Ala Leu Leu Ser Ala Ser Ala Ser Gln Ser Pro Met
210        215        220
Glu Asp Gln Gly Glu Met Glu Arg Glu Val Lys Ser Pro Gly Ala Arg
225        230        235        240
His Cys Tyr Gln Asp Asn Gln Asp Leu Ala Val Pro His Asn Arg Lys
245        250        255
Ser His Pro Gln Pro His Ser Ala Leu His Phe Pro Ala Ala Pro Arg
260        265        270
Phe Ile Gln Lys Leu Arg Ser Gln Glu Val Ala Glu Gly Ser Arg Val
275        280        285
Tyr Leu Glu Cys Arg Val Thr Gly Asn Pro Thr Pro Arg Val Arg Trp
290        295        300
Phe Cys Glu Gly Lys Glu Leu His Asn Thr Pro Asp Ile Gln Ile His
305        310        315        320
Cys Glu Gly Gly Asp Leu His Thr Leu Ile Ile Ala Glu Ala Phe Glu

```

-continued

325					330										335
Asp	Asp	Thr	Gly	Arg	Tyr	Thr	Cys	Leu	Ala	Thr	Asn	Pro	Ser	Gly	Ser
340					345					350					
Asp	Thr	Thr	Ser	Ala	Glu	Val	Phe	Ile	Glu	Gly	Ala	Ser	Ser	Thr	Asp
355					360					365					
Ser	Asp	Ser	Glu	Ser	Leu	Ala	Phe	Lys	Ser	Arg	Ala	Gly	Ala	Met	Pro
370					375					380					
Gln	Ala	Gln	Lys	Lys	Thr	Thr	Ser	Val	Ser	Leu	Thr	Ile	Gly	Ser	Ser
385					390					395					400
Ser	Pro	Lys	Thr	Gly	Val	Thr	Thr	Ala	Val	Ile	Gln	Pro	Leu	Ser	Val
405					410					415					
Pro	Val	Gln	Gln	Val	His	Ser	Pro	Thr	Ser	Tyr	Leu	Cys	Arg	Pro	Asp
420					425					430					
Gly	Thr	Thr	Thr	Ala	Tyr	Phe	Pro	Pro	Val	Phe	Thr	Lys	Glu	Leu	Gln
435					440					445					
Asn	Thr	Ala	Val	Ala	Glu	Gly	Gln	Val	Val	Val	Leu	Glu	Cys	Arg	Val
450					455					460					
Arg	Gly	Ala	Pro	Pro	Leu	Gln	Val	Gln	Trp	Phe	Arg	Gln	Gly	Ser	Glu
465					470					475					480
Ile	Gln	Asp	Ser	Pro	Asp	Phe	Arg	Ile	Leu	Gln	Lys	Lys	Pro	Arg	Ser
485					490					495					
Thr	Ala	Glu	Pro	Glu	Glu	Ile	Cys	Thr	Leu	Val	Ile	Ala	Glu	Thr	Phe
500					505					510					
Pro	Glu	Asp	Ala	Gly	Ile	Phe	Thr	Cys	Ser	Ala	Arg	Asn	Asp	Tyr	Gly
515					520					525					
Ser	Ala	Thr	Ser	Thr	Ala	Gln	Leu	Val	Val	Thr	Ser	Ala	Asn	Thr	Glu
530					535					540					
Asn	Cys	Ser	Tyr	Glu	Ser	Met	Gly	Glu	Ser	Asn	Asn	Asp	His	Phe	Gln
545					550					555					560
His	Phe	Pro	Pro	Pro	Pro	Pro	Ile	Leu	Glu	Thr	Ser	Ser	Leu	Glu	Leu
565					570					575					
Ala	Ser	Lys	Lys	Pro	Ser	Glu	Ile	Gln	Gln	Val	Asn	Asn	Pro	Glu	Leu
580					585					590					
Gly	Leu	Ser	Arg	Ala	Ala	Leu	Gln	Met	Gln	Phe	Asn	Ala	Ala	Glu	Arg
595					600					605					
Glu	Thr	Asn	Gly	Val	His	Pro	Ser	Arg	Gly	Val	Asn	Gly	Leu	Ile	Asn
610					615					620					
Gly	Lys	Ala	Asn	Ser	Asn	Lys	Ser	Leu	Pro	Thr	Pro	Ala	Val	Leu	Leu
625					630					635					640
Ser	Pro	Thr	Lys	Glu	Pro	Pro	Pro	Leu	Leu	Ala	Lys	Pro	Lys	Leu	Gly
645					650					655					
Phe	Pro	Lys	Lys	Ala	Ser	Arg	Thr	Ala	Arg	Ile	Ala	Ser	Asp	Glu	Glu
660					665					670					
Ile	Gln	Gly	Thr	Lys	Asp	Ala	Val	Ile	Gln	Asp	Leu	Glu	Arg	Lys	Leu
675					680					685					
Arg	Phe	Lys	Glu	Asp	Leu	Leu	Asn	Asn	Gly	Gln	Pro	Arg	Leu	Thr	Tyr
690					695					700					
Glu	Glu	Arg	Met	Ala	Arg	Arg	Leu	Leu	Gly	Ala	Asp	Ser	Ala	Thr	Val
705					710					715					720
Phe	Asn	Ile	Gln	Glu	Pro	Glu	Glu	Glu	Thr	Ala	Asn	Gln	Glu	Tyr	Lys
725					730					735					

-continued

Val Ser Ser Cys Glu Gln Arg Leu Ile Ser Glu Ile Glu Tyr Arg Leu
 740 745 750
 Glu Arg Ser Pro Val Asp Glu Ser Gly Asp Glu Val Gln Tyr Gly Asp
 755 760 765
 Val Pro Val Glu Asn Gly Met Ala Pro Phe Phe Glu Met Lys Leu Lys
 770 775 780
 His Tyr Lys Ile Phe Glu Gly Met Pro Val Thr Phe Thr Cys Arg Val
 785 790 795 800
 Ala Gly Asn Pro Lys Pro Lys Ile Tyr Trp Phe Lys Asp Gly Lys Gln
 805 810 815
 Ile Ser Pro Lys Ser Asp His Tyr Thr Ile Gln Arg Asp Leu Asp Gly
 820 825 830
 Thr Cys Ser Leu His Thr Thr Ala Ser Thr Leu Asp Asp Asp Gly Asn
 835 840 845
 Tyr Thr Ile Met Ala Ala Asn Pro Gln Gly Arg Ile Ser Cys Thr Gly
 850 855 860
 Arg Leu Met Val Gln Ala Val Asn Gln Arg Gly Arg Ser Pro Arg Ser
 865 870 875 880
 Pro Ser Gly His Pro His Val Arg Arg Pro Arg Ser Arg Ser Arg Asp
 885 890 895
 Ser Gly Asp Glu Asn Glu Pro Ile Gln Glu Arg Phe Phe Arg Pro His
 900 905 910
 Phe Leu Gln Ala Pro Gly Asp Leu Thr Val Gln Glu Gly Lys Leu Cys
 915 920 925
 Arg Met Asp Cys Lys Val Ser Gly Leu Pro Thr Pro Asp Leu Ser Trp
 930 935 940
 Gln Leu Asp Gly Lys Pro Val Arg Pro Asp Ser Ala His Lys Met Leu
 945 950 955 960
 Val Arg Glu Asn Gly Val His Ser Leu Ile Ile Glu Pro Val Thr Ser
 965 970 975
 Arg Asp Ala Gly Ile Tyr Thr Cys Ile Ala Thr Asn Arg Ala Gly Gln
 980 985 990
 Asn Ser Phe Ser Leu Glu Leu Val Val Ala Ala Lys Glu Ala His Lys
 995 1000 1005
 Pro Pro Val Phe Ile Glu Lys Leu Gln Asn Thr Gly Val Ala Asp Gly
 1010 1015 1020
 Tyr Pro Val Arg Leu Glu Cys Arg Val Leu Gly Val Pro Pro Pro Gln
 1025 1030 1035 1040
 Ile Phe Trp Lys Lys Glu Asn Glu Ser Leu Thr His Ser Thr Asp Arg
 1045 1050 1055
 Val Ser Met His Gln Asp Asn His Gly Tyr Ile Cys Leu Leu Ile Gln
 1060 1065 1070
 Gly Ala Thr Lys Glu Asp Ala Gly Trp Tyr Thr Val Ser Ala Lys Asn
 1075 1080 1085
 Glu Ala Gly Ile Val Ser Cys Thr Ala Arg Leu Asp Val Tyr Ile Ser
 1090 1095 1100
 Arg His
 1105

<210> SEQ ID NO 88

<211> LENGTH: 40

-continued

```

<212> TYPE: PRT
<213> ORGANISM: H. sapiens
<220> FEATURE:
<221> NAME/KEY: VARIANT
<222> LOCATION: 11
<223> OTHER INFORMATION: Xaa = Any Amino Acid

```

```

<400> SEQUENCE: 88

```

```

Val Ser Gly Leu Pro Thr Pro Asp Leu Ser Xaa Gln Leu Asp Gly Lys
1           5           10           15

```

```

Pro Val Arg Pro Asp Ser Ala His Lys Met Leu Val Arg Glu Asn Gly
20          25          30

```

```

Val His Ser Leu Ile Ile Glu Pro
35          40

```

```

<210> SEQ ID NO 89
<211> LENGTH: 40
<212> TYPE: PRT
<213> ORGANISM: H. sapiens

```

```

<400> SEQUENCE: 89

```

```

Val Ser Gly Leu Pro Thr Pro Asp Leu Ser Trp Gln Leu Asp Gly Lys
1           5           10           15

```

```

Pro Val Arg Pro Asp Ser Ala His Lys Met Leu Val Arg Glu Asn Gly
20          25          30

```

```

Val His Ser Leu Ile Ile Glu Pro
35          40

```

```

<210> SEQ ID NO 90
<211> LENGTH: 40
<212> TYPE: PRT
<213> ORGANISM: mouse

```

```

<400> SEQUENCE: 90

```

```

Val Ser Gly Leu Pro Thr Pro Asp Leu Ser Trp Gln Leu Asp Gly Lys
1           5           10           15

```

```

Pro Ile Arg Pro Asp Ser Ala His Lys Met Leu Val Arg Glu Asn Gly
20          25          30

```

```

Val His Ser Leu Ile Ile Glu Pro
35          40

```

```

<210> SEQ ID NO 91
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

```

```

<400> SEQUENCE: 91

```

```

tcttgta cta ctgaaggagg aatttatgc

```

29

```

<210> SEQ ID NO 92
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

```

```

<400> SEQUENCE: 92

```

```

ttttccgtat tggtagtaa tgtagaatta g

```

31

-continued

<210> SEQ ID NO 93
 <211> LENGTH: 343
 <212> TYPE: DNA
 <213> ORGANISM: H. sapiens
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: 228
 <223> OTHER INFORMATION: n = A,T,C or G

<400> SEQUENCE: 93

```

ccaacactga aaactgtagt tacgagtcaa tgggagaatc caacaatgac cacttccaac    60
actttccacc tccccctcca atcttgagaga caagttcctt ggagttggct tcaaagaaac   120
catctgagat ccagcaggtg aacaaccctg agttaggctt gacgagggca gcccttcaaa   180
tgcaattcaa tgctgctgag agggaaacga acggagtcca tcccagcngt ggagtaaatag   240
gactgattaa cggcaaagct aacagtaata aatctcttcc aacaccagct gtctgcttt    300
caccactaa  ggagccacca cctctgcttg ccaaaccaaa act                          343

```

<210> SEQ ID NO 94
 <211> LENGTH: 113
 <212> TYPE: PRT
 <213> ORGANISM: H. sapiens
 <220> FEATURE:
 <221> NAME/KEY: VARIANT
 <222> LOCATION: 76
 <223> OTHER INFORMATION: Xaa = Any Amino Acid

<400> SEQUENCE: 94

```

Asn Thr Glu Asn Cys Ser Tyr Glu Ser Met Gly Glu Ser Asn Asn Asp
 1           5           10           15

His Phe Gln His Phe Pro Pro Pro Pro Pro Ile Leu Glu Thr Ser Ser
20           25           30

Leu Glu Leu Ala Ser Lys Lys Pro Ser Glu Ile Gln Gln Val Asn Asn
35           40           45

Pro Glu Leu Gly Leu Ser Arg Ala Ala Leu Gln Met Gln Phe Asn Ala
50           55           60

Ala Glu Arg Glu Thr Asn Gly Val His Pro Ser Xaa Gly Val Asn Gly
65           70           75           80

Leu Ile Asn Gly Lys Ala Asn Ser Asn Lys Ser Leu Pro Thr Pro Ala
85           90           95

Val Leu Leu Ser Pro Thr Lys Glu Pro Pro Pro Leu Leu Ala Lys Pro
100          105          110

Lys

```

<210> SEQ ID NO 95
 <211> LENGTH: 22
 <212> TYPE: PRT
 <213> ORGANISM: H. sapiens

<400> SEQUENCE: 95

```

Ala Glu Arg Glu Thr Asn Gly Val His Pro Ser Arg Gly Val Asn Gly
 1           5           10           15

Leu Ile Asn Gly Lys Ala
20

```

<210> SEQ ID NO 96
 <211> LENGTH: 22

-continued

<212> TYPE: PRT
 <213> ORGANISM: mouse

<400> SEQUENCE: 96

Ala Glu Arg Glu Thr Asn Gly Val His Pro Ser His Gly Val Asn Gly
 1 5 10 15

Leu Ile Asn Gly Lys Ala
 20

<210> SEQ ID NO 97
 <211> LENGTH: 24
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 97

cacaacacag ggattctcag aaga 24

<210> SEQ ID NO 98
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 98

agctggtggtt ggaagagatt 20

<210> SEQ ID NO 99
 <211> LENGTH: 772
 <212> TYPE: PRT
 <213> ORGANISM: H. sapiens

<400> SEQUENCE: 99

Pro Glu Ser Pro Gly Gly Arg Gly Ile Lys Pro Asp Thr Cys Pro Ala
 1 5 10 15

Pro Gly Pro Arg Ser Pro Leu Gln Leu Pro Leu Ala Pro Asp Ala Glu
 20 25 30

Ser Gly Ser Ser Gly Arg Arg Pro Gly Glu Pro Arg Asp Pro Leu Lys
 35 40 45

Leu Gln Gln Leu Gln Asn Gln Ile Arg Leu Glu Gln Glu Ala Gly Ala
 50 55 60

Arg Gln Pro Pro Pro Ala Pro Arg Ser Ala Pro Pro Ser Pro Pro Phe
 65 70 75 80

Pro Pro Pro Pro Ala Phe Pro Glu Leu Ala Ala Cys Thr Pro Pro Ala
 85 90 95

Ser Pro Glu Pro Met Ser Ala Leu Ala Ser Arg Ser Ala Pro Ala Met
 100 105 110

Gln Ser Ser Gly Ser Phe Asn Tyr Ala Arg Pro Lys Gln Phe Ile Ala
 115 120 125

Ala Gln Asn Leu Gly Pro Ala Ser Gly His Gly Thr Pro Ala Ser Ser
 130 135 140

Pro Ser Ser Ser Ser Leu Pro Ser Pro Met Ser Pro Thr Pro Arg Gln
 145 150 155 160

Phe Gly Arg Ala Pro Val Pro Pro Phe Ala Gln Pro Phe Gly Ala Glu
 165 170 175

-continued

Pro	Glu	Ala	Pro	Trp	Gly	Ser	Ser	Ser	Pro	Ser	Pro	Pro	Pro	Pro	Pro	Pro
180					185					190						
Pro	Pro	Val	Phe	Ser	Pro	Thr	Ala	Ala	Phe	Pro	Val	Pro	Asp	Val	Phe	
195					200					205						
Pro	Leu	Pro	Pro	Pro	Pro	Pro	Pro	Leu	Pro	Ser	Pro	Gly	Gln	Ala	Ser	
210					215					220						
His	Cys	Ser	Ser	Pro	Ala	Thr	Arg	Phe	Gly	His	Gly	Gln	Thr	Pro	Ala	
225					230					235					240	
Ala	Phe	Leu	Ser	Ala	Leu	Leu	Pro	Ser	Gln	Pro	Pro	Pro	Ala	Ala	Val	
245					250					255						
Asn	Ala	Leu	Gly	Leu	Pro	Lys	Gly	Val	Thr	Pro	Ala	Gly	Phe	Pro	Lys	
260					265					270						
Lys	Ala	Ser	Arg	Thr	Ala	Arg	Ile	Ala	Ser	Asp	Glu	Glu	Ile	Gln	Gly	
275					280					285						
Thr	Lys	Asp	Ala	Val	Ile	Gln	Asp	Leu	Glu	Arg	Lys	Leu	Arg	Phe	Lys	
290					295					300						
Glu	Asp	Leu	Leu	Asn	Asn	Gly	Gln	Pro	Arg	Leu	Thr	Tyr	Glu	Glu	Arg	
305					310					315					320	
Met	Ala	Arg	Arg	Leu	Leu	Gly	Ala	Asp	Ser	Ala	Thr	Val	Phe	Asn	Ile	
325					330					335						
Gln	Glu	Pro	Glu	Glu	Glu	Thr	Ala	Asn	Gln	Glu	Tyr	Lys	Val	Ser	Ser	
340					345					350						
Cys	Glu	Gln	Arg	Leu	Ile	Ser	Glu	Ile	Glu	Tyr	Arg	Leu	Glu	Arg	Ser	
355					360					365						
Pro	Val	Asp	Glu	Ser	Gly	Asp	Glu	Val	Gln	Tyr	Gly	Asp	Val	Pro	Val	
370					375					380						
Glu	Asn	Gly	Met	Ala	Pro	Phe	Phe	Glu	Met	Lys	Leu	Lys	His	Tyr	Lys	
385					390					395					400	
Ile	Phe	Glu	Gly	Met	Pro	Val	Thr	Phe	Thr	Cys	Arg	Val	Ala	Gly	Asn	
405					410					415						
Pro	Lys	Pro	Lys	Ile	Tyr	Trp	Phe	Lys	Asp	Gly	Lys	Gln	Ile	Ser	Pro	
420					425					430						
Lys	Ser	Asp	His	Tyr	Thr	Ile	Gln	Arg	Asp	Leu	Asp	Gly	Thr	Cys	Ser	
435					440					445						
Leu	His	Thr	Thr	Ala	Ser	Thr	Leu	Asp	Asp	Asp	Gly	Asn	Tyr	Thr	Ile	
450					455					460						
Met	Ala	Ala	Asn	Pro	Gln	Gly	Arg	Ile	Ser	Cys	Thr	Gly	Arg	Leu	Met	
465					470					475					480	
Val	Gln	Ala	Val	Asn	Gln	Arg	Gly	Arg	Ser	Pro	Arg	Ser	Pro	Ser	Gly	
485					490					495						
His	Pro	His	Val	Arg	Arg	Pro	Arg	Ser	Arg	Ser	Arg	Asp	Ser	Gly	Asp	
500					505					510						
Glu	Asn	Glu	Pro	Ile	Gln	Glu	Arg	Phe	Phe	Arg	Pro	His	Phe	Leu	Gln	
515					520					525						
Ala	Pro	Gly	Asp	Leu	Thr	Val	Gln	Glu	Gly	Lys	Leu	Cys	Arg	Met	Asp	
530					535					540						
Cys	Lys	Val	Ser	Gly	Leu	Pro	Thr	Pro	Asp	Leu	Ser	Trp	Gln	Leu	Asp	
545					550					555					560	
Gly	Lys	Pro	Val	Arg	Pro	Asp	Ser	Ala	His	Lys	Met	Leu	Val	Arg	Glu	
565					570					575						
Asn	Gly	Val	His	Ser	Leu	Ile	Ile	Glu	Pro	Val	Thr	Ser	Arg	Asp	Ala	

-continued

580	585	590	
Gly Ile Tyr Thr Cys	Ile Ala Thr Asn Arg	Ala Gly Gln Asn Ser Phe	
595	600	605	
Ser Leu Glu Leu Val	Val Ala Ala Lys Glu	Ala His Lys Pro Pro Val	
610	615	620	
Phe Ile Glu Lys Leu	Gln Asn Thr Gly Val	Ala Asp Gly Tyr Pro Val	
625	630	635	640
Arg Leu Glu Cys Arg	Val Leu Gly Val Pro	Pro Pro Gln Ile Phe Trp	
645	650	655	
Lys Lys Glu Asn Glu	Ser Leu Thr His Ser	Thr Asp Arg Val Ser Met	
660	665	670	
His Gln Asp Asn His	Gly Tyr Ile Cys Leu	Leu Ile Gln Gly Ala Thr	
675	680	685	
Lys Glu Asp Ala Gly	Trp Tyr Thr Val Ser	Ala Lys Asn Glu Ala Gly	
690	695	700	
Ile Val Ser Cys Thr	Ala Arg Leu Asp Val	Tyr Thr Gln Trp His Gln	
705	710	715	720
Gln Ser Gln Ser Thr	Lys Pro Lys Lys Val	Arg Pro Ser Ala Ser Arg	
725	730	735	
Tyr Ala Ala Leu Ser	Asp Gln Gly Leu Asp	Ile Lys Ala Ala Phe Gln	
740	745	750	
Pro Glu Ala Asn Pro	Ser His Leu Thr Leu	Asn Thr Ala Leu Val Glu	
755	760	765	
Ser Glu Asp Leu			
770			

<210> SEQ ID NO 100
 <211> LENGTH: 1080
 <212> TYPE: DNA
 <213> ORGANISM: H. sapiens

<400> SEQUENCE: 100

```

tgagagtga atggggcagc atacttcact gacttagcgc ttggtccaga atgaactgct    60
ggccccacac ttgggcctgc taactgtgtc cgttccctg ggtgttctct gcagctggcc    120
taagtgccaa gcgatgtcca caagctgagc tcaactcctg aatacacggt cctggccggt    180
ccatctctga aggtgtcact tctcttttcc ccccagga ccctctgaag ctccagcaac    240
tccagaacca aatccgactg gagcaggagg cgggcgctcg gcagcctccg ccagcccccgc    300
gcagcgcgcc gccctcgccc cccttcccgc cgcgcgccgc ctccccgag ctgcggcct    360
gcacgcgcgc cgcgtcccgc gagcccatga gcgcgctggc ctcccgtcc gcccccgcca    420
tgcagtcctc cggtccttc aactaocgc gccccaagca gttcatcgcc gcgcagaacc    480
tcgggcccgc gtcgggccac ggcaagccgg cctccagccc cagctcgtcc agcctcccgt    540
cgcccatgtc ccgagcgcg aggcagttcg gccgcgccc cgtgcccgcc ttgcgcgagc    600
ccttcggcgc tgagcccag gcccctggg gctcctctc gccgtcgccc ccgccccgc    660
cacccccggc cttcagcccc acggtgcct tcccgggtcc cgactgttcc cactgcccgc    720
cgccaccacc gccgtcccgc agcccgggac aggcgtccca ctgctcgtcg cctgccacc    780
gcttcggcca cagccagacg cccgcggcct tctcagcgc tctgctgccc tcgagccgc    840
cgccggcggc cgtcaacgcc ctggggctgc ccaagggtgt cacccccgcg tgagtaaccg    900

```

-continued

```

ccgcggctcct ccacttcctt gccctccgc ctccggctcgc cctgggactc ccacatctcc 960
atacacgcgc tcccatcagc ctgcaacca gagcgcccca gtaacatttc acacatttct 1020
ctccgtgcga tgtaaaaatt cttaacggca atttgactca gtgattcttg cgtagccaat 1080

```

```

<210> SEQ ID NO 101
<211> LENGTH: 410
<212> TYPE: DNA
<213> ORGANISM: H. sapiens

```

```

<400> SEQUENCE: 101

```

```

agccacctct tgtactactg aaggaggaat ttatgcagac ttcttagcta ccagtgattt 60
caactctgttt taatacaaaa atttacatgt atttctttta tgatttaggt cagtgggtta 120
ccaaccccg atctaagctg gcaactagat ggaaagcccg tacgccctga cagtgtcac 180
aagatgctgg tgcgtgagaa cgggggtgcac tctctgatca tagagccagt cagtcacgt 240
gatgccggca tctacacatg tatagctacc aaccgagcag gacagaactc attcagcctg 300
gagcttgtag ttgctgtag gctcatctgt gaatccttgc tctctgacag aatgaacatc 360
agacttaca atgtaaaacta attctacatt actaaccaat acggaaaata 410

```

```

<210> SEQ ID NO 102
<211> LENGTH: 480
<212> TYPE: DNA
<213> ORGANISM: H. sapiens

```

```

<400> SEQUENCE: 102

```

```

tgcgtcagat gagagcagca caatcacctc ttctttaaca acttcacaca acacagggat 60
tctcagaaga ctctgacagt gtgaaatcac ttgtgaaact agtggcatct tcttatgttt 120
ttctctctt tccccttct tagccaacac tgaaaactgt agttacgagt caatgggaga 180
atccaacaat gaccacttcc aacactttcc acctccccct ccaatcttgg agacaagttc 240
cttgagttg gttcaaaga aacctctga gatccagcag gtgaacaacc ctgagttagg 300
cctgagcagg gcagcccttc aatgcaatt caatgctgct gagagggaaa cgaacggagt 360
ccatcccagc cgtggagtaa atggactgat taacggcaaa gctaacagta ataatctct 420
tccaacacca gctgtctgc ttccaccac taaggagcca ccacctctgc ttgcaaac 480

```

```

<210> SEQ ID NO 103
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: H. sapiens

```

```

<400> SEQUENCE: 103

```

```

Ser Gln Thr Pro Ala Ala Phe
1 5

```

```

<210> SEQ ID NO 104
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: chicken

```

```

<400> SEQUENCE: 104

```

```

Gly Gln Ser Pro Ala Ala Phe
1 5

```

```

<210> SEQ ID NO 105

```

-continued

<211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: zebra fish

<400> SEQUENCE: 105

Cys Gln Thr Pro Pro Ala Pro
 1 5

<210> SEQ ID NO 106
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: H. sapiens

<400> SEQUENCE: 106

Ser Gln Thr Ser Ala Ala Phe
 1 5

<210> SEQ ID NO 107
 <211> LENGTH: 19
 <212> TYPE: DNA
 <213> ORGANISM: H. sapiens

<400> SEQUENCE: 107

cagccagacg tccgcgcc

19

<210> SEQ ID NO 108
 <211> LENGTH: 6
 <212> TYPE: PRT
 <213> ORGANISM: H. sapiens

<400> SEQUENCE: 108

Ser Gln Thr Pro Ala Ala
 1 5

<210> SEQ ID NO 109
 <211> LENGTH: 6
 <212> TYPE: PRT
 <213> ORGANISM: H. sapiens

<400> SEQUENCE: 109

Ser Gln Thr Ser Ala Ala
 1 5

<210> SEQ ID NO 110
 <211> LENGTH: 2695
 <212> TYPE: DNA
 <213> ORGANISM: H. sapiens

<400> SEQUENCE: 110

aaatgcttta ctcttctcac cctaagggt cctgaattta gctttttatt acacagaacc 60
 cagtaaacta ataatggaa atctaaaaat tacccttct cttcttagaa aagaaatata 120
 ctgcatgatg atttgtgctt ttcgatttaa tgcacacaa tcgtttttac actgtatgtg 180
 ctttccctcc ccacagcccc agtcccagcc ccagtctcgc agttccattc cttcagggca 240
 gtttccattc taccgtgagt ttttattgct tccccctaaa acgctgcctt ggagtcgctg 300
 ttattggaca ctctttgatc tgtaaaggaa aaaaacacaa acaccggagg agaatcatcc 360
 taacaagccc acaacatadc tatgggccta aaaaacttaa attctttccc gttattagca 420
 cttctcgatc gttccaggaa agcccgattt gtagcttggc tgccccagga aggggcctgc 480

-continued

gaccagaaag atgcaaaggg cgggacaagg ttacgagtgg gtgcggggaa gcctccctaa 540
cctggggggc gcgtcccaga gctgcgagcc acgcctctc tctcccgcc gccgcgccgc 600
cggactctta ttttgaaggg cggcgggtga aggctcggag cctcctgagt caccggcgg 660
gcgaggtata aagcccgata cctgccccgc gcccggtccg cggagcccgc tgcagctccc 720
gctcgtccg gacgcggaat cgggcagcag cgggagcgg cccggagagc cgaggtaggc 780
gcggggaatc ggccctgagg ctggtggaag aaatgtgtga attaaaggga gtgcagcgtg 840
cagggtgggc cgcgagtcgg gagtgcaggg ctccggacag ggtggggggc gggaggacaa 900
gagactcgtc agcgcacttt ccaccggcc ggttccttc cgagccctct ccctcccccg 960
gcccgcggga gcacagatcc ctgaaagcgg agtcccagat aggaatgtct gcatcatctt 1020
ggcagggccc agacgggaga agatgctctg ctgccacgga cgatccctcc cgctctccta 1080
actcttctc gaacacctcg cccccccag gcccccggc cctctcttaa tcttgccac 1140
gtccctgag gggtcagacc tgggtggga gagcagagaa agcggggctg gcagagacag 1200
gggtgagggc cgggagagcc gatgacatca ttccctggca gctggcgggg agaggggtggg 1260
gacggagggg tccccccag ctgagccgag ccctcccttc ccaagccat cctcctttct 1320
ttgggatggt tcaccccacc tgggcgccat tcaaaggaag tttgcgcgta actcggggcg 1380
tcctttccc gcccggtgtc tgtgcccgc ccgcagtcgg tgtttctctt gggcctgggg 1440
gcggcgtgga gccggcggt cagtcccctc agtcccaggc cttggcagcc gctttgttca 1500
cgcggggcgc gggccgaacc cgaacctcag ctgcagcact ggcagcgcgc accggccaag 1560
cgtgggttcc cggccacggc cgggccaaca agggagggtc ccgtgagcag ccaggctggc 1620
cctccccggg cctgggtgag ctcccaaagc tggagcgcag ggctctaggc cgccccgcc 1680
cggggctgg gggcttcccg ccctttgtc tcattttaac tttgggggc agctctgtcc 1740
cgaatggtga gtggtccaga ctctagaag gggtttggtc gcttgagccc ggattgaggg 1800
ttgggacctt tgetctctg ctggggacgg aagggggcgc gctgggcagg agagggggcg 1860
tctcggtcgg gtctcctgg gctcgggcat cttcccacc tgcggagccg ggtcttgccg 1920
tgcgtgcccc gaacaggccc gggcacacc agtcccgtta ctggtcttca ggcgaaattc 1980
caggacgggt gtgcattcat caacctggga gtctctagga gccagagatg actggctcta 2040
cattttaatg cgtaatcaca gagaaactag cacgagaaa ggacagaggg accgacccaa 2100
gcttcagaga catggaataa tatgatcctg gttcataagc cttgaaaagc tcccctccgc 2160
caggttaagg aatggtgtca ccagccttg ctgtgaggat aatttcttga tttgtgttat 2220
gcaggctcct gtcaggactg tctcctaac toccactacc taaggaaaag atagccagag 2280
aattgtccag ggatttgggt ttgggttgc tctagaaaga cagcatcttt cttgtattt 2340
ttttttcac atcccccaa aacaattttt tcccccttc agacttggca gactttgatt 2400
tagataagag gtttgagtta atttgattt cccatgtct gtttagatca atggatatag 2460
gagatagttt gaagctaag taccctatg ctttgtgtg agtgaagat ttataacttg 2520
agggagcag ttaataaata gcacatcctt gaaccataaa catagagat agacaagagc 2580
tagataataa acagtttttc ccccaatccc taagctttac ttattttttt tactgattt 2640
tgtactttgt cattagaaaa aggagtgggg ttggtttccc agtgaccgca ttcatt 2695

-continued

```

<211> LENGTH: 1475
<212> TYPE: DNA
<213> ORGANISM: H. sapiens

<400> SEQUENCE: 111
cgggacaagg ttacgagtgg gtgcggggaa gcctccctaa cctgggggcc gcgtcccaga    60
gtcgcgagcc acgcctcctc tctcccgccc gccgcgccgc cggactctta tttgaaggg    120
cggcgggtga aggctcggag cctcctgagt caccgcggcg gcgaggtata aagcccgata    180
cctgccccgc gcccggtccg cggatcccgc tcgctccgga cgcggaatcg ggcagcagcg    240
ggaggcggcc cggagagccg aggtaggcgc ggggaatcgg ccctgaggct ggtggaagaa    300
atgtgtgaat taaagggagt gcagcgtgca ggggtgggccc cgagtcggga gtgcagggct    360
cgggacaggg tggggcgggg gaggacaaga gactcgtcag cgcactttcc acccggccgg    420
ttccttcccg agccctctcc ctccccggc cgcgggagc acagatccct gaaagcggag    480
ctccagatag gaatgtctgc atcatcttgg cagggcccag acgggagaag atgctctgct    540
gccacggagc atccctcccg ctctcctaac tcttctctga acacctgca cccccacggc    600
ccccgcccc tctcttaate ttggccacgc tccctgaggg gtcagacctg ggggtgggaga    660
gcagagaaa cggggctggc agagacaggg gtgagggccc ggagaggcga tgacatcatt    720
ccctggcagc tggcggggag aggggtggga cggaggggtc cccccgacct gagccgagcc    780
ctccctccc aagcccctcc tcccttcttt gggatggttc accccacctg ggcgccattc    840
aaaggaagt tgcgcgtaac tcggggcgtc ctttcccgg cgggtgctg tgcccggccc    900
gcagtcggtg tttcttctgg gcctgggggc ggcgtggagc cggcggctca gtcccctcag    960
tcccaggcct tggcagccc tttgttcacg cggggcggc gccgaacccg aacctcagct    1020
gcagcactgg cagcgcgcac cggccacgcg tgggttccc gccacggccc ggccaacaag    1080
ggagggtccc gtgagcagcc aggtggccc tccccgggcc tgggtgagct cccaaagctg    1140
gagcgcaggg ctctaggccc gcccccccc ggggtgggg gcttcccgc cgtttgtctc    1200
atthtaact ttggggcgag ctctgtccc aatggtgagt ggtccagact ctagaaggg    1260
gtttggtcgc ttgagcccg attgaggett gggacctttg ctctcttct ggggacggaa    1320
gggggcgcgc tggcagggag agggggcgtc tcggtcgggt ctccctgggc tcgggcatct    1380
tcccacctg cggagccggg tcttgcgctg cgtgcccga acaggcccgg gcacaccgag    1440
tcccgttact ggtcttcagg cgaaattcca ggacg    1475

```

```

<210> SEQ ID NO 112
<211> LENGTH: 118
<212> TYPE: DNA
<213> ORGANISM: H. sapiens

<400> SEQUENCE: 112
cgggcgaggt ataaagccc atacctgcc cgcgccgggt ccgaggagcc cgetgcagct    60
cccgtcgtc ccgacgcgg aatcgggcag cagcgggagg cggcccggag agccgagg    118

```

```

<210> SEQ ID NO 113
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

```

-continued

```

<400> SEQUENCE: 113
gcggagcccg ctgcagctcc cg                               22

<210> SEQ ID NO 114
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic probe

<400> SEQUENCE: 114
cggagcccgc tgcagctccc                               20

<210> SEQ ID NO 115
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 115
ccccgcgccc ggtccgcgga                               20

<210> SEQ ID NO 116
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic primer

<400> SEQUENCE: 116
cgcgtccgga gcgagcggga                               20

```

1. A method of detecting a pre-cancerous or cancerous cell in a biological sample, the method comprising detecting a palladin target nucleic acid comprising a nucleotide sequence alteration that is associated with cancer.

2. The method of claim 1, wherein the nucleotide sequence alteration is a nucleotide substitution, and wherein the nucleotide substitution is a C→T substitution at position 715 of a human palladin nucleic acid, a G→T substitution at position 1671 of a human palladin nucleic acid, or a C→T substitution at position 2060 of a human palladin nucleic acid.

3. The method of claim 1, wherein said detecting comprises nucleic acid-based amplification, and wherein an amplification product is generated.

4. The method of claim 3, wherein said amplification product comprises the nucleotide substitution that is associated with cancer.

5. The method of claim 1, wherein said detecting comprises hybridization with a palladin nucleic acid probe.

6. The method of claim 1, wherein the pre-cancerous or cancerous cell is a pancreatic cell, a breast cell, an axillary lymph node cell, and a squamous cell of the upper aerodigestive tract.

7. A method of identifying an individual at risk of developing cancer, the method comprising the method comprising detecting a palladin target nucleic acid comprising a nucleotide sequence alteration that is associated with cancer.

8. The method of claim 7, wherein the nucleotide sequence alteration is a nucleotide substitution, and wherein the nucle-

otide substitution is a C→T substitution at position 715 of a human palladin nucleic acid, a G→T substitution at position 1671 of a human palladin nucleic acid, or a C→T substitution at position 2060 of a human palladin nucleic acid.

9. The method of claim 8, wherein the target palladin nucleic acid is genomic DNA.

10. The method of claim 8, wherein said detecting comprises nucleic acid-based amplification, and wherein an amplification product is generated.

11. The method of claim 10, wherein said amplification product comprises the nucleotide sequence alteration that is associated with cancer.

12. The method of claim 8, wherein said detecting comprises hybridization with a palladin nucleic acid probe.

13. The method of claim 8, wherein the cancer is pancreatic cancer, breast cancer, or a head and neck cancer.

14. A method of detecting a pre-cancerous or cancerous cell in a biological sample, the method comprising detecting a mutated palladin polypeptide in the biological sample.

15. The method of claim 14, wherein the mutated palladin polypeptide comprises a Pro→Ser mutation at position 239 of a palladin polypeptide, a Trp→Cys mutation at position 557 of a palladin polypeptide, or an Arg→Cys mutation in a palladin polypeptide.

16. The method of claim 14, wherein said method is an immunological assay.

17. The method of claim 14, wherein said detecting comprises contacting the biological sample with an antibody specific for palladin, and detecting binding of the antibody to a protein in the sample.

18. The method of claim 17, wherein the antibody comprises a detectable label.

19. A method of detecting a pre-cancerous or cancerous cell in a biological sample, the method comprising detecting a palladin target nucleic acid that is abnormally expressed in a cell in the sample.

20. The method of claim 19, wherein the method comprises detecting a palladin target nucleic acid that is over-expressed in a cancerous or pre-cancerous cell.

21. The method of claim 1, wherein a detected level of palladin target nucleic acid that is at least about 2-fold higher than a control level of the palladin target nucleic acid indicates that the cell is pre-cancerous or cancerous.

22. The method of claim 19, wherein said detecting comprises hybridization with a palladin nucleic acid probe.

23. The method of claim 19, wherein the pre-cancerous or cancerous cell is a pancreatic cell.

24. A method of detecting a pre-cancerous or cancerous cell in a biological sample, the method comprising detecting an abnormal level of a palladin polypeptide in a cell in the sample.

25. The method of claim 24, wherein said method is an immunological assay.

26. The method of claim 24, wherein said detecting comprises contacting the biological sample with an antibody specific for palladin, and detecting binding of the antibody to a protein in the sample.

27. The method of claim 26, wherein the antibody comprises a detectable label.

28. The method of claim 24, wherein a detected level of palladin polypeptide that is at least about 2-fold higher than a control level of the palladin polypeptide indicates that the cell is pre-cancerous or cancerous.

29. The method of claim 24, wherein the cancerous or pre-cancerous cell is a pancreatic cell, a breast cell, an axillary lymph node cell, and a squamous cell of the upper aerodigestive tract.

30. A method of detecting cancer in an individual, the method comprising:

detecting a level of a palladin target nucleic acid in an indicator cell from the individual, wherein the indicator cell is other than a cancer cell,

wherein detection of a level of a palladin target nucleic acid that is lower than normal levels in the indicator cell indicates the presence in the individual of cancer.

31. The method of claim 30, wherein the indicator cell is a white blood cell.

32. The method of claim 30, wherein the cancer is pancreatic cancer, breast cancer, or a head and neck cancer.

33. The method of claim 30, wherein a level of a palladin target nucleic acid that is at least 2-fold lower than normal levels in the indicator cell indicates the presence in the individual of cancer.

34. A method of staging a pancreatic cancer cell, the method comprising determining the level of a palladin mRNA in the pancreatic cancer cell, wherein the level of palladin mRNA in the cell indicates a stage of neoplastic progression.

35. The method of claim 34, wherein said pancreatic cancer cell is a pancreatic ductal epithelial cell.

36. The method of claim 34, wherein the level of palladin mRNA indicates that the cell is a pancreatic intraepithelial neoplasia (PanIN) pre-cancerous cell.

37. The method of claim 34, wherein the level of palladin mRNA indicates that the cell is a pancreatic intraepithelial neoplasia (PanIN) cancerous cell.

38.-51. (canceled)

* * * * *